

Iterated Function Systems, Moments, and Transformations of Infinite Matrices

Palle E. T. Jorgensen

Keri A. Kornelson

Karen L. Shuman

Author address:

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY,
IA 52242-1419, U.S.A.

E-mail address: jorgen@math.uiowa.edu

URL: <http://www.math.uiowa.edu/~jorgen/>

DEPARTMENT OF MATHEMATICS AND STATISTICS, GRINNELL COLLEGE, GRINNELL,
IA 50112-1690, U.S.A.

Current address: Department of Mathematics, University of Oklahoma, Norman,
OK 73019-0315, U.S.A.

E-mail address: kkornelson@math.ou.edu

URL: <http://www.math.ou.edu/~kkornelson/>

DEPARTMENT OF MATHEMATICS AND STATISTICS, GRINNELL COLLEGE, GRINNELL,
IA 50112-1690, U.S.A.

E-mail address: shumank@math.grinnell.edu

URL: <http://www.math.grinnell.edu/~shumank/>

Contents

Preface	vi
Acknowledgements	ix
Chapter 1. Notation	1
1.1. Hilbert space notation	1
1.2. Unbounded operators	1
1.3. Multi-index notation	2
1.4. Moments and moment matrices	2
1.5. Computations with infinite matrices	4
1.6. Inverses of infinite matrices	7
Chapter 2. The moment problem	9
2.1. The moment problem $M = M^{(\mu)}$	9
2.2. A Parthasarathy-Kolmogorov approach to the moment problem	12
2.3. Examples	16
2.4. Historical notes	18
Chapter 3. A transformation of moment matrices: the affine case	21
3.1. Affine maps	21
3.2. IFSs and fixed points of the Hutchinson operator	23
3.3. Preserving Hankel matrix structure	27
Chapter 4. Moment matrix transformation: measurable maps	31
4.1. Encoding matrix A for τ	31
4.2. Approximation of A with finite matrices	38
Chapter 5. The Kato-Friedrichs operator	43
5.1. The quadratic form Q_M	43
5.2. The closability of Q_M	44
5.3. A factorization of the Kato-Friedrichs operator	46
5.4. Kato connection to A matrix	47
5.5. Examples	52
Chapter 6. The integral operator of a moment matrix	55
6.1. The Hilbert matrix	55
6.2. Integral operator for a measure supported on $[-1, 1]$	59
Chapter 7. Boundedness and spectral properties	65
7.1. Bounded Kato operators	65
7.2. Projection-valued measures	69
7.3. Spectrum of the Kato operator	71

7.4. Rank of measures	76
7.5. Examples	77
Chapter 8. The moment problem revisited	85
8.1. The shift operator and three incarnations of symmetry	85
8.2. Self-adjoint extensions of a shift operator	87
8.3. Self-adjoint extensions and the moment problem	89
8.4. Jacobi representations of matrices	92
8.5. The triple recursion relation and extensions to higher dimensions	98
8.6. Concrete Jacobi matrices	100
Bibliography	103

Abstract

We study the moments of equilibrium measures for iterated function systems (IFSs) and draw connections to operator theory. Our main object of study is the infinite matrix which encodes all the moment data of a Borel measure on $\mathbb{R}^d$ or $\mathbb{C}$. To encode the salient features of a given IFS into precise moment data, we establish an interdependence between IFS equilibrium measures, the encoding of the sequence of moments of these measures into operators, and a new correspondence between the IFS moments and this family of operators in Hilbert space. For a given IFS, our aim is to establish a functorial correspondence in such a way that the geometric transformations of the IFS turn into transformations of moment matrices, or rather transformations of the operators that are associated with them.

We first examine the classical existence problem for moments, culminating in a new proof of the existence of a Borel measure on $\mathbb{R}$ or $\mathbb{C}$ with a specified list of moments. Next, we consider moment problems associated with affine and non-affine IFSs. Our main goal is to determine conditions under which an intertwining relation is satisfied by the moment matrix of an equilibrium measure of an IFS. Finally, using the famous Hilbert matrix as our prototypical example, we study boundedness and spectral properties of moment matrices viewed as Kato-Friedrichs operators on weighted ℓ^2 spaces.

Received by the editor September 11, 2008.

2000 *Mathematics Subject Classification.* Primary 28A12, 34B45, 42C05, 42A82, 46E22, 47L30, 47L60, 47C10, 47S50, 54E70, 60J10, 60J20, 78M05, 81S30, 81Q15, 81T75, 82B44, 90B15.

Key words and phrases. moments, measure, iterated function system, moment matrix, Hankel matrix distribution, fractals, orthogonal polynomials, operators in Hilbert space, Hilbert matrix, positive definite functions, spectral representation, spectral measures, rank-one perturbations, spectrum, absolutely continuous, Hermitian operator, self-adjoint operator, unbounded operator, Hilbert space, deficiency indices, self-adjoint extension.

This material is based upon work partially supported by the U.S. National Science Foundation under grants DMS-0457581, DMS-0503990, and DMS-0701164, by the University of Iowa Department of Mathematics NSF VIGRE grant DMS-0602242, and by the Grinnell College Committee for Support of Faculty Scholarship. The second author was supported in part by the Woodrow Wilson Fellowship Foundation and the Andrew W. Mellon Foundation.

Preface

Moments of Borel measures μ on $\mathbb{R}^d$ have numerous uses both in analysis and in applications. For example, moments are used in the computation of orthogonal polynomials, in inverse spectral problems, in wavelets, in the analysis of fractals, in physics, and in probability theory. In this paper, we study some well-known and perhaps not-so-well-known aspects of moment theory which have been motivated by the study of both the classical literature on moments and the newer literature on iterated function systems (IFSs).

Over the last hundred years, since the time of Lebesgue, mathematicians have adopted two approaches to measures on a topological (Hausdorff) space X . In the first, measures are treated as functions on some sigma-algebra of “measurable” subsets of X . In the second, measures are realized as positive linear functionals on a suitable linear space C of functions on X . If we take C to be the compactly supported continuous functions of X , then Riesz’s theorem states that the two versions are equivalent. Starting with a measure μ on the Borel sigma algebra, integration with respect to μ yields a functional L defined on C , and conversely (by Riesz), every positive linear functional L on C takes the form of integration against some Borel measure μ . The conclusion from Riesz’s theorem asserts the existence of μ .

The moment problem is an analog, taking X to be $\mathbb{R}^d$ and replacing C with the linear space of polynomials in d variables. The issue, as before, is to construct a measure from a functional. Since the polynomials are spanned by the monomials, a functional L is then prescribed by a sequence of moments. Hence, we have the moment problem: determine a measure μ from the sequence of moments. Given the moments, one can ask about the existence of a measure having those moments, the uniqueness of the measure if one exists, and the process for constructing such a measure and determining some of its properties. We will touch on each of these topics in this Memoir.

In Chapter 1 we introduce our notation, definitions, and conventions. We have collected here for the reader’s convenience some of the tools from operator theory and harmonic analysis which we will use throughout. In Chapter 2, we review the classical moment existence problem, which asks the following question: Given a list of moments $\{m_k\}_{k \geq 0}$, does there exist a Borel measure μ such that the k^{th} moment of μ is m_k ? Alternately, we can arrange the moment sequence in a Hankel matrix in order to apply operator-theoretic techniques to this problem—a theme we carry throughout the Memoir. Using tools of Kolmogorov, Parthasarathy, and Schmidt, we provide a new approach to this old problem, which was first settled by Riesz (see [Rie23]) nearly 100 years ago. Our main result here is a new proof for the moment existence problem in both $\mathbb{R}$ and $\mathbb{C}$. We return to this same theme—the classical moment uniqueness problem—in Chapter 8. There, we describe in detail the theory of self-adjoint extensions of symmetric operators, which helps us determine precisely when a list of moments has more than one associated measure.

In Chapters 3 and 4, we explore the difficult problem of computing moments directly for an equilibrium measure arising from an iterated function system (IFS). An IFS is a finite set of contractive transformations in a metric space. A theorem of Hutchinson [Hut81] tells us that for each IFS, there exists a unique normalized equilibrium measure μ which is the solution to a fixed point problem for measures. We show that every IFS corresponds to a non-abelian system of operators and a

fixed point problem for infinite matrices. We then prove that the moment matrix $M = M^{(\mu)}$ is a solution to this matrix fixed point problem, and in turn we exploit this fixed point property to compute or approximate the moments for μ in a more general setting than the affine cases studied in [EST06].

As shown in [EST06], it is not straightforward to compute the moments for even the simplest Cantor equilibrium measures, but we *can* compute the moments of an equilibrium measure μ directly in the affine IFS case. However, we generally have no choice but to approximate the moments in non-affine examples. In particular, our results can be applied to real and complex Julia sets, in which there is much current interest. The non-affine moment approximation problem is surprisingly subtle, and as a result, we turn to operator-theoretic methods. Affine IFSs are considered in Chapter 3, while non-affine IFSs and operator theory are considered in Chapter 4. In addition, there are associated results about spectral properties of moment matrices for equilibrium measures in Chapter 7.

Infinite Hankel matrices cannot always be realized directly by operators in the ℓ^2 sequence space. We have been able to apply operator theoretic results more widely by allowing matrices to be realized as operators on a renormalized Hilbert space when necessary. We turn to this problem in Chapter 5, where we introduce the operator theoretic extensions of quadratic forms by Kato and Friedrichs [Kat80]. The quadratic form we use in this context is induced by the moment matrix $M^{(\mu)}$. Using the Kato-Friedrichs theorem, we obtain a self-adjoint operator with dense domain. This generally unbounded Kato-Friedrichs operator can be used to obtain a spectral decomposition which helps us understand the properties of the quadratic form which gave rise to the operator. Often, renormalized or weighted spaces allow us to use the Kato-Friedrichs theorem in greater generality.

We continue to use Kato-Friedrichs theory in the remaining chapters to understand the spectral properties of the moment matrix operator. In Chapter 6, we use the classical example of the Hilbert matrix and its generalizations from Widom's work [Wid66] in order to explore spectral properties of the moment matrix $M^{(\mu)}$ for general measures. In particular, we show that the moment matrix is unitarily equivalent to a certain integral operator. In Chapter 7, we further explore the spectral properties of the moment matrix and present some detailed examples. Finally, Chapter 8 uses spectral theory as a tool to reexamine the classical moment problem, this time considering not only the existence of a measure having prescribed moments, but also the uniqueness.

Readers not already familiar with the theory of moments may find some of the classical references useful. They treat both the theory and the applications of moments of measures, and they include such classics as [ST43], [Sho47], and [Akh65]. The early uses of moments in mathematics were motivated to a large degree by applications to orthogonal polynomials [Sze75]. More recent applications of orthogonal polynomials are numerical analysis, and random matrix theory, random products, and dynamics. These applications are covered in [Lan87b], [Lan87a], [Dei99], while the edited and delightful volume [Lan87a] includes additional applications to geometry, to signal processing, to probability and to statistics.

Acknowledgements

The authors thank Professor Christopher French for several helpful conversations and the proof of Proposition 3.10. In addition, one or more of the authors had helpful conversations with Professors Ken Atkinson, Dorin Dutkay, Erin Pearse, and Myung-Sin Song.

The third author is pleased to thank the orthogonal polynomial workshop students and graduate students at the University of Iowa 2007 REU for their energy and enthusiasm. She especially thanks Bobby Elam, Greg Ongie, and Mark Tucker, whose questions led the authors to the paper [EST06].

The second author is grateful to the Mathematics Department at the University of Oklahoma - Norman for their hospitality during her research leave from Grinnell College.

CHAPTER 1

Notation

In this section, we introduce some notation, conventions, and definitions needed throughout the paper.

1.1. Hilbert space notation

We use the convention that the inner product, denoted $\langle u|v\rangle$, is linear in the second position v and conjugate linear in the first position u . This choice results in fewer matrix transpositions in the type of products we will be computing and therefore yields more straightforward computations.

We will use Dirac's notation of "bras" and "kets". For vectors u and v in a Hilbert space, the inner product is denoted "bra-ket" $\langle u|v\rangle$. In contrast, the "ket-bra" notation $|u\rangle\langle v|$ denotes the rank-one operator which sends a vector x in the Hilbert space to a scalar multiple of u . Specifically, it is the operator given by $x \mapsto \langle v|x\rangle u$. The Dirac notation for this operation is

$$|u\rangle\langle v|x\rangle = \langle v|x\rangle u,$$

or also is sometimes written to preserve the order $|u\rangle\langle v|x\rangle = u\langle v|x\rangle$.

If we consider these Hilbert space operations from the point of view of matrices and Euclidean vectors, we denote a column vector by $|u\rangle$, which we call a "ket". Similarly, we denote a row vector by $\langle u|$ and call it "bra". The notation for the inner product and rank-one operators above now are consistent with the actual matrix operations being performed. Further, the algebraic manipulations with operators and vectors are done by simply merging "bras" and "kets" in the position they naturally have as we write them.

1.2. Unbounded operators

We state here some definitions and properties related to unbounded operators on a Hilbert space. For more details, see [RS80, Con90].

DEFINITION 1.1. An *operator* F which maps a Hilbert space $\mathcal{H}_1$ to another Hilbert space $\mathcal{H}_2$ is a linear map from a subspace of $\mathcal{H}_1$ (the *domain* of F) to $\mathcal{H}_2$.

It will be assumed here that the domain of F is dense in $\mathcal{H}_1$ with respect to the norm arising from the inner product on $\mathcal{H}_1$. If F is not a bounded operator, we call it an *unbounded operator*.

DEFINITION 1.2 ([Con90]). An operator F from $\mathcal{H}_1$ to $\mathcal{H}_2$ is *closed* if the graph of F , $\{(x, Fx) \mid x \in \text{dom}(F)\}$ is a closed set in the Hilbert space $\mathcal{H}_1 \times \mathcal{H}_2$ with inner product $\langle (x_1, x_2) | (y_1, y_2) \rangle_{\mathcal{H}_1 \times \mathcal{H}_2} = \langle x_1 | y_1 \rangle_{\mathcal{H}_1} + \langle x_2 | y_2 \rangle_{\mathcal{H}_2}$. If there exists a closed extension to the operator F , then F is called *closable*. In that case, there exists a smallest closed extension, which is called the *closure* of F and is denoted $\bar{F}$.

DEFINITION 1.3. Let F be an unbounded operator with dense domain $\text{dom}(F)$ from $\mathcal{H}_1$ to $\mathcal{H}_2$. Let D be the set of all $y \in \mathcal{H}_2$ such that there exists an $x \in \mathcal{H}_1$ such that

$$\langle Fz|y \rangle_{\mathcal{H}_2} = \langle z|x \rangle_{\mathcal{H}_1}$$

for all $z \in \text{dom}(F)$. We define the operator F^* on the domain D by $F^*y = x$. Note that x is uniquely determined because F is densely defined. F^* is called the *adjoint* of F .

It follows from these definitions (see [RS80]) that F is closable if and only if F^* is densely defined, and that F^* is always closed.

DEFINITION 1.4. An operator F on a Hilbert space $\mathcal{H}$ is *symmetric* if $\text{dom}(F) \subset \text{dom}(F^*)$ and $Fx = F^*x$ for all $x \in \text{dom}(F)$. F is *self-adjoint* if F is symmetric and $\text{dom}(F) = \text{dom}(F^*)$; i.e. $F^* = F$.

Often, the term *hermitian* is also used for a symmetric operator. We see that a symmetric operator F must be closable, since the domain of F^* contains the dense set $\text{dom}(F)$, and is therefore dense.

DEFINITION 1.5. A symmetric operator F is *essentially self-adjoint* if its closure $\overline{F}$ is self-adjoint.

If an operator F is bounded on its dense domain and symmetric, it is essentially self-adjoint.

1.3. Multi-index notation

In $\mathbb{R}^d$, for $d > 1$, we will need to use multi-index notation. Here, α and β denote multi-indices belonging to $\mathbb{N}_0^d$, where $\mathbb{N}_0^d$ is the Cartesian product

$$(1.1) \quad \mathbb{N}_0^d = \underbrace{\mathbb{N}_0 \times \cdots \times \mathbb{N}_0}_{d \text{ times}}.$$

Following standard conventions, the sum $\alpha + \beta$ is defined pointwise:

$$(1.2) \quad \alpha + \beta = (\alpha_i + \beta_i)_{i=1}^d.$$

Using this notation, we have the following integral expression:

$$(1.3) \quad \int_{\mathbb{R}^d} x^\alpha d\mu(x) = \int_{\mathbb{R}^d} x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_d^{\alpha_d} d\mu(x).$$

1.4. Moments and moment matrices

Let X be one of $\mathbb{R}, \mathbb{C}, \mathbb{R}^d$ with Borel measure μ . When $X = \mathbb{R}$, we define the i^{th} order moment with respect to μ to be

$$(1.4) \quad m_i = \int_{\mathbb{R}} x^i d\mu(x).$$

If the moments of all orders are finite, we will denote by $M^{(\mu)}$ an infinite matrix called the *moment matrix*. The moment matrix in the real case has entries

$$(1.5) \quad M_{i,j}^{(\mu)} = m_{i+j} = \int x^{i+j} d\mu(x).$$

Throughout this monograph, the real moment matrices will be indexed by $\mathbb{N}_0 \times \mathbb{N}_0$ – in particular, the row and column indexing both start with 0.

DEFINITION 1.6. An infinite real matrix M whose entries are indexed by $\mathbb{N}_0 \times \mathbb{N}_0$ is called a *Hankel matrix* if

$$M_{i,j} = M_{i+k,j-k} = M_{i-k,j+k}$$

for all values of k for which these entries are defined.

The moment matrix defined in Equation (1.5) is a Hankel matrix. This justifies our notation above referencing the $(i, j)^{\text{th}}$ entry of $M^{(\mu)}$ with $i + j$. The following examples will be used throughout the paper to illustrate our techniques and results.

EXAMPLE 1.7. Lebesgue measure on $[0, 1]$.

Let μ be the Lebesgue measure supported on $[0, 1]$. Then the moment matrix for μ is

$$M_{i,j}^{(\mu)} = \int_0^1 x^{i+j} dx = \frac{1}{i+j+1}.$$

This matrix is often called the *Hilbert matrix*. We will examine the properties of the Hilbert matrix in greater detail in Section 6.1. $\diamond$

EXAMPLE 1.8. Dirac point mass measure $\mu = \delta_1$.

Let μ be the Dirac probability mass δ_1 on $\mathbb{R}$. Then the moments for μ are

$$M_{i,j}^{(\mu)} = \int x^{i+j} d\mu = 1^{i+j} = 1,$$

so the moment matrix entries are all 1. In this case, we see that the moment matrix does not represent a bounded operator on ℓ^2 . This example will be studied further in Chapters 5 and 7. $\diamond$

EXAMPLE 1.9. The measure $\mu = e^{-x} dx$.

Let μ be $e^{-x} dx$ on $\mathbb{R}^+ = (0, \infty)$. Using integration by parts, a quick induction proof shows that the moments for μ are

$$M_{i,j}^{(\mu)} = (i+j)!.$$

Again, we see that the moments increase rapidly as i, j increase, so $M^{(\mu)}$ cannot be a bounded operator on ℓ^2 . We will discover more properties for this matrix in Chapters 5 and 8. $\diamond$

EXAMPLE 1.10. Measures with moments from the gamma function.

Let

$$\Gamma(k) = \int_0^\infty s^{k-1} e^{-s} ds = (k-1)!$$

when $k \in \mathbb{Z}_+$. Set $\mu = e^{-p^2 x^2} dx$. μ is a Gaussian measure with support on $\mathbb{R}$. We have the odd moments equal to zero, and the even moments are given by

$$\int_{\mathbb{R}} x^{2k} e^{-p^2 x^2} dx = \frac{\Gamma(k)}{2p^{2k+1}}$$

for $k \in \mathbb{Z}_+$. $\diamond$

If $X = \mathbb{C}$, the moments are now indexed by $\mathbb{N}_0 \times \mathbb{N}_0$ and are given by

$$(1.6) \quad m_{ij} = \int_{\mathbb{C}} \bar{z}^i z^j d\mu(z).$$

If every moment is finite, then the complex moment matrix $M^{(\mu)}$ is given by

$$(1.7) \quad M_{i,j}^{(\mu)} = m_{ij} = \int_{\mathbb{C}} \bar{z}^i z^j d\mu(z).$$

Notice that a complex moment matrix will not in general have the Hankel property that arises in real measures. We also observe that the complex moments are equivalent to the inner products of monomials in the Hilbert space $L^2(\mu)$:

$$(1.8) \quad m_{ij} = \int_{\mathbb{C}} \bar{z}^i z^j d\mu(z) = \langle z^i | z^j \rangle_{L^2(\mu)}.$$

Moments and moment matrices arise naturally in the study of orthogonal polynomials in $L^2(\mu)$.

EXAMPLE 1.11. If μ is the Lebesgue measure supported on the unit circle $\mathbb{T}$ in $\mathbb{C}$, then the moment matrix is the identity matrix:

$$M_{j,k}^{(\mu)} = \int_{\mathbb{T}} \bar{z}^j z^k d\mu = \int_0^1 e^{2\pi i(k-j)x} dx = \delta_{jk}.$$

As we noted above, the identity matrix is not Hankel.

◇

In the case where $X = \mathbb{R}^d$, we index the moments using the multi-index notation defined in Section 1.3. Given $\alpha \in \mathbb{N}_0^d$, we have the α -moment

$$(1.9) \quad m_\alpha = \int_{\mathbb{R}^d} x^\alpha d\mu(x).$$

The moment matrix for $\mathbb{R}^d$ is actually indexed by $\mathbb{N}_0^d \times \mathbb{N}_0^d$ and has entries

$$(1.10) \quad M_{\alpha,\beta}^{(\mu)} = m_{\alpha+\beta} = \int_{\mathbb{R}^d} x^{\alpha+\beta} d\mu(x).$$

When M has real entries indexed by $\mathbb{N}_0^d \times \mathbb{N}_0^d$, we will also call M a Hankel matrix if

$$M_{\alpha,\beta} = M_{\alpha-\gamma,\beta+\gamma} = M_{\alpha+\gamma,\beta-\gamma}$$

for all γ for which these entries are defined. We see from Equation (1.10) that the moment matrix for a measure on $\mathbb{R}^d$ ($d > 1$) also has the Hankel property.

1.5. Computations with infinite matrices

In a number of applications throughout this paper, we will have occasion to use infinite matrices. They will be motivated by the familiar correspondence between linear transformations and matrices from linear algebra. Given a linear transformation T between two Hilbert spaces, assumed to be infinite dimensional, then a choice of ONBs in the respective Hilbert spaces produces an infinite matrix which represents T . Moreover a number of facts from (finite-dimensional) linear algebra carry over: for example composition of two transformations (and a choice of ONBs) corresponds to multiplication of the associated two infinite matrices. Moreover by taking advantage of the orthogonality, one sees in Lemma 1.13 that the matrix multiplication is convergent.

Given an infinite matrix M , it is a separate problem to determine when there exists a linear transformation T between two Hilbert spaces and a choice of ONBs such that M represents T . This problem is only known to have solutions in special cases. We will address this further in Chapters 5 and 6. We will encounter infinite

matrices which cannot be realized directly by operators in the ℓ^2 -sequence spaces, but for which an operator representation may be found after a certain renormalization is introduced.

We use the following standard language and notation: if $G = G_{i,j}$ is an infinite matrix indexed by $\mathbb{N}_0 \times \mathbb{N}_0$, and x is an infinite vector indexed by $\mathbb{N}_0$, we will define the matrix-vector product Gx componentwise using the usual rule, provided the sums all converge.

$$(Gx)_i = \sum_{j \in \mathbb{N}_0} G_{i,j} x_j.$$

Similarly, for infinite matrices G and H , the matrix product GH is also defined componentwise using the same summation formula used for finite matrices, provided these infinite sums all converge in the appropriate sense. Specifically, the $(i,j)^{\text{th}}$ entry of the matrix GH is given by

$$(1.11) \quad (GH)_{i,j} = \sum_{k \in \mathbb{N}_0} G_{i,k} H_{k,j}.$$

As an aside, there is a parallel notion for such products which applies to matrices indexed by $\mathbb{N}_0^d \times \mathbb{N}_0^d$ and vectors indexed by $\mathbb{N}_0^d$. Given M indexed by $\mathbb{N}_0^d \times \mathbb{N}_0^d$, a vector c , and given $\alpha \in \mathbb{N}_0^d$, we have

$$(1.12) \quad (Mc)_\alpha = \sum_{\beta \in \mathbb{N}_0^d} M_{\alpha,\beta} c_\beta,$$

provided that this sum converges absolutely. We particularly need absolute convergence here so that the sum can be appropriately reordered to sum over $\mathbb{N}_0^d$.

DEFINITION 1.12. When the formal summation rules for a product of infinite matrices or a matrix-vector product yield convergent sums for each entry, we say that the matrix operations are *well defined*.

Using our matrices G and H above, if the sums $\sum_{k \in \mathbb{N}_0} G_{i,k} H_{k,j}$ converge for all $i, j \in \mathbb{N}_0$, then the matrix product GH is well defined.

If G and H are operators on a separable Hilbert space, we can determine necessary conditions on their corresponding matrix representations so that the matrix products are well defined. If we take the Hilbert space to be $\mathcal{H} = \ell^2(\mathbb{N}_0)$ and let $\{e_j : j \in \mathbb{N}_0\}$ be the standard orthonormal basis in $\ell^2(\mathbb{N}_0)$, i.e., $e_j(k) = \delta_{j,k}$ for $j, k \in \mathbb{N}_0$, then we must first assume that G and H are defined on a dense domain which includes $\{e_j\}$. This allows us to write the matrix representations of G and H . From there, we have the following result.

LEMMA 1.13. *Let G and H be linear operators densely defined on $\ell^2(\mathbb{N}_0)$ such that Ge_j , G^*e_j , and He_j are well defined and in $\ell^2(\mathbb{N}_0)$ for every element of the standard orthonormal basis $\{e_j\}_{j \in \mathbb{N}_0}$. Then $G = (G_{i,j})$ and $H = (H_{i,j})$, the infinite matrix representations of G and H respectively, are defined and the matrix product $((GH)_{i,j})$ is well defined.*

PROOF. Recall that the matrix representation of the operator G is $G_{i,j} = \langle e_i | Ge_j \rangle_2$ for $i, j \in \mathbb{N}_0$, provided Ge_j is in ℓ^2 so that the inner product is finite. Our hypotheses imply, then, that the matrix representations of G and H exist.

Given an operator G , the adjoint operator G^* satisfies $\langle G^*u | v \rangle = \langle u | Gv \rangle$ for all u, v which are in the appropriate domains and for which G^*u and Gv are in $\ell^2(\mathbb{N}_0)$.

We now check for absolute convergence of the matrix product sums from Equation (1.11):

$$\begin{aligned}
\sum_{k=0}^{\infty} |G_{ik} H_{kj}| &= \sum_{k=0}^{\infty} |\langle e_i | G e_k \rangle \langle e_k | H e_j \rangle| \\
&= \sum_{k=0}^{\infty} |\langle G^* e_i | e_k \rangle \langle e_k | H e_j \rangle| \\
&\leq \left(\sum_{k=0}^{\infty} |\langle G^* e_i | e_k \rangle|^2 \right)^{1/2} \left(\sum_{l=0}^{\infty} |\langle e_l | H e_j \rangle|^2 \right)^{1/2} \\
&= \|G^* e_i\|_2 \|H e_j\|_2 < \infty.
\end{aligned}$$

We use the Cauchy-Schwarz inequality above since we know the vectors $G^* e_i$ and $H e_j$ are in ℓ^2 . Note that if G, H are bounded operators, the sum above is bounded by $\|G^*\|_{op} \|H\|_{op}$ for any choice of $i, j \in \mathbb{N}_0$.

Once we know the sums from Equation (1.11) are absolutely convergent, they can be computed:

$$\begin{aligned}
\sum_{k=0}^{\infty} G_{i,k} H_{k,j} &= \sum_{k=0}^{\infty} \langle e_i | G e_k \rangle \langle e_k | H e_j \rangle \\
&= \sum_{k=0}^{\infty} \langle G^* e_i | e_k \rangle \langle e_k | H e_j \rangle \\
&= \langle G^* e_i | H e_j \rangle \quad \text{by Parseval's Identity} \\
&= \langle e_i | (GH) e_j \rangle \\
&= (GH)_{i,j}.
\end{aligned}$$

□

Note that the conditions in Lemma 1.13 imply that the operator G is automatically closable. It is an immediate corollary that bounded operators satisfy the hypotheses of Lemma 1.13, and thus their matrices will always have well defined products.

COROLLARY 1.14. *If G and H are bounded operators on $\ell^2(\mathbb{N}_0)$, then the product of their matrix representations is well defined.*

In some of the applications which follow - for example the cases for which both G and H are lower triangular matrices - the range of the summation index will be finite. (See, for example, the proof of Lemma 3.1.) But we will also have occasion to compute products GH for pairs of infinite matrices G and H where the range of the summation index is infinite. In those cases, we must check that the necessary sums are convergent.

There are two approaches to working with the product GH of infinite matrices G and H . One is the computational approach we have described above, and the other involves associating the matrices to operators on appropriate Hilbert spaces (see Chapters 5 and 6). For many applications, the first method is preferred. In fact, there are no known universal, or canonical, procedures for turning an infinite

matrix into an operator on a Hilbert space (see e.g., [Hal67] and [Jor06]), so often the computational approach is the only one available.

1.6. Inverses of infinite matrices

Computations in later sections will require the notion of inverse for infinite matrices. Let G be an infinite matrix indexed, as discussed in Section 1.3, by the set $\mathbb{N}_0^d \times \mathbb{N}_0^d$. We will now make precise our (admittedly abusive) use of the notation G^{-1} for an inverse. First, in order to discuss computations, the index set for rows and columns must be equipped with an order. If $d > 1$, we will use the order of the set $\mathbb{N}_0^d$ along successive finite diagonals. Our goal is to find an algorithm for the entries in the infinite matrix we denote G^{-1} . This does turn out to be possible for the infinite matrices we will be using in our analysis of moments and of transformations.

DEFINITION 1.15. Let E be an infinite matrix. We say that E is an *idempotent* if the matrix product E^2 is well defined and if $E^2 = E$.

DEFINITION 1.16. Let G, H, E_1 and E_2 be infinite matrices. Assume that the matrix products GH and HG are well defined. We say that G is a *left inverse* of H if there is an idempotent matrix E_1 such that $GH = E_1$. G is called a *right inverse* of H if there is an idempotent E_2 such that $HG = E_2$. If G is both a left and right inverse of H , G is called an *inverse* of H .

While this notion of inverses for infinite matrices is not symmetric, the following lemma does justify the use of the term “inverse”.

LEMMA 1.17. *Let G and H be infinite matrices such that both matrix products GH and HG are well defined. Suppose there is an idempotent E_1 such that $GH = E_1$ and $E_1G = GE_1 = G$. Then the infinite matrix HG is an idempotent, denoted E_2 , which satisfies $E_2H = HE_1$ and $E_2E_1 = E_2$.*

PROOF. First, we see that HG is idempotent:

$$HGHG = H(E_1)G = HG.$$

The formulas are also readily verified:

$$E_2H = HGH = HE_1$$

and

$$E_2E_1 = HGE_1 = HG = E_2.$$

□

A result of Lemma 1.17 is that if we know the matrix product $GH = E_1$ is an idempotent such that $E_1G = GE_1 = G$, then by Definition 1.16, G and H are inverses.

EXAMPLE 1.18. The following matrices arise in Example 7.17 with respect to measures which are convex combinations of Dirac masses.

We are given the matrices

$$G = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ -1 & 2 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix} \quad \text{and} \quad H_1 = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

We see that $GH_1 = E$ where E is the idempotent (in fact, projection):

$$E = GH_1 = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

Therefore, H_1 is an inverse of G . Note that this inverse is not unique. The matrix

$$H_2 = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \cdots \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \ddots \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix}$$

also satisfies $GH_2 = E$ hence is also an inverse of G .

◇

CHAPTER 2

The moment problem

In this chapter we introduce the *moment problem*, in which we describe the various positivity conditions that an infinite matrix M must satisfy in order to imply that there is a Borel measure μ on an ambient space X such that M is the moment matrix of μ , i.e. $M = M^{(\mu)}$. These conditions will be different in the cases where X is a subset of $\mathbb{R}$, $\mathbb{C}$, or $\mathbb{R}^d$ for $d > 1$. We also examine whether or not the measures found will be unique. The existence of the measure μ is discussed in Section 2.1, and a procedure by Parthasarathy using a Kolmogorov construction is used in Section 2.2 to discuss a more concrete construction of μ . The connections of the moment problem to other areas of interest to mathematicians, physicists, and engineers are mentioned in Section 2.4.

2.1. The moment problem $M = M^{(\mu)}$

Given an infinite matrix M , the moment problem addresses whether there exist measures μ such that $M = M^{(\mu)}$. There are known conditions on M which ensure that such a measure μ exists. Our presentation will be brief and we will omit the proofs of the results stated here, as they are available in the literature, albeit scattered within a variety of journal articles. Fuglede's paper [Fug83] offers a very readable survey of the literature on moments in several variables, up to 1983. The two papers [BD06, BD07] emphasize the complex case and include new results.

Our intention in this section is primarily to give the definitions of the positive semidefinite properties on the infinite matrices in the real and complex cases. We will only treat the existence part of the moment problem, but there are a variety of interesting uniqueness results in the literature as well. Here we shall only need the simplest version of uniqueness: these are the cases when the measures μ are known *a priori* to be compactly supported. In this case, uniqueness follows as a result of an application of the Stone-Weierstrass Theorem. An example of nonuniqueness is seen in Section 8.3.

We begin with the real case $\mathbb{R}^d$, for $d \geq 1$. Let $\mathcal{D}$ be the space of all infinite sequences c which are finite linear combinations of elements of the standard orthonormal basis $\{e_\alpha\}_{\alpha \in \mathbb{N}_0^d}$. Clearly $\mathcal{D}$ is a dense subspace of the Hilbert space $\ell^2(\mathbb{N}_0^d)$, and matrix-vector products are well defined on vectors in $\mathcal{D}$.

One of the ways to test whether a given sequence, or a given infinite Hankel matrix, is composed of the moments of a measure μ is to test for a positive semidefinite condition. While there are several such conditions in the literature, we isolate condition (2.1) as it summarizes a variety of features that will be essential for our point of view. Note that (2.1) entails a separate verification for every finite system

of numbers, hence it amounts to checking that all the finite truncated square matrices have positive spectrum. This in turn can be done by checking determinants of finite submatrices.

DEFINITION 2.1. A matrix M with real entries indexed by $\mathbb{N}_0^d \times \mathbb{N}_0^d$ is said to be *positive semidefinite* if

$$(2.1) \quad \langle c | M c \rangle_{\ell_2} = \sum_{\alpha \in \mathbb{N}_0^d} \sum_{\beta \in \mathbb{N}_0^d} \bar{c}_\alpha M_{\alpha, \beta} c_\beta \geq 0$$

for all $c \in \mathcal{D}$.

We say that a given infinite matrix M is a $(\mathbb{N}_0^d, \mathbb{R}^d)$ -moment matrix if there exists a positive Borel measure μ on $\mathbb{R}^d$ such that

$$(2.2) \quad M_{\alpha, \beta} = M_{\alpha, \beta}^{(\mu)} = \int_{\mathbb{R}^d} x^{\alpha+\beta} d\mu(x)$$

for all $\alpha, \beta \in \mathbb{N}_0^d$. Note that, as we observed in Section 1.4, if M is a moment matrix, it must have the Hankel property.

The solution to the existence part of the moment problem in the $\mathbb{R}^d$ case is given in a theorem due to M. Riesz [Rie23] for $d = 1$ and Haviland [Hav35, Hav36] for general d .

THEOREM 2.2 (Riesz, Haviland). *For any positive semidefinite real Hankel matrix $M = (M_{\alpha, \beta})$ indexed by $\mathbb{N}_0^d \times \mathbb{N}_0^d$, there is a positive Borel measure μ on $\mathbb{R}^d$ having moment matrix M .*

If M is an infinite matrix with complex entries, we say M is a complex moment matrix if there exists a positive Borel measure μ on $\mathbb{C}$ such that

$$(2.3) \quad M_{i, j} = M_{i, j}^{(\mu)} = \int_{\mathbb{C}} \bar{z}^i z^j d\mu(z)$$

for all $i, j \in \mathbb{N}_0$. We define a property on the complex matrix M which is stronger than the positive semidefinite property.

DEFINITION 2.3 ([BD06, BD07]). We say a complex infinite matrix M indexed by $\mathbb{N}_0 \times \mathbb{N}_0$ has *Property PDC* if given any doubly-indexed sequence $c = \{c_{i, j}\}$ having only finitely many nonzero entries, we have

$$(2.4) \quad \sum_{i, j, k, \ell \in \mathbb{N}_0} \bar{c}_{i, j} M_{i+\ell, j+k} c_{k, \ell} \geq 0.$$

Note that if we consider the special case where $j, \ell = 0$ in Definition 2.3, we get exactly the positive semidefinite condition on M given in Definition 2.1.

THEOREM 2.4 ([Fug83, BD06, BD07]). *If M is an infinite complex matrix indexed by $\mathbb{N}_0 \times \mathbb{N}_0$ which satisfies Property PDC, then there exists a positive Borel measure μ on $\mathbb{C}$ such that $M = M^{(\mu)}$.*

REMARK 2.5. The condition in Theorem 2.2 that the matrix entries in M are real numbers cannot be dropped. To see this, let $d = 1$ and take ξ be a fixed nonreal

number. Let $M = (\bar{\xi}^j \xi^k)_{j,k \in \mathbb{N}_0}$. We see that

$$\begin{aligned} \langle c | M c \rangle_{\ell^2} &= \sum_j \sum_k \bar{c}_j M_{j,k} c_k \\ &= \left(\sum_j \bar{c}_j \bar{\xi}^j \right) \left(\sum_k c_k \xi^k \right) \\ &= \left| \sum_j c_j \xi^j \right|^2 \\ &\geq 0 \end{aligned}$$

for all $c \in \mathcal{D}$. The infinite matrix M is therefore positive semidefinite. Without the condition that the components of M be real, then, we would conclude that there is a measure on $\mathbb{R}$ with M as its moment matrix. We can also verify, however, that M has Property PDC from Theorem 2.4. Let $c = \{c_{i,j}\}$ be a doubly indexed sequence in $\mathcal{D}$ (i.e. c has only finitely many nonzero entries). Then

$$\begin{aligned} \sum_{i,j,k,\ell \in \mathbb{N}_0} \bar{c}_{i,j} M_{i+\ell, j+k} c_{k,\ell} &= \sum_{i,j,k,\ell \in \mathbb{N}_0} \bar{c}_{i,j} \bar{\xi}^{i+\ell} \xi^{j+k} c_{k,\ell} \\ &= \left(\overline{\sum_{i,j} c_{i,j} \bar{\xi}^j \xi^i} \right) \left(\sum_{k,\ell} c_{k,\ell} \bar{\xi}^\ell \xi^k \right) \\ &\geq 0. \end{aligned}$$

This leads to the conclusion that there is a measure on $\mathbb{C}$ having moment matrix M . In fact, given $\xi \in \mathbb{C}$, we note that the Dirac measure δ_ξ at ξ has the matrix M as its moment matrix, since the entries of M are the evaluation of the monomials $\bar{z}^j z^k$ at ξ :

$$M_{j,k} = \int_{\mathbb{C}} \bar{z}^j z^k d\delta_\xi(z) = \bar{\xi}^j \xi^k.$$

It is shown in [Fug83] that measures with compact support are uniquely determined by their moment matrices. The Dirac measure, therefore, is the unique Borel measure on $\mathbb{C}$ having moment matrix M . If we have $\xi \in \mathbb{C} \setminus (R)$, then even though M satisfies the positive semidefinite condition from Definition 2.1, there cannot be a Borel measure μ on $\mathbb{R}$ such that $M = M^{(\mu)}$.

In some sense, what we have shown is that there is a measure, but its support is not restricted to $\mathbb{R}$. One of the classical moment questions is whether one can determine the support of a measure μ from the geometric properties of its moment matrix $M^{(\mu)}$.

The simplest moment situation arises when the measure $\mu = \delta_\xi$ is a Dirac mass. We will revisit this example frequently in the remainder of the Memoir. Observe that the infinite moment matrix for δ_ξ has rank one. We will later show that if μ is a finite convex combination of Dirac masses, then the associated moment matrix will be of finite rank.

2.2. A Parthasarathy-Kolmogorov approach to the moment problem

The traditional existence proofs of Theorems 2.2 and 2.4 are not constructive. In this section, we carefully explain how the Parthasarathy-Schmidt theorem, Theorem 2.7 below, can be applied to the moment problems from Theorems 2.2 and 2.4. Theorem 2.10 and Corollary 2.11 are restatements of the moment problem in the real and complex cases, respectively. We use Theorem 2.7 to produce the measures which appear in Theorem 2.10 and Corollary 2.11. In Corollary 2.11, the condition PDC (Definition 2.3) appears in a more natural fashion. In addition, the proof of Theorem 2.10 clearly shows why the Hankel assumption for the infinite positive definite matrix M is essential in the real case.

The use of the Kolmogorov ideas is motivated by our applications to iterated function systems (IFSs) which begin in Chapter 3. We will be interested in moments of IFS measures. A key tool in the analysis of these IFS measures will be infinite product spaces, precisely such as those that arise in the Parthasarathy-Schmidt construction.

To simplify the main idea, we carry out the details only in the real case, and only for $d = 1$. The reader will be able to generalize to the remaining real cases in $\mathbb{R}^d$, $d > 1$. We conclude with the application to the complex moment problem.

DEFINITION 2.6. Let S be a set, and let $M : S \times S \rightarrow \mathbb{C}$. We say that M is a *positive semidefinite function* if

$$\sum_{s \in S} \sum_{t \in S} \overline{c_s} M(s, t) c_t \geq 0$$

for all sequences $\{c_s\}_{s \in S} \in \mathcal{D}$. (Recall this means the sequences have only finitely many nonzero coordinates.) In the real moment problem, the matrix M is real, and we restrict to sequences $\{c_s\}_{s \in S}$ with entries in $\mathbb{R}$.

THEOREM 2.7. ([PS72, Theorem 1.2]) *Suppose S is a set, and suppose $M : S \times S \rightarrow \mathbb{C}$ is a positive semidefinite function. Then there is a Hilbert space $\mathcal{H}$ and a function $X : S \rightarrow \mathcal{H}$ such that $\mathcal{H} = \overline{\text{span}}\{X(s) : s \in S\}$ and*

$$(2.5) \quad M(s, t) = \langle X(s) | X(t) \rangle_{\mathcal{H}}.$$

Moreover, the pair $(\mathcal{H}, X)$ is unique up to unitary equivalence.

REMARK 2.8. We outline here two choices for the pair $(\mathcal{H}, X)$ in the special case where $S = \mathbb{N}_0$ and M is given by an infinite positive semidefinite matrix. The first pair is the one constructed in [PS72] using Kolmogorov's extension principle on an infinite product space. This particular choice will allow us in Theorem 2.10 to express concretely the measure satisfying the moment problem in the special case where M is Hankel.

The second pair is formed by constructing a Hilbert space from a quadratic form using the matrix M . We will be using similar completions later in the paper to work with moment matrices, so this is a natural approach. Note that Theorem 2.7 tells us that our two Hilbert spaces are isometrically isomorphic.

Proof 1. Let S be the natural numbers $\mathbb{N}_0$ and let Ω be the set of all functions from $\mathbb{N}_0$ into the one-point compactification $\overline{\mathbb{R}}$ of $\mathbb{R}$:

$$\Omega = \prod_{\mathbb{N}_0} \overline{\mathbb{R}} = (\overline{\mathbb{R}})^{\mathbb{N}_0}.$$

(Ω must be compact in order to use a Stone-Weierstrass argument in the construction.)

We now describe the Gaussian construction and its Kolmogorov consistency. For now, assume that M is positive definite in the strict sense. Let $J = \{i_1, \dots, i_p\}$ be a finite subset of $\mathbb{N}_0$. The positive definite function M gives rise to a (strictly) positive definite matrix M_J which is formed by choosing elements from rows and columns in M indexed by J . Let P_J be the Gaussian measure on $\Omega_J := \mathbb{R}^J$ with zero mean and covariance matrix M_J , such that P_J has Radon-Nikodym derivative f_J with respect to Lebesgue measure, where f_J is given by

$$(2.6) \quad f_J(\omega_J) = \frac{1}{(\sqrt{2\pi})^p \sqrt{\det M_J}} \exp\left(-\frac{1}{2} \omega_J^t M_J^{-1} \omega_J\right),$$

where we denote $\omega_J = (\omega_{i_1}, \dots, \omega_{i_p})$.

Note that Equation (2.6) uses the strict positive definite condition because the inverse M_J^{-1} is required. The density in Equation (2.6) is the standard multivariate normal density for random variables $X(i_1), \dots, X(i_p)$ with mean 0.

By Kolmogorov's extension theorem [Kol50, "Fundamental Theorem," p. 29], in order for the family of measures $\{P_J : J \text{ finite}\}$ to define a measure P on all of Ω , the measures P_J must be consistent. Suppose J and K are both finite subsets of $\mathbb{N}_0$, where $J \subset K$. In this context, *consistency* means that if g is a function on Ω_J which is extended to a function G on Ω_K depending only on the variables in J , then

$$\int_{\Omega_J} g f_J d\omega_J = \int_{\Omega_K} G f_K d\omega_K.$$

If we consider this problem in the coordinates which diagonalize the matrix M_K , we see that the variables in $K \setminus J$ will integrate to 1, and the measures P_J are indeed consistent. Note that consistency in one set of coordinates does not imply consistency in another set of coordinates.

If M is not strictly positive definite, then for any finite set $J \subset \mathbb{N}_0$, we can change coordinates to diagonalize M_J and write $\Omega_J = \ker(M_J) \oplus \ker(M_J)^\perp$. We take the measure P_J to be the Gaussian given by f_J on $\ker(M_J)^\perp$ and δ_0 , the distribution having mean 0 and variance 0, on $\ker(M_J)$. Then the covariance matrix of P_J is exactly the diagonalized M_J , and Kolmogorov consistency is maintained.

By the consistency condition, the measures P_J defined on finite subspaces of Ω can be extended to a measure P on Ω . We now take the Hilbert space in Theorem 2.7 to be $\mathcal{H} = L^2(\Omega, P)$ and the map $X : \mathbb{N}_0 \rightarrow \mathcal{H}$ to map $n \in \mathbb{N}_0$ to the projection map onto the n^{th} coordinate of the element $\omega \in \Omega$:

$$X(n)(\omega) = \omega(n).$$

Then we see that $\mathcal{H}$ is the closed linear span of the maps $\{X(n) : n \in \mathbb{N}_0\}$ and

$$\langle X(n) | X(m) \rangle_{\mathcal{H}} = \int_{\Omega} X(n)(\omega) X(m)(\omega) dP(\omega) = M(m, n).$$

We can now call M the covariance function.

To show the uniqueness statement, suppose Hilbert spaces $\mathcal{H}_1$ and $\mathcal{H}_2$ with corresponding functions X_1 and X_2 satisfy $\langle X_i(s) | X_i(t) \rangle_{\mathcal{H}_i} = M(s, t)$ and $H_i = \overline{\text{span}}\{X_i(s) : s \in S\}$ for $i = 1, 2$. For each $s \in S$, let W be the linear operator such that $W(X_1(s)) = X_2(s)$. W defined this way is an operator, since if $\sum_i c_i X_1(s_i) = 0$

then

$$\begin{aligned}
\left\| W\left(\sum_i c_i X_1(s_i)\right) \right\|_{\mathcal{H}_2}^2 &= \left\langle W\left(\sum_i c_i X_1(s_i)\right) \middle| W\left(\sum_i c_i X_1(s_i)\right) \right\rangle_{\mathcal{H}_2} \\
&= \sum_{i,j} \overline{c_i} c_j \langle X_2(s_i) | X_2(s_j) \rangle_{\mathcal{H}_2} \\
&= \sum_{i,j} \overline{c_i} c_j M(i, j) \\
&= \sum_{i,j} \overline{c_i} c_j \langle X_1(s_i) | X_1(s_j) \rangle_{\mathcal{H}_1} \\
&= \left\| \sum_i c_i X_1(s_i) \right\|_{\mathcal{H}_1}^2 = 0.
\end{aligned}$$

By linearity and density, W extends to a map from $\mathcal{H}_1$ onto $\mathcal{H}_2$, and by the association of the inner products with the matrix M , we see that W is unitary. $\square$

Proof 2. As in the first proof, let $S = \mathbb{N}_0$. Let $\mathcal{D}$ be the set of all maps from S to $\mathbb{C}$ (i.e. complex sequences) such that only finitely many coordinates are nonzero. Given the positive semidefinite function M , define the sesquilinear form

$$S(v_1, v_2) = \sum_{s \in \mathbb{N}_0} \sum_{t \in \mathbb{N}_0} \overline{v_1(s)} M(s, t) v_2(t).$$

This is clearly a sesquilinear form and therefore yields a quadratic form Q which is a seminorm

$$Q(v) = \|v\|_M^2 = S(v, v).$$

The set $\text{Null}_M = \{v \in \mathcal{D} : Q(v) = 0\}$ is a subspace of $\mathcal{D}$, so the quotient space $\mathcal{D}/\text{Null}_M$ is an inner product space. We complete this space to form a Hilbert space which we denote $\mathcal{H}_Q$ to emphasize the dependence on the quadratic form Q arising from the matrix M . We note here that this construction of a Hilbert space from a given positive semidefinite function M is unique up to unitary equivalence. We will make use of this Hilbert space completion of the quadratic form Q again in Chapters 5 and 8.

We next define the map $X : \mathbb{N}_0 \rightarrow \mathcal{H}_Q$ by

$$[X(s)](t) = \delta_s(t) = \begin{cases} 1 & s = t \\ 0 & s \neq t \end{cases}.$$

Clearly $\delta_s \in \mathcal{D} \subseteq \mathcal{H}_Q$ for all $s \in \mathbb{N}_0$, and in fact, $\mathcal{D}$ is the linear span of $\{\delta_s : s \in \mathbb{N}_0\}$. Therefore, $\mathcal{H}_Q = \overline{\text{span}}\{X(s) : s \in \mathbb{N}_0\}$. We also see that

$$\begin{aligned}
\langle X(s) | X(t) \rangle &= \sum_{u, v \in \mathbb{N}_0} \overline{\delta_s(u)} M(u, v) \delta_t(v) \\
&= M(s, t).
\end{aligned}$$

$\square$

REMARK 2.9. There are subtleties involved in these Hilbert space completions. In the pre-Hilbert space (before the completion), we will typically be working with a space $\mathcal{D}$ of all finitely supported functions on S . In other words, $\mathcal{D}$ is the space of all finite linear combinations from the orthonormal basis $\{\delta_s\}_{s \in S}$ for $\ell^2(S)$. When

the completion to a Hilbert space is done, $\mathcal{D}$ will be a dense linear subspace in the completion. Here, “dense” refers to the norm being used. In Chapter 5, for example, the norm is a weighted ℓ^2 norm.

Caution: In these alternative Hilbert space completions, say $\mathcal{H}_Q$ above, explicit representations of the vectors in $\mathcal{H}_Q$ are often not transparent. The useful geometric representations of limits of concretely given functions may be subtle and difficult. Notions of boundary constructions reside in these completions. In fact, this subtlety is quite typical when Hilbert space completions are used in mathematical physics problems. Such examples are seen in [JÓ00, Jor00], where symmetries result in separate positive definite quadratic forms, and hence two very different Hilbert space completions.

We are now ready to state our existence result for the real moment problem, using the language of Kolmogorov and Parthasarathy-Schmidt (the first proof above). Given a matrix M , our goal is to find a measure μ such that M is the moment matrix for μ . The previous theorem gives us a Hilbert space and a map $(\mathcal{H}, X)$, unique up to unitary equivalence, which we can use under the right condition (M a Hankel matrix) to determine a solution to the moment problem. This solution will not, in general, be a unique solution.

THEOREM 2.10. *Let M be an infinite matrix satisfying the positive semidefinite condition (2.1), and let $M(0, 0) = 1$. Let Ω_{Han} be the measurable subset of Ω given by*

$$(2.7) \quad \Omega_{Han} = \{\omega \in \Omega : \omega(k) = [\omega(0)]^k \text{ for all } k \in \mathbb{N}_0\}.$$

Then Kolmogorov’s extension construction yields a probability measure P_{Han} on Ω_{Han} with M as its moment matrix if and only if M satisfies the Hankel property from Definition 1.6.

PROOF. ($\Rightarrow$) Suppose a measure P_{Han} exists as stated. Theorem 2.7 defines the map X such that $X(k)(\omega) = \omega(k) = [\omega(0)]^k$ for $\omega \in \Omega_{Han}$. For the covariance function we have

$$(2.8) \quad M(j, k) = \int_{\Omega_{Han}} X(j)X(k) dP_{Han}.$$

Using Equation (2.7), we then find

$$\begin{aligned} M(j, k) &= \int_{\Omega_{Han}} [X(0)(\omega)]^j [X(0)(\omega)]^k dP_{Han}(\omega) \\ &= \int_{\mathbb{R}} x^j x^k d(P_{Han} \circ X(0)^{-1})(x) \\ &= \int_{\mathbb{R}} x^{j+k} d\mu(x) \end{aligned}$$

where we define the measure by

$$(2.9) \quad \mu = P_{Han} \circ X(0)^{-1}.$$

Since this formula shows that M is a Hankel matrix, we have proved one implication.

($\Leftarrow$) Conversely, if M is a Hankel matrix, then Kolmogorov consistency holds on the subset $\Omega_{Han} \subset \Omega$ because Ω_{Han} is measurable. By the extension principle, we get a measure space (Ω_{Han}, P_{Han}) such that the measure $\mu = P_{Han} \circ X(0)^{-1}$ on $\mathbb{R}$ is a solution to the moment problem for $M = M^{(\mu)}$. $\square$

There is an analogous result for the complex case, which we describe briefly. Let $M : \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{C}$ be a function satisfying Property PDC, i.e.,

$$(2.10) \quad \sum_{i,j,k,\ell \in \mathbb{N}_0} \bar{c}_{i,j} M_{i+\ell, j+k} c_{k,\ell} \geq 0$$

for all doubly-indexed sequences $c = \{c_{ij}\} \in \mathcal{D}$. Define an induced function $\widehat{M} : \mathbb{N}_0^2 \times \mathbb{N}_0^2 \rightarrow \mathbb{C}$ by

$$(2.11) \quad \widehat{M}((i, j), (k, l)) = M(i + l, j + k).$$

One can readily verify that $\widehat{M}$ satisfies the positive definite condition given in Theorem 2.7.

COROLLARY 2.11. *When Theorem 2.7 is applied to the function $\widehat{M}$ above, we get a pair $(X, \mathcal{H})$ where the Hilbert space may be taken to be $\mathcal{H} = L^2(\mathbb{C}, \mu)$, and the function can be given by $X(i, j) = z^i \bar{z}^j$ for all $(i, j) \in \mathbb{N}_0^2$.*

PROOF. Given these choices of $\mathcal{H}$ and X , we see that

$$\langle X(i, j) | X(k, l) \rangle_{\mathcal{H}} = \int_{\mathbb{C}} \overline{z^i \bar{z}^j} z^k \bar{z}^l d\mu(z)$$

by Theorem 2.7. Then,

$$\begin{aligned} \int_{\mathbb{C}} \overline{z^{i+l}} z^{j+k} d\mu(z) &= \langle z^{i+l} | z^{j+k} \rangle_{L^2(\mu)} \\ &= M(i + l, j + k) = \widehat{M}((i, j), (k, l)). \end{aligned}$$

This verifies our desired result and also shows that μ is a solution to the complex moment problem for the matrix $M_{ij} = M(i, j)$. $\square$

2.3. Examples

EXAMPLE 2.12. The measure $\mu = e^{-x} dx$.

We showed in Example 1.9 that the measure $e^{-x} dx$ on the positive reals has moment matrix

$$M_{i,j}^{(\mu)} = (i + j)!$$

for all $i, j \in \mathbb{N}_0$. This measure does not have compact support, but it is an example of a case where the solution to the moment problem $M = M^{(\mu)}$ is unique. We know this because the Laguerre system of orthogonal polynomials is dense in $L^2(\mu, \mathbb{R}^+)$.

Recall our earlier observation that this is also an example which illustrates that infinite Hankel matrices cannot always be realized directly by operators on ℓ^2 -sequence spaces. However we will show that an operator representation may be found after a certain renormalization is introduced. This will be an example of a general operator theoretic framework to be introduced in Chapter 5. $\diamond$

We next describe a series of examples of measures having moments involving the Catalan numbers. We will revisit these examples in Chapter 7, where we will be able to say more about the operator properties of the moment matrices.

The k^{th} Catalan number C_k is defined

$$C_k := \frac{1}{k+1} \binom{2k}{k} = \frac{(2k)!}{k!(k+1)!},$$

where the first Catalan number, C_0 , is 1. We define B_k to be

$$B_k := \binom{2k}{k}.$$

The Catalan numbers satisfy the following relation:

$$C_{k+1} = \sum_{n=0}^k C_n C_{k-n}.$$

There is an explicit formula for the generating function associated with the Catalan numbers:

$$(2.12) \quad G_{\text{Cat}}(x) = \sum_{k=0}^{\infty} C_k x^k = \frac{1 - \sqrt{1 - 4x}}{2x}.$$

The radius of convergence for $G_{\text{Cat}}(x)$ is $\frac{1}{4}$.

The generating function $G_{\text{Bin}}(x)$ associated with the B_k 's has the same radius of convergence; in fact,

$$(2.13) \quad G_{\text{Cat}}(x) = \frac{1}{x} \int_0^x G_{\text{Bin}}(y) dy,$$

and

$$(2.14) \quad G_{\text{Bin}}(x) = \frac{2}{\sqrt{1 - 4x}}.$$

The Hankel matrix $M^{\text{Cat}} = (C_{j+k})$ is positive definite because every principal submatrix has determinant 1.

EXAMPLE 2.13. Wigner's semicircle measure.

The measure $d\mu$ is given on $(-2, 2)$ by

$$d\mu(x) = \frac{\sqrt{4 - x^2}}{2\pi} dx$$

and is 0 otherwise. A simple calculation shows that

$$m_{2k} = \int_{-2}^2 x^{2k} d\mu(x) = C_k$$

and

$$m_{2k+1} = \int_{-2}^2 x^{2k+1} d\mu(x) = 0.$$

◇

EXAMPLE 2.14. The secant measure.

The measure $d\mu$ is given on $(-2, 2)$ by

$$d\mu(x) = \frac{1}{\pi \sqrt{4 - x^2}} dx$$

and is 0 otherwise. In this case,

$$m_{2k} = \int_{-2}^2 x^{2k} d\mu(x) = B_k$$

and

$$m_{2k+1} = \int_{-2}^2 x^{2k+1} d\mu(x) = 0.$$

◇

EXAMPLE 2.15. The half-secant measure.

The measure $d\mu$ is given on $(0, 2)$ by

$$d\mu(x) = \frac{2}{\pi\sqrt{4-x^2}} dx$$

and is 0 otherwise. In this example, both the even and odd moments are nonzero:

$$m_{2k} = \int_0^2 x^{2k} d\mu(x) = B_k,$$

and

$$m_{2k+1} = \int_0^2 x^{2k+1} d\mu(x) = \frac{4^{k+2}}{\pi(k+1)} \binom{2k+1}{k}^{-1}.$$

◇

2.4. Historical notes

Our positive semidefinite functions M in the form (2.1) have a history in a variety of guises under the names *positive semidefinite kernels*, *positive hermitian matrices*, *reproducing kernels*, or *positive definite functions*, among others. In these settings, a positive semidefinite function is just a function in two variables, or, equivalently, a function M defined on $S \times S$ where S is a set.

There is a broad literature covering many aspects of positive semidefinite kernels, especially in the case when S is a domain in a continuous manifold; S may even be a function space. Function spaces lead to the theory of reproducing kernels, which are also called Bergman-Shiffer-Aronszajn kernels. See [Aro50], [BS52] for a classical exposition and [BCR84], [Dut04], [DJ06] for a modern view.

If S is a group or a semigroup, and if M is a positive semidefinite kernel, we will adopt additive notation “+” for the operation in S and restrict attention to the abelian case. Of special interest are the cases when the function M has one of two forms:

- (i) $M(s, t) = F_1(s - t)$ for some function F_1 on S
- (ii) $M(s, t) = F_2(s + t)$, F_2 a function on S .

In the first case, we say that the function F_1 is *positive semidefinite* on S . Equivalently, we say that M is a *Toeplitz matrix*. There is a substantial theory of positive semidefinite functions on groups (and semigroups), see e.g., [HR79], [BCR84]; positive semidefinite functions play a central role in harmonic analysis.

In the second case (ii), M is often called a *Hankel matrix*. In the following chapters, we develop operator theoretic duality techniques to study positive semidefinite kernels in the form of infinite Hankel matrices.

- (1) We define a (generally unbounded) operator F which maps sequence spaces ℓ^2 or their weighted variants $\ell^2(w)$ into function spaces $L^2(\mu)$. This crucial operator $F : \ell^2(w) \rightarrow L^2(\mu)$ is defined on the dense subspace $\mathcal{D}$ of finite sequences in ℓ^2 and sends a sequence into the generating function (or polynomial).

- (2) Given the Hilbert spaces $\ell^2(w)$ and $L^2(\mu)$, we introduce a duality with the use of adjoint operators. If F is a given operator from $\ell^2(w)$ to $L^2(\mu)$, its adjoint operator F_w^* maps from $L^2(\mu)$ to $\ell^2(w)$.

As a byproduct of our operator analysis for the spaces $\ell^2(w)$ and $L^2(\mu)$, we get operations on the two sides which are unitarily equivalent. The unitary equivalence is critical: it allows us to derive spectral properties of measures μ from matrix operations with infinite matrices, and vice versa.

As a final historical note, we mention that because the operator F associates elements of sequence spaces with generating functions (or polynomials), we initially obtained some of our results with formal power series in the spirit of Rota's umbral calculus [RR78], [Rom84]. We then found conditions (such as renormalizing sequence spaces) which guaranteed not just formal convergence but actual convergence. In Rota's umbral calculus, the "umbra" is a space of formal power series. In this Memoir, the "umbra" consists of the Hilbert spaces $\ell^2(w)$ and $L^2(\mu)$ and the closed linear operators F and F_w^* between them. For more about the book [Rom84] and the umbral calculus's relation to other parts of mathematics, see the 2007 review by Jorgensen [Jor07].

CHAPTER 3

A transformation of moment matrices: the affine case

In this chapter we use the maps from an iterated function system to describe a corresponding transformation on moment matrices. In the case of an affine IFS, we show that the matrix transformation is a sum of triple products of infinite matrices.

Given a measurable endomorphism τ mapping a measure space (X, μ) to itself, we make use of the measure transformation

$$(3.1) \quad \mu \mapsto \mu \circ \tau^{-1}.$$

If (X, μ) and $(X, \mu \circ \tau^{-1})$ have finite moments of all orders, we can also study the transformation of moment matrices

$$(3.2) \quad M^{(\mu)} \mapsto M^{(\mu \circ \tau^{-1})}.$$

We are interested in the circumstances under which this transformation of moment matrices (3.2) can be expressed explicitly as a well-defined intertwining of the form

$$(3.3) \quad M^{(\mu \circ \tau^{-1})} = A^* M^{(\mu)} A$$

for some suitable infinite matrix A . (We are using the notation A^* here to represent the conjugate transpose of the infinite matrix A .) We then ask whether there are conditions under which A might be an operator on the Hilbert space ℓ^2 .

We begin by stating the result from [EST06] that the matrix A can be described explicitly in the case where τ is an affine map of a single variable. We give a proof of this result in Section 3.1 and then examine in Section 3.2 the connections of this result to affine iterated function systems (IFSs) in one dimension. We also prove a stronger uniqueness result for the fixed point of iterations of the moment matrix transformation arising out of a Bernoulli IFS. We examine in Section 3.3 the types of transformations that preserve infinite Hankel matrices, as a precursor to Chapter 4 in which we explore the existence of matrices A for more general measurable maps τ .

3.1. Affine maps

We consider μ a probability measure on $\mathbb{R}$ or $\mathbb{C}$. We will consider the case where τ is an affine map, noting that a finite set $\{\tau_b\}$ of affine maps on $\mathbb{R}$ or $\mathbb{C}$ can comprise an affine iterated function system (IFS). Such IFSs are used in the study of infinite convolution problems; see e.g., [Erd39, JKS07b, JKS08]. We begin with some definitions.

Recall that the moment matrix $M^{(\mu)}$ of μ is the infinite matrix defined by $M_{i,j}^{(\mu)} = \int_{\mathbb{R}} x^{i+j} d\mu(x)$ when x is a real variable, and by $M_{i,j}^{(\mu)} = \int_{\mathbb{C}} \overline{z^i} z^j d\mu(z)$ when z is a complex variable. We naturally must assume that moments of all orders exist.

Certainly moments of all orders exist when μ has compact support, but we will not always be restricted to compactly supported measures. Recall that the row and column indexing of $M^{(\mu)}$ start at row 0 and column 0.

In the real case, we recall that every moment matrix $M^{(\mu)}$ is a Hankel matrix. Our moment matrix transformation corresponding to τ in this real setting must preserve the Hankel property, then, since $M^{(\mu \circ \tau^{-1})}$ is also a moment matrix.

The following result is stated in [EST06]. Because one of our goals is to generalize the lemma, we include a proof here for completeness. This result gives a matrix A corresponding to an affine map τ on $\mathbb{C}$, so that the moment transformation is an intertwining by matrix multiplication. Note that the same matrix A works in the real case if we take τ to be an affine map of a single real variable.

LEMMA 3.1. [EST06, Proposition 1.1, p. 80] *Suppose $\tau : \mathbb{C} \rightarrow \mathbb{C}$ is an affine map (not necessarily contractive) given by*

$$\tau(z) = cz + b,$$

where $c, b \in \mathbb{C}$. Let $M^{(\mu \circ \tau^{-1})}$ be the moment matrix associated with the measure $\mu \circ \tau^{-1}$. We have

$$(3.4) \quad M^{(\mu \circ \tau^{-1})} = A^* M^{(\mu)} A,$$

where $A = (a_{i,j})$ is the upper triangular matrix with

$$(3.5) \quad a_{i,j} = \begin{cases} \binom{j}{i} c^i b^{j-i} & i \leq j \\ 0 & \text{otherwise} \end{cases}.$$

PROOF. The $(i, j)^{\text{th}}$ entry of the moment matrix $M^{(\mu \circ \tau^{-1})}$ is given by

$$(3.6) \quad \begin{aligned} M_{i,j}^{(\mu \circ \tau^{-1})} &= \int_{\mathbb{C}} \overline{z^i} z^j d(\mu \circ \tau^{-1})(z) \\ &= \int_{\mathbb{C}} \overline{(cz + b)^i} (cz + b)^j d\mu(z). \end{aligned}$$

Even though A and $M^{(\mu)}$ are infinite matrices, the sum defining the $(i, j)^{\text{th}}$ entry of the triple product $A^* M^{(\mu)} A$ is finite because A is upper triangular. With this observation, the matrix product is well defined and we can compute the $(i, j)^{\text{th}}$ entry of the product $A^* M^{(\mu)} A$ by

$$\begin{aligned} (A^* M^{(\mu)} A)_{i,j} &= \sum_{k=0}^i \sum_{\ell=0}^j A_{i,k}^* M_{k,\ell}^{(\mu)} A_{\ell,j} \\ &= \sum_{k=0}^i \sum_{\ell=0}^j \binom{i}{k} \overline{c^k b^{i-k}} \left(\int_{\mathbb{C}} \overline{z^k} z^\ell d\mu(z) \right) \binom{j}{\ell} c^\ell b^{j-\ell} \\ &= \sum_{k=0}^i \binom{i}{k} \overline{c^k b^{i-k}} \sum_{\ell=0}^j \binom{j}{\ell} c^\ell b^{j-\ell} \left(\int_{\mathbb{C}} \overline{z^k} z^\ell d\mu(z) \right). \end{aligned}$$

Taking advantage of the linearity of the integral, we compute the sums in k and ℓ to obtain

$$\begin{aligned} (A^* M^{(\mu)} A)_{i,j} &= \int_{\mathbb{C}} \sum_{k=0}^i \binom{i}{k} \overline{c^k z^k b^{i-k}} \sum_{\ell=0}^j \binom{j}{\ell} c^\ell z^\ell b^{j-\ell} d\mu(z) \\ &= \int_{\mathbb{C}} \overline{(cz + b)^i} (cz + b)^j d\mu(z). \end{aligned}$$

This gives us our desired result. $\square$

3.2. IFSs and fixed points of the Hutchinson operator

An iterated function system has an associated compact set X and a measure μ supported on X , where both X and μ arise as unique solution to a fixed-point problem described in the following paragraph. A central theme in this Memoir is that every IFS in the classical sense corresponds to a non-abelian system of operators and a version of Equations (3.7) and (3.8) in which the moment matrix $M = M^{(\mu)}$ will be a solution to an associated fixed-point problem for moment matrices. (See Proposition 3.3 and Sections 4.1 and 5.4.) One of the consequences is that we will be able to use the formula for M in a recursive computation of the moments, which is not easy to do directly for even the simplest Cantor measures.

Let I be a finite index set. An *iterated function system* (IFS) is a finite collection $\{\tau_i\}_{i \in I}$ of contractive transformations on $\mathbb{R}$ and a set of probabilities $\{p_i\}_{i \in I}$. Using Banach's fixed point theorem and the Hausdorff metric topology, it is proved in Hutchinson's paper [Hut81] that there is a unique compact subset X of $\mathbb{R}^d$, called the *attractor* of the IFS, which satisfies the equation

$$(3.7) \quad X = \bigcup_{i \in I} \tau_i(X).$$

There is also a unique measure μ supported on X which arises from Banach's theorem.

THEOREM 3.2 (Hutchinson, [Hut81]). *Given a contractive IFS $\{\tau_i\}_{i \in I}$ in $\mathbb{R}^d$ and probabilities $\{p_i\}$, there is a unique Borel probability measure μ on $\mathbb{R}^d$ satisfying*

$$(3.8) \quad \mu = \sum_{i \in I} p_i (\mu \circ \tau_i^{-1}).$$

The measure μ is called an equilibrium measure. Moreover if $p_i > 0$ for all $i \in I$, the support of μ is the unique compact attractor X for $\{\tau_i\}_{i \in I}$.

First, let us examine an IFS on the real line $\mathbb{R}$ consisting of contractive maps. We will find that the moment matrix for the equilibrium measure of the IFS also satisfies an invariance property corresponding to Equations (3.7) and (3.8).

Let I be a finite index set. Let a contractive IFS on $\mathbb{R}$ be given by the maps $\{\tau_i\}_{i \in I}$ and the nonzero probability weights $\{p_i\}_{i \in I}$. Suppose μ is the invariant Hutchinson measure associated with this IFS. Then μ is a probability measure satisfying (3.8) which is supported on the attractor set X .

We now observe, as also noted in [EST06], that the moment matrix for the equilibrium measure μ of an IFS on the real line also satisfies an invariance property $\mathcal{R}(M^{(\mu)}) = M^{(\mu)}$ under the transformation

$$(3.9) \quad \mathcal{R} : M^{(\nu)} \mapsto M^{(\sum_{i \in I} p_i (\nu \circ \tau_i^{-1}))}.$$

In other words, $M^{(\mu)}$ is a fixed point of the transformation $\mathcal{R}$. We now state a uniqueness result.

PROPOSITION 3.3. *Given a contractive IFS $\{\tau_i\}_{i \in I}$ on $\mathbb{R}$ and probability weights $\{p_i\}_{i \in I}$ with $p_i \in (0, 1)$ and $\sum_{i \in I} p_i = 1$, there exists a unique moment matrix $M^{(\mu)}$ for a probability measure μ which satisfies*

$$(3.10) \quad \mathcal{R}(M^{(\mu)}) = M^{(\mu)}.$$

This unique solution is exactly the moment matrix for the Hutchinson equilibrium measure for the IFS. Furthermore, for any Borel regular probability measure ν supported on the attractor set X of the IFS, $\mathcal{R}^n(M^{(\nu)})$ converges componentwise to $M^{(\mu)}$.

PROOF. We begin by proving the convergence of the moments. Let S be the transformation of measures supported on X given by

$$(3.11) \quad S(\nu) = \sum_{i \in I} p_i (\nu \circ \tau_i^{-1}).$$

From [Hut81], we know that $S(\nu)$ maps probability measures to probability measures, and that for any Borel regular probability measure ν supported on X , $S^n \nu \rightarrow \mu$ as $n \rightarrow \infty$, where convergence is in the metric ρ which Hutchinson calls the L -metric on the space of measures [Hut81].

To define the L -metric, first recall the space of Lipschitz functions $\text{Lip}(X, \mathbb{R})$ on a metric space X with metric d . $\text{Lip}(X, \mathbb{R})$ consists of all functions $\phi : X \rightarrow \mathbb{R}$ such that there exists a constant $C = C_\phi < \infty$ with

$$(3.12) \quad |\phi(x) - \phi(y)| \leq C_\phi d(x, y) \quad \text{for all } x, y \in X.$$

We then define the constant

$$(3.13) \quad L(\phi) = \inf\{C_\phi \mid C_\phi \text{ satisfies (3.12)}\},$$

and hence the space of functions

$$\text{Lip}_1 = \{\phi \in \text{Lip}(X, \mathbb{R}) \mid L(\phi) \leq 1\}.$$

We can now define the L -metric:

$$(3.14) \quad \rho(\mu, \nu) = \sup \left\{ \int_X \phi d\mu - \int_X \phi d\nu : \phi \in \text{Lip}_1(X, \mathbb{R}) \right\}.$$

When the measures are defined on compact spaces, the L -metric topology is equivalent to a weak topology, and therefore we have

$$\int_X f d(S^n \nu) \rightarrow \int_X f d\mu$$

for all functions $f : X \rightarrow \mathbb{R}$ which are bounded on bounded sets [Hut81].

Let $f(x) = x^{i+j}$, which is bounded on bounded sets for all choices of $i, j \in \mathbb{N}_0$. Inserting f into the convergence above yields the componentwise convergence of the moment matrix for $S^n \nu$ to the moment matrix for μ .

Given that μ satisfies the invariance property

$$S(\mu) = \mu,$$

we next wish to conclude that $\mathcal{R}(M^{(\mu)}) = M^{(\mu)}$. We defined the transformation $\mathcal{R}$ by

$$\mathcal{R}(M^{(\nu)}) = M^{(\sum_i p_i \nu \circ \tau_i^{-1})} = M^{S\nu}.$$

Since $S\mu = \mu$, it is clear that $S\mu$ and μ have the same moment matrices. Therefore $\mathcal{R}(M^{(\mu)}) = M^{(\mu)}$.

It remains to be shown that the matrix $M^{(\mu)}$ is the unique solution among moment matrices for probability measures on X to the equation $\mathcal{R}(M) = M$. We have from [Hut81] that the measure μ is the unique probability measure satisfying $S\mu = \mu$. Suppose there exists another matrix $M^{(\nu)}$ which is a moment matrix for a probability measure ν and which is a fixed point for $\mathcal{R}$. Since $\mathcal{R}(M^{(\nu)}) = M^{(S\nu)}$, the measures ν and $S\nu$ have the same moments of all orders. By Stone-Weierstrass, since the measures have compact support and agree on the moments (hence the polynomials), the measures must be the same. Since $S\nu = \nu$, we must have $\nu = \mu$. $\square$

COROLLARY 3.4. *Let $\{\tau_i\}_{i \in I}$ be an affine contractive IFS on $\mathbb{R}$, such that $\tau_i(x) = c_i x + b_i$. Let $\{p_i\}_{i \in I}$ be probability weights and let A_i be the matrix given in Lemma 3.1 which encodes τ_i for each $i \in I$. The moment matrix transformation*

$$(3.15) \quad M^{(\nu)} \mapsto \sum_{i \in I} p_i A_i^* M^{(\nu)} A_i$$

has a unique fixed point among moment matrices. Moreover, the fixed point is the moment matrix $M^{(\mu)}$ for the Hutchinson equilibrium measure of the IFS.

Given an affine IFS and its associated moment matrix transformation, we can state a uniqueness result for finite matrices which corresponds to and extends Corollary 3.4. Given any infinite matrix M , denote by M_n the $(n+1) \times (n+1)$ matrix which is the upper left truncation $[M_{ij}]_{i,j=0,1,\dots,n}$ of M . Let $\{\tau_i\}_{i=0}^k$ be a contractive affine IFS on $\mathbb{R}$, and let μ be the unique Hutchinson measure corresponding to the IFS and probability weights $\{p_i\}_{i=0}^k$. Note that we have $\tau_i(x) = c_i x + b_i$, where $c_i < 1$ for each $i = 0, 1, \dots, k$ in the most general affine IFS.

We know from Corollary 3.4 that $M^{(\mu)}$ is the unique infinite matrix which is fixed under the transformation in Equation (3.15). We now give the result that the truncated form of the transformation in Equation (3.15) has the truncated moment matrix as a fixed point, and moreover, that it is a unique fixed point among Hankel matrices. We remark here that some of the computations required for this proof are done in [EST06], but they don't use them to state this uniqueness result.

We begin by observing that, because the matrices A_i which encode affine maps are triangular (recall Lemma 3.1), a truncation of the matrix product $A_i^* M A_i$ is exactly equal to the product of the truncated matrices $(A_i)_n^* M_n (A_i)_n$.

LEMMA 3.5. *Given infinite matrices A, M such that A is upper triangular,*

$$(3.16) \quad (A^* M A)_n = A_n^* M_n A_n.$$

PROOF. This follows immediately by the properties of block matrices assuming that the associated block products are well defined. In the matrices below, B and D are upper triangular:

$$\begin{bmatrix} B^* & 0 \\ C^* & D^* \end{bmatrix} \begin{bmatrix} E & F \\ G & H \end{bmatrix} \begin{bmatrix} B & C \\ 0 & D \end{bmatrix} = \begin{bmatrix} B^* E B & * \\ * & * \end{bmatrix}.$$

$\square$

Given the affine IFS $\{\tau_i\}_{i=0}^k, \{p_i\}_{i=0}^k$ mentioned above, let A_i be the matrix given by Lemma 3.1 which encodes each affine map $\tau_i(x) = c_i x + b_i$. Let μ be the

unique Hutchinson equilibrium measure for this IFS. Using our truncation notation, define the matrix transformation $\mathcal{R}_n$ on $(n+1) \times (n+1)$ matrices M by

$$(3.17) \quad \mathcal{R}_n(M) = \sum_{i=0}^k p_i (A_i^*)_n M (A_i)_n.$$

We then have the following result.

PROPOSITION 3.6. *For each $n \in \mathbb{N}$, $M = M_n^{(\mu)}$ is the unique Hankel matrix having $M_{0,0} = 1$ which is a fixed point for $\mathcal{R}_n$.*

PROOF. We know from Corollary 3.4 and Lemma 3.5 that for each n , $M_n^{(\mu)}$ is a fixed point of $\mathcal{R}_n$. We need only to prove the uniqueness.

For $n = 1$, assume M is a 2×2 Hankel matrix with $M_{0,0} = 1$ such that $\mathcal{R}_1(M) = M$. M is of the form

$$M = \begin{bmatrix} 1 & x \\ x & y \end{bmatrix}.$$

Then,

$$\begin{aligned} \mathcal{R}_1(M) &= \sum_{i=0}^k p_i \begin{bmatrix} 1 & 0 \\ b_i & c_i \end{bmatrix} \begin{bmatrix} 1 & x \\ x & y \end{bmatrix} \begin{bmatrix} 1 & b_i \\ 0 & c_i \end{bmatrix} \\ &= \sum_{i=0}^k p_i \begin{bmatrix} 1 & b_i + c_i x \\ b_i + c_i x & b_i^2 + 2b_i c_i x + c_i^2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & x \sum p_i c_i + \sum p_i b_i \\ x \sum p_i c_i + \sum p_i b_i & y \sum p_i c_i^2 + 2x \sum p_i b_i c_i + \sum p_i b_i^2 \end{bmatrix} \\ &= \begin{bmatrix} 1 & x \\ x & y \end{bmatrix} = M. \end{aligned}$$

There is a unique solution for x and y in the equations $x = x \sum p_i c_i + \sum p_i b_i$ and $y = y \sum p_i c_i^2 + 2x \sum p_i b_i c_i + \sum p_i b_i^2$, thus $M = M_1^{(\mu)}$.

Assume next that $M_{n-1}^{(\mu)}$ is the unique $n \times n$ Hankel matrix which is a fixed point of $\mathcal{R}_{n-1}$. Take M to be an $(n+1) \times (n+1)$ Hankel matrix with $M_{0,0} = 1$ such that $\mathcal{R}_n(M) = M$. By Lemma 3.5 and our hypothesis, M is of the form

$$M = \begin{bmatrix} M_{n-1}^{(\mu)} & B \\ B^{tr} & y \end{bmatrix},$$

where B is $n \times 1$. Due to the Hankel structure of M , all but one of the entries in B are known from $M_n^{(\mu)}$:

$$B = \begin{bmatrix} m_n \\ \vdots \\ m_{2n-2} \\ x \end{bmatrix}.$$

It is straightforward to verify that, as in the $n = 1$ case, the terms in the matrix equation $S(M) = M$ yield two linear equations in x and y which have unique solutions. Therefore $M = M_n^{(\mu)}$. $\square$

The computations in Proposition 3.6 to solve for x and y give a recursive equation to compute the moments of an equilibrium measure in terms of the previous moments.

COROLLARY 3.7 ([EST06]). *Given the IFS as above with equilibrium measure μ ,*

$$M_{m,n}^{(\mu)} = \frac{1}{1 - \sum_{i=0}^k p_i c_i^2} \sum_{i=0}^k p_i \sum_{j=0, \ell=0, (j,\ell) \neq (m,n)}^{m,n} \binom{m}{j} \binom{n}{\ell} b_i^{m+n-j-\ell} c_i^{j+\ell} M_{j,\ell}^{(\mu)}$$

3.3. Preserving Hankel matrix structure

When we work with affine IFSs for a real variable x , the moment matrix of a probability measure is a Hankel matrix whose $(0,0)^{\text{th}}$ entry is 1. We will examine here the sorts of matrix transformations of the form

$$M \mapsto A^* M A$$

which preserve Hankel structure, so that in the next section we can look for moment matrix transformations corresponding to more general maps τ . Let $\mathcal{H}^{(1)}$ denote the set of positive definite Hankel matrices whose $(0,0)^{\text{th}}$ entry is 1.

We have already shown that the transformation $\mathcal{R}$ corresponding to an affine IFS given by

$$(3.18) \quad \mathcal{R}(M) = \sum_{i \in I} p_i (A_i^* M A_i) = \sum_{b \in B} p_b M^{(\mu \circ \tau_b^{-1})}$$

maps a moment matrix $M^{(\mu)}$ to another moment matrix $M^{(\sum_{i \in I} p_i \mu \circ \tau_i^{-1})}$, and therefore maps $\mathcal{H}^{(1)}$ into itself.

DEFINITION 3.8. We say that the matrix A *preserves* $\mathcal{H}^{(1)}$ if $A^* M A \in \mathcal{H}^{(1)}$ for all matrices $M \in \mathcal{H}^{(1)}$.

In the following lemma, we refer to inverses of infinite matrices and to invertible matrices. A careful definition of these concepts is given in Definition 1.16.

LEMMA 3.9. *The following matrices preserve $\mathcal{H}^{(1)}$:*

$$(3.19) \quad D(\delta) = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & \delta & 0 & 0 & \cdots \\ 0 & 0 & \delta^2 & 0 & \cdots \\ 0 & 0 & 0 & \delta^3 & \cdots \\ \vdots & \vdots & \vdots & & \ddots \end{bmatrix} \quad \text{and} \quad G(\gamma) = \begin{bmatrix} 1 & \gamma & \gamma^2 & \gamma^3 & \cdots \\ 0 & 1 & 2\gamma & 3\gamma^2 & \cdots \\ 0 & 0 & 1 & 3\gamma & \cdots \\ 0 & 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & & \ddots \end{bmatrix}.$$

The inverse of $D(\delta)$ is $D(\delta^{-1})$ ($\delta \neq 0$) and the inverse of $G(\gamma)$ is $G(-\gamma)$.

We are pleased to thank Christopher French for the proof of the following proposition. See also [Fre07], which implicitly uses Proposition 3.10 throughout, and the papers [SS06] and [Lay01].

PROPOSITION 3.10. *Suppose $A = (a_{i,j})$ is an infinite upper triangular, invertible matrix which preserves $\mathcal{H}^{(1)}$. Then either A or $-A$ is the product of matrices of the form $D(\delta)$ and $G(\gamma)$, where D and G are defined in Lemma 3.9.*

PROOF. If $A^*MA \in \mathcal{H}^{(1)}$ for all $M \in \mathcal{H}^{(1)}$, then $a_{0,0}^2 = 1$, so $a_{0,0} = \pm 1$. If $a_{0,0} = -1$, replace A with $-A$.

Denote A by

$$A = \begin{bmatrix} 1 & a_{0,1} & a_{0,2} & \cdots \\ 0 & a_{1,1} & a_{1,2} & \cdots \\ 0 & 0 & a_{2,2} & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix}.$$

Since $a_{1,1} \neq 0$, multiply A on the right by $D(1/a_{1,1})$. The upper 2×2 principal submatrix of $AD(\delta)$ is

$$\begin{bmatrix} 1 & a_{0,1}/a_{1,1} \\ 0 & 1 \end{bmatrix}.$$

Now set $\gamma = -a_{0,1}/a_{1,1}$; the matrix $AD(\delta)G(\gamma)$ now has the 2×2 identity matrix as its upper left principal submatrix.

Now set $\tilde{A} = AD(1/a_{1,1})G(-a_{0,1}/a_{1,1})$; we know that $\tilde{A}$ preserves $\mathcal{H}^{(1)}$ by Lemma 3.9. We will show that $\tilde{A}$ is the infinite identity matrix. We start with the $k = 2$ case (instead of $k = 1$) to give more intuition.

Let $\tilde{A}$ be denoted

$$(3.20) \quad \tilde{A} = \begin{bmatrix} 1 & 0 & a_0 & \cdots \\ 0 & 1 & a_1 & \cdots \\ 0 & 0 & a_2 & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix}$$

Consider the upper left 3×3 principal submatrix of the matrix $\tilde{A}^*M\tilde{A}$:

$$(3.21) \quad \begin{bmatrix} 1 & m_1 & a_0 + a_1m_1 + a_2m_2 \\ m_1 & m_2 & * \\ a_0 + a_1m_1 + a_2m_2 & * & * \end{bmatrix}.$$

Since $\tilde{A}^*M\tilde{A}$ belongs to $\mathcal{H}^{(1)}$ for all $M \in \mathcal{H}^{(1)}$, we know that the above matrix is Hankel. We can treat m_1 and m_2 as independent variables, so $a_0 = a_1 = 0$ and $a_2 = 1$. Therefore $\tilde{A}$ actually has an upper 3×3 principal submatrix which is the identity.

Continuing inductively, suppose the upper $k \times k$ principal submatrix of $\tilde{A}$ is the $k \times k$ identity matrix and $\tilde{A}$ has the form

$$(3.22) \quad \tilde{A} = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & a_0 & * & \cdots \\ 0 & 1 & 0 & \cdots & 0 & a_1 & * & \cdots \\ \vdots & & & & & \vdots & & \\ 0 & 0 & 0 & \cdots & 1 & a_{k-1} & * & \cdots \\ 0 & 0 & 0 & \cdots & 0 & a_k & * & \cdots \\ 0 & 0 & 0 & \cdots & 0 & 0 & * & \cdots \\ \vdots & \vdots & \vdots & & \vdots & \vdots & & \ddots \end{bmatrix}.$$

Multiplying $\tilde{A}^*M\tilde{A}$, we find two expressions for the $(k, 0)^{\text{th}}$ entry:

$$(3.23) \quad m_k = \sum_{i=0}^k a_i m_i.$$

Since this equation holds for all choices of $m_1, \dots, m_k$, we must have that $a_0 = \dots = a_{k-1} = 0$ and $a_k = 1$.

Therefore, $\tilde{A} = AD(1/a_{1,1})G(-a_{0,1}/a_{1,1}) = I$, so we can write

$$(3.24) \quad A = G(a_{0,1}/a_{1,1})D(a_{1,1}).$$

□

CHAPTER 4

Moment matrix transformation: measurable maps

In the previous chapter, we focused on the case of affine transformations comprising IFSs, but there is a great deal of interest in concrete applications to nonaffine examples; e.g. real and complex Julia sets. We turn our attention to nonaffine measurable maps in this chapter. We begin with an arbitrary measurable map τ on a measure space (X, μ) , where X is a subset of $\mathbb{R}^d$ for $d \geq 1$ or $\mathbb{C}$, and the moments of all orders with respect to μ and $\mu \circ \tau^{-1}$ are finite. We ask whether the transformation of moment matrices $M^{(\mu)} \mapsto M^{(\mu \circ \tau^{-1})}$ can be expressed as a matrix triple product $M^{(\mu \circ \tau^{-1})} = A^* M^{(\mu)} A$, for A an infinite matrix. In the previous section, we stated the appropriate A when τ is an affine map. We now seek to find an intertwining matrix A for more general maps τ .

We find that A , when it can be written down, need not be a triangular matrix. We therefore also will need to examine the hypotheses under which the formal matrix products we write down are actually well defined.

4.1. Encoding matrix A for τ

The following analysis will be restricted to the real one-dimensional case. We let $X \subset \mathbb{R}$ be a Borel subset and let μ be a Borel measure with moments of all orders on X . Let τ be a measurable map on X , so that $\mu \circ \tau^{-1}$ also is a measure on X with moments of all orders. We seek to find an infinite matrix A which enacts the moment matrix transformation from $M^{(\mu)}$ to $M^{(\mu \circ \tau^{-1})}$; more precisely, such that

$$(4.1) \quad M^{(\mu \circ \tau^{-1})} = A^* M^{(\mu)} A.$$

In the space $L^2(\mu)$, let $\mathcal{P}$ be the closed linear span of the monomials. (The monomials are L^2 functions since all moments are finite for μ .) The following results hinge on a careful description of the Gram-Schmidt process on the monomials, since the monomials could possibly have linear dependence relations among them in $L^2(\mu)$ (See Example 4.1 below). Let $\{v_j\}_{j \in \mathbb{N}_0}$ be the set of monomial functions, but if x^j is dependent on $1, x, x^2, \dots, x^{j-1}$, we remove x^j from the collection. In other words, v_j might not be x^j , but it is a monomial function x^k for some $k \geq j$, and there are no finite dependence relations among the set $\{v_j\}_{j \in \mathbb{N}_0}$.

EXAMPLE 4.1. Bernoulli IFSs which yield measures with linearly dependent monomials.

Consider the affine IFS on $\mathbb{R}$ of the form

$$\tau_0(x) = \lambda x, \quad \tau_1(x) = \lambda(x + 2).$$

The properties of the Hutchinson equilibrium measure depend on the choice of the parameter λ . If $\lambda < \frac{1}{2}$, the measure is supported on a generalized Cantor set, while when $\lambda \geq \frac{1}{2}$ the measure is supported on an interval in the real line.

We can transform these into measures supported on subsets of the unit circle $\mathbb{T} \subset \mathbb{C}$ via the map $x \mapsto e^{2\pi i x}$. By [JP98], when $\lambda = \frac{1}{4}$, there is an orthonormal basis of monomials

$$\{z^k : k = 0, 1, 4, 5, 16, 17, 20, 21, \dots\}$$

for the corresponding measure contained in $\mathbb{T}$. The collection of all the monomials, however, does not have finite linear dependence relations, as described in Remark 4.2. On the other hand, if $\lambda = \frac{1}{3}$, the corresponding circle measure does have finite linear dependence relations among the monomials. These cases also arise when $\lambda > \frac{1}{2}$ [JKS08]. $\diamond$

REMARK 4.2. A central theme in our study of moments in Chapters 3 and 4 is how the study of moments relates to the self-similarity which characterizes equilibrium measures for IFSs. Because we use operator-theoretic methods to study moment matrices, we encounter spectral information along the way. Historically (see [Akh65]) the approach to spectral theory of moments in $\mathbb{R}$ went as follows:

- (1) Start with the monomials $\{x^k\}_{k \in \mathbb{N}_0}$ viewed as a dense subset of (a subspace of) $L^2(\mu)$.
- (2) Apply Gram-Schmidt to obtain the associated orthonormal polynomial basis $\{p_k\}_{k \in \mathbb{N}_0}$.
- (3) The tri-diagonal matrix J representing multiplication by x in the ONB $\{p_k\}_{k \in \mathbb{N}_0}$ (see Chapter 8) gives rise to spectral information.

However, the classical approach does not take into account the self-similar properties that μ may have. Following [JP98] and [Jor06], we can seek instead to encode the IFS structure directly into the analysis of moments. A case in point is the Cantor system mentioned above with scaling by $\frac{1}{4}$ and two subdivisions (a measure μ with scaling dimension $\frac{1}{2}$). It was shown in [JP98] that it is better to realize μ as a complex measure which is supported on the circle in the complex plane. The monomials we are then led to study are the complex monomials $\{z^k\}_{k \in \mathbb{N}_0}$ in $L^2(\mu)$.

This set of all the monomials has no finite linear dependence relations. To see this, we assume a polynomial $p(z) = \sum_{j=0}^n a_j z^j$ supported on the circle is zero in $L^2(\mu)$. Then for almost every point z in the support of μ , the polynomial is zero. Since this measure has no atoms (see [DJ07],[JKS07a]), the complement of a set of measure zero must be infinite. We conclude that since p has infinitely many zeros, it is the identically zero polynomial and hence $a_0 = a_1 = \dots = a_n = 0$.

The standard application of Gram-Schmidt on the monomials would produce an ONB of polynomials, but it would be different from an ONB of monomials such as $\{z^k | k = 0, 1, 4, 5, 16, 17, 20, 21, \dots\}$. Moreover, this approach misses the IFS scaling property of μ .

Returning to our quest to encode the map τ via a matrix A , we perform the Gram-Schmidt process on the finitely linearly independent monomials $\{v_j\}_{j \in \mathbb{N}_0}$ to construct an orthonormal basis $\{p_k\}_{k \in \mathbb{N}_0}$ of polynomials for $\mathcal{P}$. Part of the definition of the Gram-Schmidt process gives that each polynomial p_k is in the span of $\{v_0, v_1, \dots, v_k\}$, and hence is orthogonal to $sp\{v_0, \dots, v_{k-1}\} = sp\{p_0, \dots, p_{k-1}\}$.

We can write the lower triangular matrix G which enacts Gram-Schmidt as follows:

$$(4.2) \quad \sum_{i=0}^k G_{k,i} v_i = p_k.$$

Also, assume that μ is a probability measure, so p_0 is the constant function 1, i.e. $p_0(x) \equiv 1$.

In the cases where some of the monomials have been left out of the sequence $\{v_j\}$, we define a moment matrix $N^{(\mu)}$ for μ to be

$$(4.3) \quad N_{j,k}^{(\mu)} = \langle v_j | v_k \rangle_{L^2(\mu)}.$$

This adjusted moment matrix $N^{(\mu)}$ will be symmetric in the real cases but will not have the Hankel property. The moments contained in $N^{(\mu)}$ are total in $\mathcal{P}$.

We place the condition on the map τ that the powers τ^j are in the space $\mathcal{P}$ for all $j \in \mathbb{N}_0$. From here, we define the following transformations on $\mathcal{P}$:

$$(4.4) \quad Rp_k = v_k$$

$$(4.5) \quad Tp_k = v_k \circ \tau.$$

R and T are well-defined operators on $\mathcal{P}$ since they are defined on an orthonormal basis. They might be unbounded, but their domains do contain the ONB elements $\{p_k\}_{k \in \mathbb{N}_0}$ by definition. Therefore, we can express R and T in matrix form with respect to $\{p_k\}_{k \in \mathbb{N}_0}$. With a slight abuse of notation, we will also refer to these matrices as R and T respectively:

$$(4.6) \quad R_{j,k} = \langle p_j | Rp_k \rangle_{L^2(\mu)} = \langle p_j | v_k \rangle_{L^2(\mu)}$$

$$(4.7) \quad T_{j,k} = \langle p_j | Tp_k \rangle_{L^2(\mu)} = \langle p_j | v_k \circ \tau \rangle_{L^2(\mu)}.$$

Observe that the matrix for R is upper triangular, since each p_j is orthogonal to $sp\{v_0, \dots, v_{j-1}\} = sp\{p_0, \dots, p_{j-1}\}$.

LEMMA 4.3. *The matrix R is invertible in the sense of Definition 1.16, and the matrix of R^{-1} is exactly the transpose of the matrix which enacts the Gram-Schmidt process on the monomials, i.e.*

$$\sum_{i=0}^k R_{i,k}^{-1} v_i = p_k.$$

PROOF. We compute the matrix product RG^{tr} . Note that the triangular structure of G^{tr} gives a finite sum, so the product is well-defined:

$$\begin{aligned}
(RG^{tr})_{i,j} &= \sum_{k=0}^{\infty} R_{i,k} G_{j,k} \\
&= \sum_{k=0}^{\infty} \langle p_j | v_k \rangle_{L^2(\mu)} G_{j,k} \\
&= \left\langle p_j \left| \sum_{k=0}^j G_{j,k} v_k \right. \right\rangle_{L^2(\mu)} \\
&= \langle p_i | p_j \rangle \\
&= \delta_{i,j}.
\end{aligned}$$

We need to show that this composition has a dense domain. One can readily verify that when a monomial v_j is expressed as a column vector with respect to $\{p_k\}_{k \in \mathbb{N}_0}$, that column vector only has nonzero entries $\langle p_k | v_j \rangle$ for $k \leq j$. Moreover, by the upper triangular structure of G^{tr} , the matrix-vector product $G^{tr} v_j$ must be in $\mathcal{D}$. These are in the domain of the matrix for R , so we can now conclude that the monomials v_k are all in the domain of the operator RG^{tr} . Therefore, RG^{tr} has dense domain and on that domain, RG^{tr} is the identity. By Lemma 1.17, G^{tr} and R are inverses of each other, and we write $G^{tr} = R^{-1}$. $\square$

Next, we will discuss the adjoints R^* and T^* . These can be written down as matrices, but it is not always true that their domains are dense in $\mathcal{P}$. We will show that there exists a *renormalization* of $L^2(\mu)$ such that both R^* and T^* have dense domains in $\mathcal{P}$. This is equivalent (see [Con90, Proposition 1.6, Chapter 10]) to saying R and T are closable operators.

Given a set of nonzero weights $w = \{w_i\}_{i \in \mathbb{N}_0}$, define the space $\mathcal{P}_w$ to be the set of all measurable functions which are in the span of the monomials with respect to the weighted norm

$$\|f\|_{\mathcal{P}_w}^2 = \sum_{i=0}^{\infty} w_i |\langle f | p_i \rangle_{L^2(\mu)}|^2.$$

To be precise, we see that $\int f p_i d\mu$ must be finite for all $i \in \mathbb{N}_0$, but depending on the weights, observe that $\mathcal{P}_w$ could include functions which are not in $L^2(\mu)$. The inner product, then, on $\mathcal{P}_w$ is given by

$$\langle f | g \rangle_{\mathcal{P}_w} = \sum_k w_k \langle f | p_k \rangle_{L^2(\mu)} \langle p_k | g \rangle_{L^2(\mu)}.$$

If we consider the operator R as defined above, but as a map from $\mathcal{P}_w$ to $\mathcal{P}$, then the adjoint $R_w^* : \mathcal{P} \rightarrow \mathcal{P}_w$ is given by

$$(4.8) \quad R_w^* g = \sum_k \frac{1}{w_k} \langle v_k | g \rangle_{L^2(\mu)} p_k$$

on every g in the domain of R_w^* (i.e. every g such that $R_w^* g \in \mathcal{P}_w$.) Observe that the weights change the adjoint, so we denote it by R_w^* . To verify Equation (4.8), we compute for $f \in \mathcal{P}_w \cap \text{Dom}(R)$ and $g \in \mathcal{P}$;

$$Rf = \sum_{k=0}^{\infty} \langle p_k | f \rangle_{L^2(\mu)} v_k,$$

which gives

$$\langle g | Rf \rangle_{L^2(\mu)} = \left\langle g \left| \sum_{k=0}^{\infty} \langle p_k | f \rangle_{L^2(\mu)} v_k \right. \right\rangle_{L^2(\mu)} = \sum_{k=0}^{\infty} \langle p_k | f \rangle_{L^2(\mu)} \langle g | v_k \rangle_{L^2(\mu)}.$$

We then compute

$$\begin{aligned} \left\langle \sum_k \frac{1}{w_k} \langle v_k | g \rangle_{L^2(\mu)} p_k \middle| f \right\rangle_{\mathcal{P}_w} &= \sum_j w_j \left\langle \sum_k \frac{1}{w_k} \langle v_k | g \rangle_{L^2(\mu)} p_k \middle| p_j \right\rangle_{L^2(\mu)} \langle p_j | f \rangle_{L^2(\mu)} \\ &= \sum_k \langle g | v_k \rangle_{L^2(\mu)} \langle p_k | f \rangle_{L^2(\mu)}. \end{aligned}$$

The equality of these two expressions verifies our definition of R_w^* , at least for the dense set of finite linear combinations of the orthogonal polynomials, hence holds for all f in the domain of R .

In the next lemma, we produce a weighted norm on the space $\mathcal{P}$ using weights $\{w_k\}$ which guarantee that R will be closable in the weighted space $\mathcal{P}_w$.

LEMMA 4.4. *Given the measure space (X, μ) as above having finite moments of all orders and given the operator R defined in Equation (4.4), there exist weights $\{w_i\}_{i \in \mathbb{N}_0}$ such that $R : \mathcal{P}_w \rightarrow \mathcal{P}$ is closable.*

PROOF. Observe that $R_w^* g \in \mathcal{P}_w$ precisely when

$$(4.9) \quad \sum_k w_k |\langle p_k | R_w^* g \rangle_{L^2(\mu)}|^2 < \infty.$$

Substituting in the definition of R_w^* shows that Equation (4.9) is true if and only if

$$(4.10) \quad \sum_k \frac{1}{w_k} \left| \langle v_k | g \rangle_{L^2(\mu)} \right|^2 < \infty.$$

Since μ has finite 0th moment, i.e. is a finite measure, we have $L^\infty(\mu) \subseteq L^2(\mu)$. Set $\mathcal{M}_k := \int |v_k| d\mu(x)$, and suppose $f \in L^\infty(\mu)$. Thenf

$$(4.11) \quad \left| \langle v_k | f \rangle_{L^2(\mu)} \right| \leq \int_{\mathbb{R}} |v_k f| d\mu \leq \|f\|_{L^\infty(\mu)} \mathcal{M}_k.$$

Therefore, if $\{\mathcal{M}_k^2/w_k\} \in \ell^1$, then

$$(4.12) \quad L^\infty(\mu) \subset \text{dom}(R^*).$$

We thus explicitly define a choice of weights $w = \{w_k\}_{k \in \mathbb{N}_0}$ by

$$(4.13) \quad w_k := (1 + k^2) \mathcal{M}_k^2.$$

Since $L^\infty(\mu)$ is dense in $L^2(\mu)$, we now know that $\text{dom}(R_w^*)$ is dense in $\mathcal{P}$, where $w = \{w_k\}_{k \in \mathbb{N}_0}$ is defined in (4.13). Therefore $R : \mathcal{P}_w \rightarrow \mathcal{P}$ is closable. $\square$

The space $\mathcal{P}_w$ has the weighted orthogonal polynomials $\left\{ \frac{p_k}{\sqrt{w_k}} \right\}_{k \in \mathbb{N}_0}$ as an orthonormal basis. We will denote these vectors $\{p_k^w\}_{k \in \mathbb{N}_0}$. We next observe the following property of the matrix $R_w^* R$ written in terms of the orthonormal basis $\{p_k^w\}_{k \in \mathbb{N}_0}$:

$$\begin{aligned}
(4.14) \quad R_w^* R_{i,j} &= \langle p_i^w | R_w^* R p_j^w \rangle_{\mathcal{P}_w} \\
&= \langle R p_i^w | R p_j^w \rangle_{L^2(\mu)} \\
&= \frac{\langle v_i | v_j \rangle_{L^2(\mu)}}{\sqrt{w_i w_j}} \\
&= \frac{1}{\sqrt{w_i w_j}} N_{i,j}^{(\mu)}
\end{aligned}$$

So $R_w^* R$ is a self-adjoint operator and is exactly a weighted version of the moment matrix.

Next, we recall our definition (Equation (4.5)) for the operator T which maps p_i to $v_i \circ \tau$ for each $i \in \mathbb{N}_0$. By our hypothesis on τ , we know T is an operator on $\mathcal{P}$. We wish T to also be closable, so that T^* is densely defined. As we discovered for R , this may require weights. Since the adjoint depends on the weights, we will denote it T_w^* .

A function f is in the domain of T_w^* with respect to weights $w = \{w_k\}_{k \in \mathbb{N}_0}$ if $T_w^* f$ is in $\mathcal{P}_w$. The same computations used for R_w^* , replacing each v_k with $v_k \circ \tau$, show

$$(4.15) \quad T_w^* g = \sum_{k=0}^{\infty} \frac{1}{w_k} \langle v_k \circ \tau | g \rangle_{L^2(\mu)}.$$

We also find that $g \in \text{dom}(T_w^*)$ if and only if

$$(4.16) \quad \sum_{k=0}^{\infty} \frac{1}{w_k} \left| \langle v_k \circ \tau | g \rangle_{L^2(\mu)} \right|^2 < \infty.$$

The condition on τ under which there are weights such that T_w^* is densely defined on $\mathcal{P}_w$ is given below.

LEMMA 4.5. *Let $\mathcal{M}'_k = \int_X |v_k \circ \tau| d\mu = \int |v_k| d(\mu \circ \tau^{-1})$. If $\mathcal{M}'_k < \infty$ for all $k \in \mathbb{N}_0$, we define the weights $w = \{w_k\}_{k \in \mathbb{N}_0}$ by*

$$w_k = (\mathcal{M}'_k)^2 (1 + k^2),$$

for which T_w^ is densely defined from $\mathcal{P}$ to $\mathcal{P}_w$.*

PROOF. Repeating the computation in Lemma 4.4, we find that $f \in L^\infty(\mu)$ is in the domain of T_w^* if and only if

$$\sum_k \frac{1}{w_k} (\mathcal{M}'_k)^2 \|f\|_\infty < \infty,$$

which holds for the given weights. □

This then gives us an expression for the matrix of the self-adjoint operator $T_w^* T$ with respect to our orthonormal basis of polynomials:

$$\begin{aligned}
(T_w^* T)_{i,j} &= \langle p_i^w | T_w^* T p_j^w \rangle_{\mathcal{P}_w} \\
&= \langle T p_i^w | T p_j^w \rangle_{L^2(\mu)} \\
&= \frac{\langle v_i \circ \tau | v_j \circ \tau \rangle_{L^2(\mu)}}{\sqrt{w_i w_j}} \\
(4.17) \qquad &= \frac{1}{\sqrt{w_i w_j}} N_{i,j}^{\mu \circ \tau^{-1}}.
\end{aligned}$$

We next define a matrix A that will give coefficients of the functions $v_k \circ \tau$ expanded in terms of the monomials $\{v_j\}_{j \in \mathbb{N}_0}$:

$$(4.18) \qquad v_k \circ \tau = \sum_{j=0}^{\infty} A_{jk} v_j.$$

The entries of A exist since we assumed that each $v_k \circ \tau$ is an element of $\mathcal{P}$ and therefore has an L^2 -convergent expansion in the monomials. We may or may not, however, be able to compute entries of the matrix A directly from this definition.

EXAMPLE 4.6. A nonaffine map: $\tau(x) = x^2 + b$

This is perhaps the simplest example on a nonaffine transformation. The matrix A which encodes $\tau(x) = x^2 + b$ can be computed from the powers of τ to satisfy Equation (4.18):

$$A = \begin{bmatrix} 1 & b & b^2 & b^3 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 2b & 3b^2 & \dots \\ 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & 3b & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

◇

We wish to think of A as the matrix representation for an operator on the weighted space $\mathcal{P}_w$. With an abuse of notation, we will also refer to this operator as A . In general, it is not even certain that the operator A has dense domain. Throughout the remainder of this section, however, we will restrict our attention to the cases in which the domain of A contains the monomials, and hence the orthogonal polynomials.

PROPOSITION 4.7. *Given A as in Equation (4.18) and such that A is the matrix representation of an operator (also denoted A) on $L^2(\mu)$ with respect to the ONB $\{p_k\}_{k \in \mathbb{N}_0}$, if the operator composition RA is densely defined, then*

$$RA = T$$

on the domain of RA , i.e. RA is a restriction of T .

PROOF. Suppose $f \in \text{dom}(R)$. Then by Parseval we have

$$\begin{aligned} Rf &= \sum_{j=0}^{\infty} \langle p_j | f \rangle_{L^2(\mu)} R p_j \\ &= \sum_{j=0}^{\infty} \langle p_j | f \rangle_{L^2(\mu)} v_j, \end{aligned}$$

with convergence in the L^2 sense.

We have assumed that, as an operator, A has the monomial functions $\{v_k\}_{k \in \mathbb{N}_0}$ and the polynomials $\{p_k\}_{k \in \mathbb{N}_0}$ in its domain. We have also assumed that the product RA is well-defined on a dense subset of $L^2(\mu)$. We can then compute

$$\begin{aligned} T p_k &= v_k \circ \tau \\ &= \sum_{j=0}^{\infty} A_{j,k} v_j \\ &= \sum_{j=0}^{\infty} \langle p_j | A p_k \rangle_{L^2(\mu)} v_j \\ &= R A p_k. \end{aligned}$$

The last line above holds for every $A p_k$ that is in the domain of R . $\square$

THEOREM 4.8. *Given (X, μ) a Borel measure space with $X \subset \mathbb{R}$ and given τ a measurable map from X to itself, let $N^{(\mu)}$ be the adjusted moment matrix for μ and $N^{(\mu \circ \tau^{-1})}$ the corresponding moment matrix for $\mu \circ \tau^{-1}$. Then, if the matrix A from Equation (4.18) satisfies the hypotheses in Proposition 4.7,*

$$(4.19) \quad A^* N^{(\mu)} A = N^{(\mu \circ \tau^{-1})}.$$

PROOF. We have $RA = T$. Find weights w such that both T_w^* and R_w^* are densely defined on $L^2(\mu)$. Let $N_w^{(\mu)}$ be the weighted moment matrix with entries $\frac{1}{\sqrt{w_i w_j}} \langle v_i | v_j \rangle_{L^2(\mu)}$. Using Equations (4.14) and (4.17), we find that

$$\begin{aligned} N_w^{(\mu \circ \tau^{-1})} &= T_w^* T \\ &= (RA)^*(RA) \\ &= A^* R_w^* R A \\ &= A^* N_w^{(\mu)} A. \end{aligned}$$

$\square$

4.2. Approximation of A with finite matrices

In this section, we explore whether we can perform computations with finite matrices which yield a finite approximation to the infinite matrix A defined in Equation (4.18), and thereby achieve an approximation of the moments for $\mu \circ \tau^{-1}$ from the moments for μ .

Let μ be a Borel measure on a set $X \subset \mathbb{R}$ such that the moments $M_{i,j}^{(\mu)} = \int_X x^{i+j} d\mu(x)$ are finite for all orders. Let $\tau : X \rightarrow X$ be a measurable endomorphism such that the moments with respect to $\mu \circ \tau^{-1}$ are also finite for all orders, and the powers of τ are in the closed span $\mathcal{P}$ of the monomials in $L^2(\mu)$. Let T

be the infinite matrix introduced in Section 4.1 with entries $T_{ij} = \langle p_i | v_j \circ \tau \rangle_{L^2(\mu)}$, where $\{p_i\}_{i \in \mathbb{N}_0}$ are the orthonormal polynomials in $L^2(\mu)$ given by performing the Gram-Schmidt method on the monomials $\{v_j\}_{j \in \mathbb{N}_0}$. Let R be the transformation taking p_i to v_i , so the matrix entries are $R_{i,j} = \langle p_i | v_j \rangle_{L^2(\mu)}$.

Fix n . Let R_n and T_n be (as in Section 3.2) the $(n+1) \times (n+1)$ truncations of the R and T matrices, respectively. Let $\mathcal{H}_n$ be the closed linear span of the monomials $\{v_0, v_1, \dots, v_n\}$ (which is also the closed linear span of $\{p_i\}_{i=0}^n$) and let P_n be the orthogonal projection onto $\mathcal{H}_n$ from $\mathcal{P}$. Note that in the Dirac notation mentioned in Chapter 1,

$$(4.20) \quad P_n = \sum_{i=0}^n |p_i\rangle\langle p_i|.$$

We have proved in Proposition 4.7 that the matrix $A = R^{-1}T$, in the cases where this product of infinite matrices is well defined, gives the coefficients of $v_j \circ \tau$ in terms of the monomials $\{v_i\}_{i \in \mathbb{N}_0}$. We now wish to show that the finite matrix product $R_n^{-1}T_n$ provides an approximation of A , in the sense that it yields coefficients for the projection of the powers of τ onto $\mathcal{H}_n$, expanded in terms of the monomials.

LEMMA 4.9. *For fixed n ,*

$$(4.21) \quad \sum_{j=0}^n (R_n^{-1}T_n)_{j,k} v_j = P_n(v_k \circ \tau).$$

Consequently,

$$(4.22) \quad \lim_{n \rightarrow \infty} \sum_{j=0}^n (R_n^{-1}T_n)_{j,k} v_j = v_k \circ \tau,$$

where convergence is in $L^2(\mu)$.

PROOF. Given our fixed n , let $k \leq n$. In the computation below, recall that for $j, k \leq n$, we have $(R_n^{-1})_{j,k} = R_{j,k}^{-1}$ and $(T_n)_{j,k} = T_{j,k}$. Also, since the orthogonal polynomials $p_0, \dots, p_j$ are in the span of the monomials $v_0, \dots, v_j$, we know that the truncated matrix R_n^{-1} maps v_j to p_j for each $j \leq n$.

$$\begin{aligned} \sum_{j=0}^n (R_n^{-1}T_n)_{j,k} v_j &= \sum_{j=0}^n \sum_{\ell=0}^n (R_n)_{j,\ell}^{-1} (T_n)_{\ell,k} v_j \\ &= \sum_{\ell=0}^n \langle p_\ell | v_k \circ \tau \rangle_{L^2(\mu)} \sum_{j=0}^n (R_n^{-1})_{j,\ell} v_j \quad \text{where both } \ell, j \leq n \\ &= \sum_{\ell=0}^n \langle p_\ell | v_k \circ \tau \rangle_{L^2(\mu)} p_\ell \\ &= P_n v_k \circ \tau \quad \text{by Equation (4.20).} \end{aligned}$$

□

REMARK 4.10. It is important to realize here that even for $j, k \leq n$,

$$(4.23) \quad (R_n^{-1}T_n)_{j,k} \neq (R^{-1}T)_{j,k}.$$

We do, however, know that because the product $R^{-1}T$ is well defined, if we fix j, k and let $n \rightarrow \infty$, we do have $(R_n^{-1}T_n)_{j,k} \rightarrow (R^{-1}T)_{j,k}$.

If we were able to conclude from the truncation result in Lemma 4.9 that $\lim_{n \rightarrow \infty} \sum_{j=0}^n (R^{-1}T)_{j,k} v_j = v_k \circ \tau$, where we take the limit in $L^2(\mu)$, we would have an alternate proof of Proposition 4.7. In other words, we would be able to write

$$(4.24) \quad \sum_{j=0}^{\infty} (R^{-1}T)_{j,k} v_j = v_k \circ \tau.$$

This convergence, however, would require hypotheses about how the sequences $(R_n^{-1}T_n)_{j,k}$ converge to $(R^{-1}T)_{j,k}$ as $n \rightarrow \infty$. To illustrate this point, let us fix k and let $a_j^{(n)} = (R_n^{-1}T_n)_{j,k}$ and $a_j = (R^{-1}T)_{j,k}$, so for each j , $a_j^{(n)} \rightarrow a_j$ as $n \rightarrow \infty$.

$$\begin{aligned} \left\| \sum_{j=0}^n a_j v_j - v_k \circ \tau \right\|_{L^2(\mu)} &= \left\| \sum_{j=0}^n a_j v_j - \sum_{j=0}^n a_j^{(n)} v_j + \sum_{j=0}^n a_j^{(n)} v_j - v_k \circ \tau \right\|_{L^2(\mu)} \\ &\leq \left\| \sum_{j=0}^n (a_j - a_j^{(n)}) v_j \right\|_{L^2(\mu)} + \left\| \sum_{j=0}^n a_j^{(n)} v_j - v_k \circ \tau \right\|_{L^2(\mu)} \end{aligned}$$

Using Lemma 4.9, the second term of the last line above can certainly be made arbitrarily small for large enough n . The first term, however, may not have that property.

For each $k \in \mathbb{N}_0$, the truncated matrix products $R_n^{-1}T_n$ produce asymptotic expansions for $v_k \circ \tau$ by giving expansions of $P_n(v_k \circ \tau)$ in terms of monomials v_j , even if there is no *a priori* known expansion of $v_k \circ \tau$ in the monomials. We assume that such an expansion exists, that is,

$$v_k \circ \tau = \sum_{j=0}^{\infty} a_j v_j.$$

We may, however, have no way of computing the actual coefficients. Lemma 4.9 gives approximations to these coefficients which get better as n increases. The next question is how these finite approximations to the matrix A interact with the moment matrix transformation given in Equation (4.19) from Theorem 4.8. In the case described in Subsection 3.1, where τ is an affine map on $\mathbb{R}$ or $\mathbb{C}$, the matrix A is upper triangular. With this added structure, it is readily demonstrated that the truncated matrix product yields exactly the truncation of the infinite matrix product, i.e.

$$A_n^* M_n^{(\mu)} A_n = [A^* M^{(\mu)} A]_n = M^{(\mu \circ \tau^{-1})}.$$

We also note that in this special case, we have Equation (3.5) from [EST06] giving a concrete expression of the entries of A . We then can compute the exact moments with respect to $\mu \circ \tau^{-1}$ using finite matrix computations.

In the more general case, however, we may not have a construction giving us the entries in A , so we may need to use the finite matrix product $\tilde{A}_n = R_n^{-1}T_n$. In this case, we find the triple product gives entries which are the inner products of projections of powers of the measurable map τ .

Given the map τ , the entries in the adjusted moment matrix for $\mu \circ \tau^{-1}$ are given by the inner product

$$(N^{(\mu \circ \tau^{-1})})_{i,j} = \langle v_i \circ \tau | v_j \circ \tau \rangle_{L^2(\mu)}.$$

COROLLARY 4.11. *Let $\tilde{A}_n = R_n^{-1}T_n$. Then*

$$(4.25) \quad (\tilde{A}_n^* N^{(\mu)} \tilde{A}_n)_{i,j} = \langle P_n(v_i \circ \tau) | P_n(v_j \circ \tau) \rangle_{L^2(\mu)} = \langle v_i \circ \tau | P_n(v_j \circ \tau) \rangle_{L^2(\mu)}.$$

PROOF. This is a consequence of Lemma 4.9.

$$\begin{aligned} (\tilde{A}_n^* N^{(\mu)} \tilde{A}_n)_{i,j} &= \sum_{k=0}^n \sum_{l=0}^n (\tilde{A}_n^*)_{i,k} (N^{(\mu)})_{k,l} (\tilde{A}_n)_{l,j} \\ &= \sum_{k=0}^n \sum_{l=0}^n (\tilde{A}_n)_{k,i} \langle v_k | v_l \rangle_{L^2(\mu)} (\tilde{A}_n)_{l,j} \\ &= \left\langle \sum_{k=0}^n (\tilde{A}_n)_{k,i} v_k \middle| \sum_{l=0}^n (\tilde{A}_n)_{l,j} v_l \right\rangle_{L^2(\mu)} \\ &= \langle P_n(v_i \circ \tau) | P_n(v_j \circ \tau) \rangle_{L^2(\mu)} \\ &= \langle v_i \circ \tau | P_n(v_j \circ \tau) \rangle_{L^2(\mu)}. \end{aligned}$$

In the last line, we use the property of projections that $P_n = P_n^* = P_n^2$. $\square$

CHAPTER 5

The Kato-Friedrichs operator

The moment matrix $M^{(\mu)}$ is an infinite matrix, and while $M^{(\mu)}$ may not be a well defined operator on ℓ^2 , we may be able to view $M^{(\mu)}$ as an operator in some other sequence space. We use the techniques of Kato and Friedrichs to turn $M^{(\mu)}$ into a self-adjoint densely defined operator on a weighted ℓ^2 space. Our main tool will be a quadratic form Q_M which is defined from the moments of μ . We obtain the weighted ℓ^2 space by finding a space in which the quadratic form Q_M is closable. In the course of showing that Q_M is closable, we introduce two key operators: F and its adjoint F^* , which we will continue to study in Chapters 6 and 7.

5.1. The quadratic form Q_M

Let $\mathcal{D}$ be the set of all finitely supported sequences indexed by $\mathbb{N}_0$. The familiar expression in Equation (1.12) for the infinite matrix-vector product Mc makes sense for every $c \in \mathcal{D}$, but Mc may not be well defined for every sequence c . Even when it is well defined, Mc may not be in the same Hilbert space as c . We will address this technicality by changing the domain of M .

Let M be the moment matrix of a positive Borel measure with finite moments, where M satisfies the positive semidefinite condition in Definition 2.1 (real case) or the PDC condition in Definition 2.3 (complex case). The quadratic form Q_M is defined on infinite sequences with only finitely many nonzero components. Specifically, given $c \in \mathcal{D}$ we have

$$(5.1) \quad Q_M(c) = \sum_i \sum_j \bar{c}_i M_{i,j} c_j.$$

Even though every operator on a Hilbert space determines a quadratic form, the converse is not necessarily true. Here, we look for conditions on the quadratic form Q_M which guarantee that the quadratic form gives rise to an operator on a Hilbert space $\mathcal{H}$. The work of Friedrichs and Kato provides the correct conditions. Friedrichs (see [Kat80, Section 6.2.3]) proved that semibounded symmetric operators in a Hilbert space $\mathcal{H}$ have self-adjoint extensions in $\mathcal{H}$. Later, Kato [Kat80, Section 6.2.1-Theorem 2.1; Section 6.2.6-Theorem 2.23] extended Friedrichs's theorem to quadratic forms.

THEOREM 5.1 (Kato). *Let Q be a densely defined, closed, positive quadratic form. Then there exists a unique self-adjoint operator H on a Hilbert space $\mathcal{H}$ such that the Hilbert space completion $\mathcal{H}_Q$ of Q is equal to the completion of H , and for all c in the domain of Q ,*

$$Q(c) = \|H^{1/2}c\|_{\mathcal{H}}^2 = \|c\|_{\mathcal{H}_Q}^2.$$

In particular, the domain of Q is equal to the domain of $H^{1/2}$.

Our quadratic form Q_M arising from a moment matrix may not be closed, but we can show that Q_M is *closable* and then apply Kato's theorem to the closure. We change our notation briefly here in order to work with sequences of elements in $\mathcal{D}$. A sequence in $\mathcal{D}$ will be denoted $\{c_n\}_{n \in \mathbb{N}_0}$, and the i^{th} component of $\{c_n\}$ is $c_n(i)$. (We retain the notation $c = \{c(i)\}_{i \in \mathbb{N}_0}$ for elements of $\mathcal{D}$ and its subsequent completion throughout the next two sections.)

DEFINITION 5.2 ([Kat80]). Let $\mathcal{H}$ be a Hilbert space in which $\mathcal{D}$ is dense. A quadratic form Q defined on $\mathcal{D}$ is *closable* if and only if

$$c_n \rightarrow 0 \text{ in } \mathcal{H}$$

and

$$Q(c_n - c_m) \rightarrow 0 \text{ as } m, n \rightarrow \infty$$

imply

$$Q(c_n) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We next define the operator F to map finite sequences $c \in \mathcal{D}$ to polynomials (a.k.a generating functions) in $L^2(\mu)$. Given $c \in \mathcal{D}$, we define

$$(5.2) \quad Fc(x) = f_c(x) = \sum_{i \in \mathbb{N}_0} c_i x^i.$$

EXAMPLE 5.3. A quadratic form which is not closable in ℓ^2 .

In order to construct a quadratic form which is not closable in ℓ^2 , we need a sequence $\{c_n\} \subset \ell^2$ and a measure μ such that

- (1) $\sum_{i \in \mathbb{N}_0} |c_n(i)|^2 \rightarrow 0$ as $n \rightarrow \infty$
- (2) $\int_{\mathbb{R}} |f_{c_m}(x) - f_{c_n}(x)|^2 d\mu(x) \rightarrow 0$ as $m, n \rightarrow \infty$
- (3) $\int_{\mathbb{R}} |f_{c_n}(x)|^2 d\mu(x) \not\rightarrow 0$ as $n \rightarrow \infty$.

Let $\mu = \delta_b$ for $b > 1$, and let $c_n(i) = \delta(n, i)b^{-i}$ (the Kronecker delta). Since $b > 1$, $\|c_n\|_{\ell^2}^2 = b^{-2n} \rightarrow 0$ as $n \rightarrow \infty$, so (1) is satisfied. Since

$$f_{c_n}(x) = b^{-n}x^n \text{ for each } n,$$

and the integral in (2) simply evaluates f_{c_n} and f_{c_m} at $x = b$, we have

$$\int_{\mathbb{R}} |f_{c_m}(x) - f_{c_n}(x)|^2 d\mu(x) = |1 - 1|^2 = 0 \text{ for all } m, n.$$

However, the integral in (3) is 1 for all n . Therefore the quadratic form associated with δ_b , $b > 1$, is not closable in ℓ^2 . $\diamond$

5.2. The closability of Q_M

In order to show that the quadratic form Q_M is closable for a particular infinite Hankel matrix M , we show that the operator F defined in Equation (5.2) is closable with respect to a weighted space $\ell^2(w)$. Recall Definition 1.2 regarding closed and closable operators. The *weighted* ℓ^2 spaces are defined by a sequence of weights $w = \{w_i\}_{i \in \mathbb{N}_0}$:

$$\ell^2(w) := \left\{ \{c(i)\} \mid \sum_i w_i |c(i)|^2 < \infty \right\}.$$

The norm on $\ell^2(w)$ is

$$\|c\|_{\ell^2(w)}^2 = \sum_{i \in \mathbb{N}_0} w_i |c(i)|^2,$$

and the inner product is given by

$$\langle c|d \rangle_{\ell^2(w)} = \sum_{i \in \mathbb{N}_0} w_i \overline{c(i)} d(i).$$

We will generally take the weights to all be strictly greater than zero.

In the next lemma, we use an argument mirroring the proof of Lemma 4.4 to produce a set of weights w which guarantee that F will be closable in the weighted space $\ell^2(w)$, and then in Theorem 5.5 we show that the quadratic form Q_M is closable in $\ell^2(w)$.

Consider F as a map with domain in $\ell^2(w)$ for some weights $w = \{w_i\}_{i \in \mathbb{N}_0}$. We then see that the adjoint F^* of F is given by

$$(5.3) \quad (F_w^* f)_k = \frac{1}{w_k} \int_{\mathbb{R}} x^k f(x) d\mu(x).$$

To verify this, we must show that F_w^* satisfies

$$(5.4) \quad \langle Fc|f \rangle_{L^2(\mu)} = \langle c|F_w^* f \rangle_{\ell^2(w)}$$

for every f such that $F_w^* f \in \ell^2(w)$.

For each $c \in \mathcal{D}$, we have

$$(5.5) \quad \begin{aligned} \langle Fc|f \rangle_{L^2(\mu)} &= \int_{\mathbb{R}} \sum_{k \in \mathbb{N}_0} \overline{c(k)} x^k f(x) d\mu(x) \\ &= \sum_{k \in \mathbb{N}_0} \overline{c(k)} \int_{\mathbb{R}} x^k f(x) d\mu(x). \end{aligned}$$

We now multiply and divide by w_k :

$$(5.6) \quad \begin{aligned} \langle Fc|f \rangle_{L^2(\mu)} &= \sum_{k \in \mathbb{N}_0} w_k \overline{c(k)} \underbrace{\frac{1}{w_k} \int_{\mathbb{R}} x^k f(x) d\mu(x)}_{(5.3)} \\ &= \langle c|F_w^* f \rangle_{\ell^2(w)}, \end{aligned}$$

where we have shown the right hand side of Equation (5.3) satisfies the definition of the adjoint F_w^* .

LEMMA 5.4. *Suppose $\int_{\mathbb{R}} |x|^k d\mu(x) < \infty$ for each $k \in \mathbb{N}_0$. There exist weights $\{w_i\}_{i \in \mathbb{N}_0}$ such that $F : \ell^2(w) \rightarrow L^2(\mu)$ (5.2) is closable.*

PROOF. The operator F is closable if and only if $\text{dom}(F^*)$ is dense in $L^2(\mu)$. (See, for example, [Con90, Proposition 1.6, Chapter 10].)

Now, $F_w^* f \in \ell^2(w)$ precisely when

$$(5.7) \quad \sum_{k \in \mathbb{N}_0} w_k |(F_w^* f)_k|^2 < \infty,$$

and (5.7) is true if and only if

$$(5.8) \quad \sum_k \frac{1}{w_k} \left| \int_{\mathbb{R}} x^k f(x) d\mu(x) \right|^2 < \infty.$$

This gives us one sufficient criterion for F to be closable. We see from Equation (5.8) that the monomial functions $\{v_j\}_{j \in \mathbb{N}_0}$ are in the domain of F_w^* if and only if

$$(5.9) \quad \sum_k \frac{1}{w_k} \left| \int_{\mathbb{R}} x^{j+k} d\mu(x) \right|^2 = \sum_k \frac{1}{w_k} |M_{j,k}|^2 < \infty$$

for all $j \in \mathbb{N}_0$.

Rather than looking at this criterion (5.9), however, we will find weights that put the essentially bounded functions in the domain of F_w^* . By hypothesis, μ is a finite measure so $L^\infty(\mu) \subseteq L^2(\mu)$. Set $\mathcal{M}_k := \int_{\mathbb{R}} |x|^k d\mu(x)$, and suppose $f \in L^\infty(\mu)$. Then

$$(5.10) \quad \left| \int_{\mathbb{R}} x^k f(x) d\mu(x) \right| \leq \|f\|_{L^\infty(\mu)} \mathcal{M}_k.$$

Therefore, if $\{\mathcal{M}_k^2/w_k\} \in \ell^1$, then

$$(5.11) \quad L^\infty(\mu) \subset \text{dom}(F_w^*),$$

and we explicitly define a choice of weights w_k :

$$(5.12) \quad w_k := (1 + k^2) \mathcal{M}_k^2.$$

Since $L^\infty(\mu)$ is dense in $L^2(\mu)$, we now know that $\text{dom}(F_w^*)$ is dense in $\ell^2(w)$ where $w = \{w_k\}_{k \in \mathbb{N}_0}$ is defined in (5.12). Therefore $F : \ell^2(w) \rightarrow L^2(\mu)$ is closable. $\square$

THEOREM 5.5. *Given μ a Borel measure on $\mathbb{R}$ with finite moments of all orders and $M = M^{(\mu)}$, there are weights $\{w(i)\}_{i \in \mathbb{N}_0}$ such that $(Q_M, \mathcal{D}, \ell^2(w))$ is closable.*

PROOF. Choose weights $w = \{w_i\}_{i \in \mathbb{N}_0}$ as in Equation (5.12) so that F is closable. Suppose $c_n \xrightarrow{\ell^2(w)} 0$ and $Q_M(c_n - c_m) \rightarrow 0$ as $m, n \rightarrow \infty$ as in Definition

5.2. Our goal is to show that $Q_M(c_n) \rightarrow 0$.

Denote the polynomial Fc_n by f_n for each $n \in \mathbb{N}_0$. Because Q_M on $\mathcal{D}$ and $\tilde{Q}_M$ defined on the polynomials satisfy

$$\tilde{Q}_M(F(c)) = Q_M(c),$$

the condition $Q_M(c_n - c_m) \rightarrow 0$ implies that $\{f_n\}_{n \in \mathbb{N}_0}$ is a Cauchy sequence in $L^2(\mu)$. Therefore, there exists $g \in L^2(\mu)$ such that $f_n \xrightarrow{L^2(\mu)} g$.

We will now use the condition that $c_n \rightarrow 0$ in $\ell^2(w)$. By Lemma 5.4, $F : \ell^2(w) \rightarrow L^2(\mu)$ is closable, so its closure $\overline{F}$ has a closed graph. Since $c_n \rightarrow 0$, $F(c_n) = \overline{F}(c_n) \rightarrow g$, and the graph of $\overline{F}$ is closed, g must be $0 \in L^2(\mu)$. $\square$

As a result of Theorem 5.5, we can apply Theorem 5.1 (Kato's Theorem) to the closure of the quadratic form Q_M . We denote by $\mathcal{H}_Q$ the Hilbert space completion with respect to the quadratic form Q_M . By Kato's theorem, there exists a self-adjoint operator H on $\ell^2(w)$ such that the Hilbert space completion is also $\mathcal{H}_Q$. This gives

$$(5.13) \quad \|c\|_{\mathcal{H}_Q}^2 = Q_M(c) = \|H^{1/2}c\|_{\ell^2(w)}^2$$

for all $c \in \mathcal{H}_Q$. Moreover, the domain of $H^{1/2}$ is equal to the domain of Q_M . It should be noted that in general, the domain of H is a subset of the domain of $H^{1/2}$ and may or may not contain $\mathcal{D}$.

5.3. A factorization of the Kato-Friedrichs operator

Given F from $\ell^2(w)$ to $L^2(\mu)$, suppose the weights w have been selected such that F is closable, i.e. F_w^* is densely defined in $L^2(\mu)$.

Let $\tilde{Q}_M$ be the quadratic form given by the $L^2(\mu)$ -norm:

$$(5.14) \quad \tilde{Q}_M(f) = \int_{\mathbb{R}} |f|^2 d\mu.$$

Then for every $c \in \mathcal{D}$, we have

$$(5.15) \quad \begin{aligned} \tilde{Q}_M(Fc) &= \int_{\mathbb{R}} |Fc|^2 d\mu \\ &= \int_{\mathbb{R}} \sum_i \sum_j \bar{c}_i c_j x^{i+j} d\mu \\ &= \sum_i \sum_j \bar{c}_i M_{i,j} c_j = Q_M(c). \end{aligned}$$

PROPOSITION 5.6. $F_w^* F$ is a self-adjoint densely defined operator on $\ell^2(w)$, and $F_w^* F$ is the Kato operator H .

PROOF. The domain of F contains $\mathcal{D}$, and the domain of F_w^* contains the polynomials. Since F maps finite sequences $c \in \mathcal{D}$ to polynomials Fc , we see that $\mathcal{D}$ is contained in the domain of $F_w^* F$. In order to prove that $F_w^* F$ is the Kato operator corresponding to the quadratic form Q_M , we must show that it satisfies

$$Q_M(c) = \|(F_w^* F)^{1/2} c\|_{\ell^2(w)}$$

for all $c \in \mathcal{D}$.

First, note that for any $c \in \mathcal{D}$,

$$\begin{aligned} \langle c | F_w^* F c \rangle_{\ell^2(w)} &= \langle (F_w^* F)^{1/2} c | (F_w^* F)^{1/2} c \rangle_{\ell^2(w)} \\ &= \|(F_w^* F)^{1/2} c\|_{\ell^2(w)}^2. \end{aligned}$$

We also note that

$$\begin{aligned} \langle c | F_w^* F c \rangle_{\ell^2(w)} &= \langle Fc | Fc \rangle_{\ell^2(w)} \\ &= \|Fc\|_2^2 \\ &= \tilde{Q}_M(Fc) = Q_M(c). \end{aligned}$$

Therefore, for $c \in \mathcal{D}$,

$$Q_M(c) = \langle c | F_w^* F c \rangle_{\ell^2(w)} = \|(F_w^* F)^{1/2} c\|_{\ell^2(w)}^2.$$

By uniqueness of the Kato operator for the closable quadratic form Q_M , we have $F_w^* F = H$. $\square$

This proposition makes the usually mysterious Kato operator much more concrete. In particular, as a result of this proposition, the domain of H includes $\mathcal{D}$.

5.4. Kato connection to A matrix

Given a moment matrix $M^{(\mu)}$ and a measurable map τ , we can use Kato theory as another way to describe a matrix A such that, under appropriate convergence criteria,

$$M^{(\mu \circ \tau)} = A^* M^{(\mu)} A.$$

This is a different approach to that used in Section 4.1. We prove that under certain hypotheses on the measure μ and the map τ , we can find an operator $\tilde{A}$ which is an isometry between the Kato completion Hilbert spaces for the quadratic forms

Q_M and Q'_M , where $M = M^{(\mu)}$ and $M' = M^{(\mu \circ \tau^{-1})}$. We demonstrate that this isometry $\tilde{A}$ will intertwine the Kato operators for Q_M and $Q_{M'}$, and therefore the corresponding matrix representation A will intertwine the moment matrices M and M' as required.

Let (X, μ) be a measure space with $X \subseteq \mathbb{R}$. For each $k \in \mathbb{N}_0$, let v_k be the monomial function on X : $v_k(x) = x^k$. The polynomials may not be dense in $L^2(\mu)$, so denote by $\mathcal{P}$ the closed span of the polynomials in $L^2(\mu)$. We take μ to be a measure such that $\{v_k\}_{k \in \mathbb{N}_0} \subseteq L^1(\mu) \cap \mathcal{P}$. Let $M = M^{(\mu)}$ be the moment matrix for μ . By Section 5.1, there exist weights $w = \{w_i\}_{i \in \mathbb{N}_0}$ such that the map F is closable on the weighted space $\ell^2(w)$. Thus the quadratic form Q_M is closable, and we can define the Kato operator H on $\ell^2(w)$. Note that the domains of Q_M , F , and F_w^* all contain the set of finite sequences $\mathcal{D}$.

Let τ be a measurable endomorphism on X such that the functions $v_k \circ \tau = \tau^k$ are all in $L^1(\mu) \cap \mathcal{P}$. Let $M' = M^{(\mu \circ \tau^{-1})}$ be the moment matrix for the measure $\mu \circ \tau^{-1}$. Define the map $F' : \mathcal{D} \rightarrow L^2(\mu \circ \tau^{-1})$ to be the analog of F , i.e. the map which takes $c \in \mathcal{D}$ to the polynomial $f_c = \sum_i c_i v_i \in L^2(\mu \circ \tau^{-1})$. Using the same reasoning as that used in Section 5.1, there exist weights such that F' is closable, hence the quadratic form $Q_{M'}$ is closable, and we have H' , the Kato operator on the corresponding weighted ℓ^2 space.

Let $w = \{w_k\}_{k \in \mathbb{N}_0}$ be weights that allow for both F and F' to be closable on $\ell^2(w)$, hence the Kato operators H and H' are both densely defined self-adjoint operators on $\ell^2(w)$. Denote the Kato completion Hilbert spaces for the quadratic forms Q_M and $Q_{M'}$ by $\mathcal{H}_Q$ and $\mathcal{H}_{Q'}$, respectively.

Notice that $\phi \in L^2(\mu \circ \tau^{-1})$ if and only if $\int_{\mathbb{R}} |\phi \circ \tau|^2 d\mu < \infty$, which holds if and only if $\phi \circ \tau \in L^2(\mu)$. In fact, there is a natural isometry α between $L^2(\mu \circ \tau^{-1})$ and $L^2(\mu)$ given by

$$\alpha(\phi) = \phi \circ \tau.$$

REMARK 5.7. The map F which we have defined above actually takes on several realizations in this section. Simply put, it is the map that takes finite sequences to the polynomial with these coefficients. But we can think of F as operating on the Kato spaces $\mathcal{H}_Q$, $\mathcal{H}_{Q'}$ or on the weighted space $\ell^2(w)$. We can also take the codomain of F to be $L^2(\mu)$ or $L^2(\mu \circ \tau^{-1})$. In order to clarify these different realizations, we will denote the operator F as follows:

$$\begin{aligned} F_Q &: \mathcal{H}_Q \rightarrow L^2(\mu) \\ F_{Q'} &: \mathcal{H}_{Q'} \rightarrow L^2(\mu \circ \tau^{-1}) \\ F &: \ell^2(w) \rightarrow L^2(\mu) \\ F' &: \ell^2(w) \rightarrow L^2(\mu \circ \tau^{-1}). \end{aligned}$$

Moreover, we also observe that the operator T (defined below) which maps standard basis elements e_i to τ^i can be expressed in terms of the F' and the isometry α . Given $c \in \mathcal{D}$,

$$\begin{aligned} Tc &= \left(\sum c_i \tau^i \right) \in L^2(\mu) \\ &= \sum c_i \alpha(v_i) \in L^2(\mu) \\ &= \alpha F' c. \end{aligned}$$

We will show below that the operators F_Q and $F_{Q'}$ are isometries, and we have already proved that for appropriately chosen weights, the operators F and F' are closable on $\ell^2(w)$.

LEMMA 5.8. *The Kato space $\mathcal{H}_Q$ is isometric to $\mathcal{P} \subseteq L^2(\mu)$, and $\mathcal{H}_{Q'}$ is isometric to a subspace of $L^2(\mu \circ \tau^{-1})$.*

PROOF. The norm on the Hilbert space $\mathcal{H}_Q$ is given by

$$\|c\|_{\mathcal{H}_Q}^2 = Q_M(c)$$

when $c \in \mathcal{D}$, and by the definition of the Kato operator, we also have

$$\|c\|_{\mathcal{H}_Q} = \|H^{1/2}c\|_{\ell^2(w)}.$$

Let F_Q be the map taking $c \in \mathcal{D}$ to $f_c = \sum_i c_i v_i$ in $\mathcal{P} \subseteq L^2(\mu)$. Then

$$\begin{aligned} \|F_Q c\|_2^2 &= \|f_c\|_2^2 \\ &= \int_{\mathbb{R}} \left| \sum_i c_i v_i \right|^2 d\mu \\ &= \sum_i \sum_j \overline{c_i} c_j \int_{\mathbb{R}} v_i v_j d\mu \\ &= \sum_i \sum_j \overline{c_i} c_j M_{i,j}^{(\mu)} \\ &= Q_M(c) = \|c\|_{\mathcal{H}_Q}^2. \end{aligned}$$

Thus F_Q is an isometry from $\mathcal{H}_Q$ to $\mathcal{P}$. An identical argument proves that the corresponding $F_{Q'}$ map from $\mathcal{H}_{Q'}$ to $L^2(\mu \circ \tau^{-1})$ is also an isometry onto the subspace $\mathcal{P}'$ of $L^2(\mu \circ \tau^{-1})$ spanned by the polynomials. $\square$

PROPOSITION 5.9. *The Kato Hilbert spaces $\mathcal{H}_Q$ and $\mathcal{H}_{Q'}$ are isometric.*

PROOF. We use the notation $\mathcal{H} \cong \mathcal{G}$ if there exists an isometry from the Hilbert space $\mathcal{H}$ onto the Hilbert space $\mathcal{G}$. Using Lemma 5.8 and the map α , we have the following isometries:

$$\mathcal{H}_{Q'} \cong \mathcal{P}' \cong \mathcal{P} \cong \mathcal{H}_Q.$$

$\square$

If we denote the isometry in Proposition 5.9 by $\tilde{A}$, then

$$\tilde{A} = F_Q^{-1} \alpha F_{Q'}.$$

The isometry property $\|c\|_{Q'} = \|\tilde{A}c\|_Q$ implies the quadratic forms satisfy $Q_M(\tilde{A}c) = Q_{M'}(c)$ for all $c \in \mathcal{D}$.

Given $\{e_i\}_{i \in \mathbb{N}_0}$ the standard orthonormal basis for ℓ^2 , we define the standard orthonormal basis $\{e_i^w\}_{i \in \mathbb{N}_0} = \{\frac{e_i}{\sqrt{w_i}}\}_{i \in \mathbb{N}_0}$ for $\ell^2(w)$. Given F from $\ell^2(w)$ to $L^2(\mu)$, suppose the weights w have been selected such that F is closable, i.e. F_w^* is densely defined in $L^2(\mu)$. Recall from Lemma 5.6 that $F_w^* F$ is equal to the Kato operator H on $\ell^2(w)$.

The operator $\tilde{A}$ has domain containing $\mathcal{D} \subseteq \mathcal{H}_Q$, but we will need to consider the analog of $\tilde{A}$ as an operator on $\ell^2(w)$. In order to do this, we define the following

maps:

$$\begin{aligned} j_w &: \ell^2(w) \rightarrow \mathcal{H}_Q \\ j'_w &: \ell^2(w) \rightarrow \mathcal{H}_{Q'}. \end{aligned}$$

Each of j_w, j'_w acts like the identity map on $\mathcal{D}$, mapping e_i^w in $\ell^2(w)$ to e_i^w in the Kato spaces $\mathcal{H}_Q, \mathcal{H}_{Q'}$, respectively. Then, we see

$$F = F_Q j_w \quad \text{and} \quad F' = F_{Q'} j'_w.$$

We then wish to define the operator on $\ell^2(w)$:

$$A = j'_w \tilde{A} j_w^{-1}.$$

REMARK 5.10. Defining A in this fashion requires the composition of operators to be defined on $\mathcal{D}$ and also requires j_w be invertible. We will assume these properties through the remainder of this section, but we note that they may not hold in general. Thus, this approach requires convergence hypotheses to define A , just as were needed in Proposition 4.7.

Given the maps α and F' defined above, we define T to be the composition

$$T = \alpha F' : \ell^2(w) \rightarrow L^2(\mu),$$

where T maps basis element e_i^w to the function $\frac{1}{\sqrt{w_i}} \tau^i = \frac{1}{\sqrt{w_i}} v_i \circ \tau$.

We now wish to write T with respect to the isometry $\tilde{A}$ and the map $F = F_Q j_w$. We will need to make use of the operator A , as shown in the following diagram.

$$\begin{array}{ccc} L^2(\mu \circ \tau^{-1}) & \xrightarrow{\alpha} & L^2(\mu) \\ F_{Q'} \uparrow & & F_Q \uparrow \\ \mathcal{H}_{Q'} & \xrightarrow{\tilde{A}} & \mathcal{H}_Q \\ j'_w \uparrow & & j_w \uparrow \\ \ell^2(w) & \xrightarrow{A} & \ell^2(w) \end{array}$$

LEMMA 5.11. *Given the operator T defined above,*

$$T = F j_w^{-1} \tilde{A} j'_w = F A.$$

PROOF. Recall from the definition of $\tilde{A}$ that $\alpha F'_Q = F_Q \tilde{A}$. We expand T using the diagram above.

$$\begin{aligned} T &= \alpha F'_w \\ &= \alpha F'_Q j'_w \\ &= F_Q \tilde{A} j'_w \\ &= F j_w^{-1} \tilde{A} j'_w = F A. \end{aligned}$$

□

LEMMA 5.12. *There exist weights such that T is a closable operator on the weighted space $\ell^2(w)$.*

PROOF. We have proven that there exist weights $w = \{w_i\}_{i \in \mathbb{N}_0}$ already that make the operators F and F' closable, and in particular, these weights ensure that the bounded functions are in the domains of both F_w^* and $(F'_w)^*$. Since α is an isometry, these weights also ensure that T_w^* has dense domain. $\square$

LEMMA 5.13. *Let w be weights such that F and T are both closable operators on $\ell^2(w)$. If the operator T_w^*T is self-adjoint and densely defined on $\ell^2(w)$, then $T_w^*T = H'$.*

PROOF. This proof repeats the argument from Lemma 5.6:

$$\begin{aligned} \langle e_i^w | T_w^* T e_j^w \rangle_{\ell^2(w)} &= \langle T e_i^w | T e_j^w \rangle_{L^2(\mu)} \\ &= \frac{1}{\sqrt{w_i w_j}} \langle \tau^i | \tau^j \rangle_{L^2(\mu)} \\ &= \frac{1}{\sqrt{w_i w_j}} \langle v_i | v_j \rangle_{L^2(\mu \circ \tau^{-1})} \\ &= M'_{i,j} = \frac{1}{\sqrt{w_i w_j}} M_{i,j}^{(\mu \circ \tau^{-1})}. \end{aligned}$$

Therefore, we have

$$Q_{M'}(c) = \langle c | M' c \rangle_{\ell^2(w)} = \|(T^*T)^{1/2}c\|_{\ell^2(w)}^2$$

for all $c \in \mathcal{D}$, which proves T_w^*T is exactly the Kato operator H' for $Q_{M'}$. This proves T_w^*T is self-adjoint and densely defined on $\ell^2(w)$. $\square$

Let A be the operator $j_w^{-1} \tilde{A} j'_w$ on $\ell^2(w)$, so $T = FA$.

THEOREM 5.14. *The operator A intertwines the Kato operators H and H' , and thus in matrix form intertwines the moment matrices $M^{(\mu)}$ and $M^{(\mu \circ \tau^{-1})}$.*

PROOF. If the product $T = FA$ is defined on $\mathcal{D}$, we have

$$T_w^*T = (FA)^*FA = A^*(F_w^*F)A,$$

which by Lemmas 5.6 and 5.13 implies that

$$H' = A^*HA.$$

If we form the matrices for A , H , and H' with respect to the orthonormal basis $\{e_i^w\}_{i \in \mathbb{N}_0}$, we have

$$A^* M^{(\mu)} A = M^{(\mu \circ \tau^{-1})}.$$

$\square$

Using the Kato approach we can therefore show that, under the appropriate conditions on τ and the measures μ and $\mu \circ \tau$, we can find a matrix A such that

$$A^* M^{(\mu)} A = M^{(\mu \circ \tau^{-1})}.$$

5.5. Examples

The following examples illustrate the situations in which infinite matrices do not have direct realizations as ℓ^2 operators, but can be realized as operators on a certain weighted Hilbert space. The simplest such case arises when the measure μ is a Dirac mass at the point $b = 1$, which we observed earlier in Example 1.8.

EXAMPLE 5.15. The Kato operator for $\mu = \delta_1$.

Let $\mu = \delta_1$ be the Dirac measure on $\mathbb{R}$ with point mass at 1. As we stated in Example 1.8, the moment matrix $M = M^{(\mu)}$ will be $M_{i,j} = 1$ for all $i, j \in \mathbb{N}_0$. The associated quadratic form Q_M is given on $\mathcal{D}$ by

$$Q_M(c) = \left| \sum_{k \in \mathbb{N}_0} c_k \right|^2.$$

If we attempt to define the operator F^* on the constant function $f(x)$ without weights, we have for each i ,

$$(F^*f)_i = \int_{\mathbb{R}} f(x)x^i d\delta_1(x) = f(1).$$

The weights are necessary in order to make any function with $f(1) \neq 1$ be in the domain of F^* .

Select weights $w = \{w_i\} \subset \mathbb{R}^+$ such that

$$\sum_{i \in \mathbb{N}_0} \frac{1}{w_i} < \infty.$$

Then F_w^* and Q_M are closable, and the resulting Kato-Friedrichs operator H_w then satisfies Equation (5.13). Given c is in the domain of H_w , we have

$$Q_M(c) = \sum_j \sum_k \overline{c_j} c_k = \langle c | H_w c \rangle_{\ell^2(w)} = \sum_j w_j \overline{c_j} (H_w c)_j$$

for all $j \in \mathbb{N}_0$. Thus, H_w is defined by

$$(H_w c)_j = \frac{1}{w_j} \sum_k c_k.$$

In other words, H_w is a rank-one operator with range equal to the span of the vector

$$\xi = \begin{bmatrix} \frac{1}{w_0} & \frac{1}{w_1} & \frac{1}{w_2} & \cdots \end{bmatrix}.$$

◇

EXAMPLE 5.16. The Laplace transform and the operator F^* associated with the measure $e^{-x} dx$ on $\mathbb{R}^+$

We first observed in Examples 1.9 and 2.12 that the moment matrix entries

$$M_{i,j} = (i+j)!$$

grow quickly as i and j increase. Therefore, we again need weights to make F a closable operator.

Let $w = \{w_i\}_{i \in \mathbb{N}_0}$ be weights given by

$$w_i = [(2i)!]^2.$$

We claim that with these weights, F is closable.

$$\sum_{i=0}^{\infty} \frac{1}{w_i} |M_{i+j}|^2 = \sum_{i=0}^{\infty} \frac{(i+j)!}{[(2i)!]^2}$$

for each fixed $j \in \mathbb{N}_0$. Using the ratio test, we see that the series converges for each j so by Equation (5.9) we have the monomials in the domain of F_w^* , which makes F closable on $\ell^2(w)$.

As a side note, we observe that F_w^* has a connection, under appropriate conditions on the function f , to the Laplace transform $\mathcal{L}$:

$$(F_w^* f)_i = \frac{1}{w_i} \int_0^{\infty} f(x) x^i e^{-x} dx = \frac{(-1)^i}{w_i} \left(\frac{d}{ds} \right)^i \mathcal{L}_f(1),$$

where

$$\mathcal{L}_f(s) := \int_0^{\infty} e^{-sx} \xi(x) dx$$

for $s \in \mathbb{R}^+$.

◇

CHAPTER 6

The integral operator of a moment matrix

In this chapter, we will first show how the Hilbert matrix M , which is the moment matrix for Lebesgue measure on $[0, 1]$, can be associated to an integral operator. It turns out that both the operator associated to M and the integral operator are bounded. In fact, we use properties of Hardy space functions and the polar decomposition to show M and its integral operator are unitarily equivalent. The bounded operator properties of the Hilbert matrix are well-known. See [Hal67, Wid66] for more details.

We will generalize Widom's technique, using a weighted Hilbert space when necessary, to find an integral operator on $L^2(\mu)$ associated with the moment matrix $M^{(\mu)}$, where μ is a positive Borel measure supported on $[-1, 1]$. In the more general setting, the integral operator may not be bounded. In Chapter 7 we will examine conditions under which the integral operator is bounded.

When we refer to the Hilbert matrix, where μ is Lebesgue measure restricted to $[0, 1)$, we will use the symbol M ; for any other measure μ supported in $[-1, 1]$, the moment matrix will be denoted $M^{(\mu)}$.

6.1. The Hilbert matrix

The Hilbert matrix M is the moment matrix for Lebesgue measure restricted to $[0, 1)$. M has $(i, j)^{\text{th}}$ entry given by

$$(6.1) \quad M_{i,j} := \frac{1}{1+i+j} = \int_0^1 x^{i+j} dx.$$

It is well-known [Hal67, Wid66] that the Hilbert matrix defines a bounded operator on ℓ^2 , and the operator norm of M as an operator from ℓ^2 to ℓ^2 is π .

Using our definition of $\mathcal{D}$ to be the set of sequences with only finitely many nonzero coordinates, recall the operator $F : \mathcal{D} \rightarrow \mathcal{P}[0, 1]$ given by

$$Fc(x) = \sum_{n \in \mathbb{N}_0} c_n x^n.$$

The operator F defines a correspondence between the finite sequences $\mathcal{D}$ and the polynomials on $[0, 1]$, and this correspondence extends by Stone-Weierstrass and the Riesz-Fisher theorem to an operator from ℓ^2 to $L^2[0, 1]$. We will often call the image of a sequence c a *generating function* $Fc = f_c$.

Suppose $f \in L^2(0, 1)$ and $c \in \ell^2$. By a Fubini argument and the Cauchy-Schwarz inequality, we can switch the sum and the integral in $\langle f | Fc \rangle_{L^2(0,1)}$ to

define the adjoint operator F^* :

$$\begin{aligned}\langle f|Fc\rangle_{L^2} &= \int_0^1 \overline{f(x)} \sum_{n=0}^{\infty} c_n x^n \, dx \\ &= \sum_{n=0}^{\infty} c_n \int_0^1 \overline{f(x)} x^n \, dx \\ &= \langle F^* f | c \rangle_{\ell^2},\end{aligned}$$

where we have the adjoint now defined by

$$(F^* f)_n = \int_0^1 f(x) x^n \, dx.$$

Furthermore, if $f \in L^2(0, 1)$, then $\{(F^* f)_n\}_{n \in \mathbb{N}_0} \in \ell^2$.

LEMMA 6.1. *Let M be the operator for the Hilbert matrix, and let F and F^* be the associated operators defined previously. Then $F^* F = M$.*

PROOF. Given $c \in \ell^2$, the matrix product Mc is well defined since M is a bounded operator on ℓ^2 . We can therefore write $(Mc)_i = \sum_{j=0}^{\infty} M(i, j) c_j = \sum_{j=0}^{\infty} \frac{c_j}{i+j+1}$, where convergence is in the ℓ^2 norm. Then

$$\begin{aligned}(F^* F c)_i &= \int_0^1 x^i (F c)(x) \, dx \\ &= \int_0^1 x^i \sum_{j=0}^{\infty} c_j x^j \, dx \\ &= \sum_{j=0}^{\infty} c_j \int_0^1 x^{i+j} \, dx \\ &= \sum_{j=0}^{\infty} \frac{c_j}{i+j+1}.\end{aligned}$$

The exchange of the sum and integral above follows from Fubini since $c \in \ell^2$ and $\left(\frac{1}{i+j+1}\right) \in \ell^2$ (as a sequence in j). $\square$

We now turn to a relation between the operator M and an integral operator K . This relationship is shown in a more general setting in [Wid66], and we will also discuss a generalized version in Section 6.2.

THEOREM 6.2 ([Wid66], Lemma 3.1). *Suppose $M = F^* F$ is the operator representing the Hilbert matrix. The self-adjoint operator $K = F F^*$ on $L^2[0, 1]$ is an integral operator with kernel*

$$k(x, y) = \frac{1}{1 - xy}.$$

Moreover, K and M satisfy the relation on ℓ^2 :

$$(6.2) \quad KF = FM.$$

PROOF. Let $f \in L^2[0, 1]$ and fix $x \in (0, 1)$. Then

$$\begin{aligned} (FF^*f)(x) &= \sum_{j=0}^{\infty} (F^*f)_j x^j \\ &= \sum_{j=0}^{\infty} \int_0^1 f(y) y^j x^j dy \\ &= \int_0^1 \sum_{j=0}^{\infty} f(y) (xy)^j dy \\ &= \int_0^1 \frac{f(y) dy}{1 - xy}. \end{aligned}$$

We again use Fubini to exchange the sum and integral above, so FF^* is the desired integral operator. Combining this equation with Lemma 6.1, we find

$$KF = FF^*F = FM.$$

□

We next develop tools which we will use in the next section to study the spectrum of the integral operator K . We will demonstrate the relationship between the integral transform and the Hardy space $\mathbb{H}_2$ on the open unit disc $\mathbb{D}$ in $\mathbb{C}$.

The Hardy space $\mathbb{H}_2$ consists of functions which are analytic on $\mathbb{D}$. The Hardy space norm of a function $F(z)$ with expansion $F(z) = \sum_{n=0}^{\infty} c_n z^n$ is given by

$$\|F\|_{\mathbb{H}_2}^2 = \sum_{n=0}^{\infty} |c_n|^2.$$

The inner product on $\mathbb{H}_2$ is exactly the ℓ^2 inner product of the expansion coefficients of two Hardy functions.

Given $z \in \mathbb{D}$, we let $k_z(x) = (1 - zx)^{-1}$. Then $\tilde{K}$ denotes the integral operator with kernel k_z , which is the extension of K defined in Theorem 6.2 to the complex unit disc $\mathbb{D}$. Given $g \in L^2[0, 1]$, we define

$$(\tilde{K}g)(z) = \int_0^1 \frac{g(x)}{1 - \bar{z}x} dx = \langle k_z | g \rangle_{L^2[0,1]}.$$

Note that as a function of $z \in \mathbb{D}$, k_z is analytic for $x \in [-1, 1]$:

$$\frac{1}{1 - zx} = \sum_{n \in \mathbb{N}_0} (zx)^n.$$

Next, we prove that the range of $\tilde{K}$ is contained in $\mathbb{H}_2$. In particular, we show that if $f = \tilde{K}g$, with $g \in L^2[0, 1]$, then f can be extended to an analytic function $F(z)$ on the open unit disc $\mathbb{D}$.

LEMMA 6.3. *Suppose $f \in L^2[0, 1]$. Then*

$$F(z) = \int_0^1 \frac{f(x)}{1 - zx} dx \in \mathbb{H}_2.$$

PROOF. Given $x \in [0, 1]$ and $|z| < 1$, we have

$$F(z) = \int_0^1 f(x) \sum_{n=0}^{\infty} x^n z^n dx.$$

Next, observe that $(\int_0^1 |f(x)| x^n dx)$ is in ℓ^2 as a sequence in n , since by Cauchy-Schwarz

$$\int_0^1 |f(x)| x^n dx = \langle |f| |x^n \rangle_{L^2[0,1]} \leq \|f\|_{L^2[0,1]} \|x^n\|_{L^2[0,1]} = \frac{\|f\|_{L^2[0,1]}}{2n+1}.$$

This allows us to use Fubini's theorem to write

$$F(z) = \sum_{n=0}^{\infty} \left(\int_0^1 f(x) x^n dx \right) z^n$$

and also shows that $F \in \mathbb{H}_2$. $\square$

The adjoint operator $\tilde{K}^*$ to our extended integral operator $\tilde{K}$ is a map from $\mathbb{H}_2$ to $L^2[0, 1]$. Let $\phi \in \mathbb{H}_2$ have expansion $\phi(z) = \sum_{n=0}^{\infty} c_n z^n$. Then,

$$\begin{aligned} \langle \tilde{K}f | \phi \rangle_{\mathbb{H}_2} &= \sum_{n=0}^{\infty} \left(\int_0^1 f(x) x^n dx \right) c_n \\ &= \int_0^1 \overline{f(x)} \sum_{n=0}^{\infty} c_n x^n dx \\ &= \langle f | \phi \rangle_{L^2[0,1]}. \end{aligned}$$

We use Fubini's Theorem above to switch the sum and the integral, and note that in the last line, ϕ is restricted to the real interval $[0, 1]$. Thus, we find that $\tilde{K}^* \phi$ is the restriction of ϕ to $[0, 1]$. We now provide an argument using the polar decomposition and Hardy spaces to show that the operator F is in fact a unitary operator, so the operators M and K have the same spectrum. Using the definitions of F and F^* , we note that the polynomials are in the domain of F^* . This proves that F^* has a dense domain, and thus F is closable. We therefore have by a result of von Neumann that both F^*F and FF^* are self-adjoint positive operators with dense domains.

By polar decomposition, given the closable operator $F : \ell^2 \rightarrow L^2[0, 1]$, there is a partial isometry $U : \ell^2 \rightarrow L^2[0, 1]$ such that

$$F = U(F^*F)^{1/2} = (FF^*)^{1/2}U.$$

Because U is a partial isometry, it is an isometry on the orthogonal complement of the kernel of F . Our goal now is to show that U is, in fact, a unitary operator, i.e. the kernels of F and F^* are trivial. This will give us a unitary equivalence between the operators M and K .

Given $c \in \ell^2$, let $Fc = 0$, that is, $\sum_{n=0}^{\infty} c_n x^n = 0$ for a.e. x in $[0, 1]$. But, by the definition of Hardy functions, $\sum_{n=0}^{\infty} c_n x^n$ is the restriction to $[0, 1]$ of an $\mathbb{H}_2$ function

$$F(z) = \sum_{n=0}^{\infty} c_n z^n.$$

Since F is analytic and is zero on a set with accumulation points, we have $F(z) = 0$ on $\mathbb{D}$. Therefore $c_n = 0$ for all n , which gives $\ker(F) = \{0\}$.

For a function $f \in L^2[0, 1]$, assume that $F^*f = 0$. Thus, $\int_0^1 f(x)x^n dx = 0$ for all $n \in \mathbb{N}_0$. This means that f is orthogonal to the monomials, and hence to the polynomials. Since the polynomials are dense in $L^2[0, 1]$ by Stone-Weierstrass, we have that $f(x) = 0$ a.e. x . Therefore, $\ker(F) = \{0\}$, and we can now conclude that the operator U is a unitary operator from ℓ^2 to $L^2[0, 1]$.

Given our definition of U from the polar decomposition, we have $U(F^*F)^{1/2} = (FF^*)^{1/2}U$. This gives

$$U(F^*F)^{1/2}U^* = (FF^*)^{1/2},$$

and squaring both sides gives

$$U(F^*F)U^* = UMU^* = FF^* = K.$$

6.2. Integral operator for a measure supported on $[-1, 1]$

In this section we study measures μ with finite moments of all orders on $\mathbb{R}$. We generalize the results found in Section 6.1 for the Hilbert matrix. Given a measure μ , we see that the moment matrix $M^{(\mu)}$ may be realized in two ways:

- (1) as an operator H having dense domain in ℓ^2 or a weighted space $\ell^2(w)$ and acting via formal matrix multiplication by $M^{(\mu)}$; and
- (2) as an integral operator K in the $L^2(\mu)$ -space of all square integrable functions.

While the resulting duality is also true in $\mathbb{R}^d$, for any value of d , for clarity we will present the details here just for $d = 1$. The reader will be able to generalize to $d > 1$. We will also assume our measures to be supported in $[-1, 1]$.

Given the moment matrix $M^{(\mu)}$, define a quadratic form Q_M as in Definition 5.1 on the dense space $\mathcal{D}$ consisting of finite sequences. Recall from Section 5.1 that if the monomial functions $\{v_k\}_{k \in \mathbb{N}_0}$ are in $L^1(\mu) \cap L^2(\mu)$, there exist weights $w = \{w_i\}_{i \in \mathbb{N}_0}$ such that Q_M is a closable quadratic form. (Note: in some cases such as the Hilbert matrix, the weights are not required.) Moreover, the operator $F : \ell^2(w) \rightarrow L^2(\mu)$ which takes $c \in \mathcal{D}$ to the generating function (a polynomial) $f_c(x) = \sum_j c_j x^j$ is also made closable by the same weights. Then, as we showed previously,

$$Q_M(c) = \int |f_c(x)|^2 d\mu(x)$$

for all $c \in \mathcal{D}$. By Kato's theory, Q_M has a corresponding closable linear operator H on $\ell^2(w)$ with dense domain satisfying $Q_M(c) = \|H^{1/2}c\|_{\ell^2(w)}^2$. In particular, the Hilbert space completion $\mathcal{H}_Q$ with respect to the quadratic form Q_M is the same as the completion with respect to the operator H . (See Remark 6.7 for another viewpoint on this space $\mathcal{H}_Q$.) We also showed in Lemma 5.6 that the Kato operator H is given by F_w^*F , which shows that the domain of H contains $\mathcal{D}$. Thus, for $c \in \mathcal{D}$, we have $Q_M(c) = \langle c|Hc \rangle_{\ell^2(w)}$.

We will show below that the operator FF_w^* on $L^2(\mu)$ is equal on its domain to the integral operator with kernel

$$k(x, y) = \sum_k \frac{(xy)^k}{w_k}.$$

Note that if weights are not required, the kernel above reduces to the same kernel which represented the Hilbert matrix in the previous section. The integral operators corresponding to moment matrices $M^{(\mu)}$ for different measures μ might have the same kernel, but they will act on different Hilbert spaces $L^2(\mu)$.

We now can state the association between the Kato operator F_w^*F for a moment matrix $M^{(\mu)}$, the operator $K = F_w^*$, and an integral operator on $L^2(\mu)$. As usual, we use the notation $v_j(x) = x^j$ for the monomials in $L^2(\mu)$.

PROPOSITION 6.4. *Let μ be a Borel measure on $\mathbb{R}$ with support contained in $[-1, 1]$ and $v_j \in L^1(\mu)$ for all $j \in \mathbb{N}_0$. Let F be the closure of the operator in (5.2) which sends a sequence $c \in \mathcal{D}$ to the polynomial $f_c \in L^2(\mu)$. Let $w = \{w_i\}_{i \in \mathbb{N}_0}$ be weights such that F_w^* has dense domain as an operator into $\ell^2(w)$, i.e. via Equation (5.9) we assume that*

$$(6.3) \quad \sum_{j \in \mathbb{N}_0} \frac{1}{w_j} |M_{j+k}|^2 < \infty \text{ for all } k \in \mathbb{N}_0.$$

Then the operator $K = \tilde{F}F_w^$ is a self-adjoint operator with dense domain. On its domain, K is an integral operator with kernel*

$$k(x, y) = \sum_j \frac{(xy)^j}{w_j}.$$

PROOF. For the weights w , F is closable and the domain of F_w^* is given by

$$(6.4) \quad \left\{ \phi \in L^2(\mu) \mid \sum_{j \in \mathbb{N}_0} w_j |(F_w^* \phi)_j|^2 < \infty \right\}.$$

We note in particular that the monomials $\{v_j\}_{j \in \mathbb{N}_0}$ are contained in the domain of F_w^* . The domain of FF_w^* is the set of all functions ϕ in the domain of F_w^* such that $F_w^*(\phi)$ is in the domain of F . This set is dense by von Neumann's polar decomposition but may not contain all of the polynomials.

Let $\phi \in \text{dom}(FF_w^*)$. Then a Lebesgue dominated convergence argument gives

$$(6.5) \quad \begin{aligned} (FF_w^*)\phi(x) &= \sum_{k \in \mathbb{N}_0} (F^* \phi)_k v_k(x) \\ &\stackrel{(5.3)}{=} \sum_{k \in \mathbb{N}_0} \frac{1}{w_k} \int_{\text{supp}(\mu)} \phi(y) y^k d\mu(y) x^k \\ &\stackrel{(6.3)}{=} \int_{\text{supp}(\mu)} \phi(y) \sum_{k \in \mathbb{N}_0} \frac{(xy)^k}{w_k} d\mu(y). \end{aligned}$$

The requirement that μ is supported on $[-1, 1]$ ensures that the sum above converges for all x, y . $\square$

The polar decomposition of F is

$$(6.6) \quad F = U(F_w^*F)^{1/2},$$

where U is a partial isometry. We therefore also have

$$(6.7) \quad F = (FF^*)^{1/2}U.$$

The partial isometry $U : \ell^2 \rightarrow L^2(\mu)$ is the same in both (6.6) and (6.7).

COROLLARY 6.5. *The two operators $H = F_w^*F$ and $K = FF_w^*$ have the same spectrum, apart from the point 0.*

PROOF. A rearrangement of (6.6) and (6.7) yields

$$(6.8) \quad UHU = K^*.$$

□

EXAMPLE 6.6. Convex combinations of Dirac masses.

Let $\mu = \sum_{i=1}^n \alpha_i \delta_{x_i}$, where $\alpha_i \geq 0$, $\sum_{i=1}^n \alpha_i = 1$ and $x_i, i = 1, 2, \dots, n$ are distinct real numbers. Recall from Example 5.15 that weights are required for any Dirac measure in order to make F closable. The same holds for convex combinations of these measures. It turns out, however, that the integral operator is not so difficult to describe. We show here that $L^2(\mu)$ has dimension n , so the corresponding integral operator must have a representation as a finite matrix.

Suppose the polynomial $f_c(x) = \sum_{k=0}^m c_k x^k$ is equal to the zero vector in $L^2(\mu)$. In other words,

$$\|f_c\|_{L^2(\mu)}^2 = \int_{\mathbb{R}} \left| \sum_{k=0}^m c_k x^k \right|^2 d\mu(x) = 0.$$

Using our usual notation, f_c is the image of a sequence $c \in \mathcal{D}$ under the map F_Q from the Hilbert space $\mathcal{H}_Q$. We recall from Lemma 5.8 that F_Q is an isometry, hence

$$\|f_c\|_{L^2(\mu)}^2 = Q_M(c) = \|c\|_{\mathcal{H}_Q}^2$$

where $M = M^{(\mu)}$ is the moment matrix for μ . The matrix M is of the form

$$M_{j,k} = \sum_{i=1}^n \alpha_i x_i^{j+k}.$$

Therefore, if $f_c = 0$ in $L^2(\mu)$, we have

$$\begin{aligned} \int_{\mathbb{R}} \left| \sum_{k=0}^m c_k x^k \right|^2 d\mu(x) &= \sum_{k,j=0}^m \overline{c_j} c_k M_{j,k} \\ &= \sum_{j,k=1}^m \overline{c_j} c_k \sum_{i=1}^n \alpha_i x_i^{j+k} \\ &= \sum_{i=1}^n \alpha_i \left| \sum_{k=0}^m c_k x_i^k \right|^2 \\ &= 0. \end{aligned}$$

Since each $\alpha_i \geq 0$ and $\sum_{i=1}^n \alpha_i = 1$, the above sum is zero if and only if

$$\sum_{k=0}^m c_k x_i^k = 0$$

for all $i = 1, 2, \dots, n$. Therefore, the distinct real numbers $x_1, \dots, x_n$ are roots of any polynomial which is equal to zero in $L^2(\mu)$.

Let $p(x)$ be the degree- n polynomial

$$p(x) = \prod_{i=1}^n (x - x_i).$$

Using the Euclidean algorithm, any polynomial q can be written in the form $q = ap + r$, where a, r are polynomials and r has degree less than n . In the equivalence classes of $L^2(\mu)$, we therefore have that $q = r$. Hence, the dimension of $L^2(\mu)$ is n . $\diamond$

REMARK 6.7 ([JÓ00]). We can describe another way to generate the Hilbert space completion $\mathcal{H}_Q$ of a quadratic form Q . So far we have looked at two ways to generate isomorphic Hilbert spaces $\mathcal{H}_Q$. The first is to complete the finite sequences $\mathcal{D}$ with respect to the quadratic form Q_M . The second is the closure of the polynomials in $L^2(\mu)$, where μ is the measure whose moments are the entries of M . Here, we briefly outline a third method to generate the same Hilbert space which consists of distributions. For now, we will restrict ourselves to the concrete case $M =$ the Hilbert matrix.

As in our first method, we start with the positive semidefinite quadratic form generated by the moment matrix M :

$$Q_M : [0, 1) \times [0, 1) \rightarrow \mathbb{R}.$$

We use M to define linear functionals. For each x in $[0, 1)$, there is a linear functional v_x defined by

$$(6.9) \quad v_x(\cdot) = Q_M(\cdot, x).$$

(The analog to v_x is a sequence in $\mathcal{D}$ with zeros in every position except one, where there is a 1.) We use the v_x s to build more linear functionals with finite sums (here, the analog is $\mathcal{D}$):

$$(6.10) \quad u(\cdot) = \sum_{x \text{ finite}} c_x v_x(\cdot) = \sum_{x \text{ finite}} Q_M(\cdot, x).$$

Finally, there is an inner product in the space of linear functionals of the form (6.10):

$$(6.11) \quad \left\langle \sum_{x \text{ finite}} c_x v_x, \sum_{y \text{ finite}} c_y v_y \right\rangle = \sum_x \sum_y \overline{c_x} c_y Q_M(x, y).$$

Now, we complete the inner product space to the Hilbert space $\tilde{\mathcal{H}}$. The Hilbert space $\tilde{\mathcal{H}}$ is a reproducing kernel Hilbert space. It is fairly easy to check that for v_x as in (6.9) and u as in (6.10), $\langle v_x, u \rangle_{\tilde{\mathcal{H}}} = u(x)$.

Recall that the Hilbert matrix is related to the integral kernel $(1 - xy)^{-1}$. We will next explain how this kernel appears in the Hilbert space of distributions $\tilde{\mathcal{H}}$.

The tensor product of two distributions $\overline{u_1}$ and u_2 is straightforward; we apply the linear functional $\overline{u_1} \otimes u_2$ to the pair of functions (f, g) by

$$(6.12) \quad (\overline{u_1} \otimes u_2)(f(x), g(y)) = \overline{u_1}(f) u_2(g).$$

Then, we can extend this definition to functions $h(x, y)$ because the span of functions of the form $f(x)g(y)$ is dense in $L^2[0, 1]$.

One important example of a tensor product of distributions which is related to Widom's theorem is the following. Suppose ϕ_1 and ϕ_2 are locally integrable functions on $[0, 1)$, and $u_1 = \phi_1 dx$ and $u_2 = \phi_2 dx$. Then

$$(6.13) \quad (\overline{u_1} \otimes u_2) \left(\frac{1}{1 - xy} \right) = \int_0^1 \int_0^1 \frac{\overline{\phi_1}(x) \phi_2(y)}{1 - xy} dx dy.$$

The Hilbert space $\tilde{\mathcal{H}}$ is spanned by distributions u on such that

$$(\bar{u} \otimes u) \left(\frac{1}{1 - xy} \right) < \infty.$$

It can be shown that the Dirac mass at 0 and all its derivatives belong to $\tilde{\mathcal{H}}$; moreover, derivatives of the Dirac mass at 0 (when scaled appropriately) form an orthonormal basis for $\tilde{\mathcal{H}}$ with respect to the inner product (6.11). The specific ONB is

$$u_n = \frac{(-1)^n}{n!} \delta_0^{(n)}, n = 0, 1, 2, \dots$$

CHAPTER 7

Boundedness and spectral properties

In cases such as the Hilbert matrix from Section 6.1, the operators F and F^* (which might be weighted or unweighted) are bounded operators, although in general they will be unbounded densely defined operators. In the first part of this chapter, we give sufficient conditions such that these operators are bounded operators, and hence the Kato operator F^*F is also bounded. In Section 7.3, we use the theory of projection-valued measures described in Section 7.2 to analyze connections among the spectrum of the Kato operator for a moment matrix $M^{(\mu)}$, the measure μ itself, and the associated integral operator. We will demonstrate these connections via the examples in Section 7.5. We define the rank of a measure in Section 7.4 and demonstrate with examples which are convex combinations of Dirac measures.

7.1. Bounded Kato operators

Given a measure μ with finite moments of all orders, denote its moment matrix by $M = M^{(\mu)}$. We will use the generating function $G_\mu(x)$ having the even moments $M_{k,k}$ as coefficients to express conditions which ensure the operator F_w^* on the weighted space $\ell^2(w)$ is bounded.

PROPOSITION 7.1. *Let μ be a measure with compact support on $\mathbb{R}$ with moments of all orders. Let the generating function*

$$G_\mu(x) = \sum_{k=0}^{\infty} M_{k,k} x^k$$

have a finite and positive radius of convergence R . Select t such that $t > R$ and $\frac{1}{t} < R$. If we choose weights $w_k := t^k$, then F_w^ is a bounded operator with*

$$\|F_w^*\|_{op} \leq \left(G_\mu \left(\frac{1}{t} \right) \right)^{1/2}.$$

PROOF. Note that since $\frac{1}{t} < R$, then $\frac{1}{t}$ is within the radius of convergence, hence

$$G_\mu \left(\frac{1}{t} \right) := \sum_{k=0}^{\infty} \frac{M_{k,k}}{t^k} < \infty.$$

We take $w_k := t^k$ for our weights.

Now,

$$(7.1) \quad (F_w^* \phi)_k = \frac{1}{w_k} \int x^k \phi(x) d\mu(x), k \in \mathbb{N}_0.$$

So

$$\left| \int x^k \phi(x) d\mu(x) \right|^2 \leq M_{2k} \|\phi\|_{L^2(\mu)}^2$$

and

$$\begin{aligned} \|F_w^* \phi\|_{\ell^2(w)}^2 &= \sum_{k=0}^{\infty} \frac{1}{w_k} \left| \int x^k \phi(x) d\mu(x) \right|^2 \\ (7.2) \quad &\leq \sum_{k=0}^{\infty} \frac{1}{w_k} M_{2k} \|\phi\|_{L^2(\mu)}^2 \\ &= G_\mu \left(\frac{1}{t} \right) \|\phi\|_{L^2(\mu)}^2. \end{aligned}$$

Therefore, F_w^* is a bounded operator with

$$\|F_w^*\|_{\ell^2(w) \rightarrow L^2(\mu)}^2 = \sup_{\|\phi\|=1} \|F_w^* \phi\|_{\ell^2(w)}^2 \leq G_\mu \left(\frac{1}{t} \right).$$

Using the properties of the adjoints of bounded operators, we also can conclude that F , $F_w^* F$ and FF_w^* are bounded operators with norms given by

$$\|F_w^*\|_{op}^2 = \|F\|_{op}^2 = \|F_w^* F\|_{op} = \|F F_w^*\|_{op} \leq G_\mu \left(\frac{1}{t} \right).$$

□

LEMMA 7.2. *Suppose μ is a measure on $\mathbb{R}$ with compact support and finite moments of all orders. Suppose in addition that $\text{supp}(\mu) \subset [-1, 1]$, with $d\mu(x) = B(x) dx$, where B is a nonnegative bounded function. Then the operators F , F^* , $F^* F$ and FF^* are bounded, and*

$$\|F^*\|_{op}^2 = \|F\|_{op}^2 = \|F^* F\|_{op} = \|FF^*\|_{op} \leq \pi \|B\|_\infty.$$

PROOF. By [Wid66] (a straightforward generalization of Theorem 6.2), we know FF^* is an integral operator of the form

$$(FF^* \phi)(x) = \int_{-1}^1 \frac{\phi(y)}{1 - xy} d\mu(y) = \int_{-1}^1 \frac{\phi(y) B(y)}{1 - xy} dy.$$

Let K be the positive integral operator defined in Section 6.1 which is equivalent to the moment matrix for Lebesgue measure. Then

$$\begin{aligned} \langle \phi | FF^* \phi \rangle_{L^2(\mu)} &= \int_{-1}^1 \int_{-1}^1 \frac{\overline{\phi(x)} B(x) \phi(y) B(y)}{1 - xy} dy dx \\ &= \langle \phi B | K \phi B \rangle_{L^2[-1,1]} \\ &= \langle K^{1/2} \phi B | K^{1/2} \phi B \rangle_{L^2[-1,1]} \\ &\leq \|K^{1/2}\|_{op} \|\phi B\|_{L^2[-1,1]} \\ &\leq \pi \int_{-1}^1 |\phi(x) B(x)|^2 dx. \end{aligned}$$

We recall that the operator norm of K is π . In turn, the last expression is bounded above by

$$\pi \|B\|_\infty \int_{-1}^1 |\phi(x)|^2 B(x) dx = \pi \|B\|_\infty \|\phi\|_{L^2(\mu)}^2.$$

Therefore, $\|FF^*\|_{op} \leq \pi\|B\|_\infty$, and $\|F\| = \|F^*\| \leq \sqrt{\pi\|B\|_\infty}$. $\square$

THEOREM 7.3. *Suppose there exists a finite $t > 1$ such that $\text{supp}(\mu) \subset [-t, t]$, μ has finite moments of all orders, and $d\mu(x) = B(x) dx$, with B a bounded non-negative function. Define weights $w = \{w_k\}_{k=0}^\infty$ where $w_k := t^{2k}$ on $\ell^2(w)$. Then the operators F , F^* , F_w^*F , FF_w^* are bounded, and the operator norm for F_w^*F and FF_w^* is bounded above by $t\pi\|B\|_\infty$.*

PROOF. For each $k \in \mathbb{N}_0$,

$$\begin{aligned} (F_w^*\phi)_k &= \frac{1}{w_k} \int_{-t}^t x^k \phi(x) d\mu(x) \\ &= \frac{1}{t^{2k}} \int_{-t}^t x^k \phi(x) B(x) dx \\ &= \frac{t}{t^k} \int_{-1}^1 u^k \phi(tu) B(tu) du, \end{aligned}$$

and

$$\begin{aligned} \|F_w^*\phi\|_{\ell^2(w)}^2 &= \sum_k \frac{1}{w_k} \left| \int_{-t}^t x^k \phi(x) B(x) dx \right|^2 \\ &= t^2 \sum_k \left| \int_{-1}^1 u^k \phi(tu) B(tu) du \right|^2. \end{aligned}$$

But by the previous lemma, this last expression is less than or equal to

$$\begin{aligned} &t^2 \pi \|B\|_\infty \int_{-1}^1 |\phi(tu)|^2 B(tu) du \\ &= t\pi \|B\|_\infty \int_{-t}^t |\phi(x)|^2 B(x) dx \\ &= t\pi \|B\|_\infty \|\phi\|_{L^2(\mu)}^2. \end{aligned}$$

So, $\|F_w^*\|_{op}^2 = \|FF_w^*\|_{op} \leq t\pi\|B\|_\infty$. $\square$

The following is an immediate result of Theorem 7.3.

COROLLARY 7.4. *Suppose there exists a finite $t > 1$ such that $\text{supp}(\mu) \subset [-t, t]$, μ has finite moments of all orders, and $d\mu(x) = B(x) dx$, with B a nonnegative bounded function. Define weights $w = \{w_k\}_{k=0}^\infty$ where $w_k := t^{2k}$ on $\ell^2(w)$. The operator FF_w^* is an integral operator of the form*

$$(FF_w^*\phi)(x) = \int_{-t}^t \frac{1}{1 - \frac{xy}{t^2}} \phi(y) B(y) dy.$$

We can use Theorem 7.3 to study operators from the examples in Chapter 2 where the moments were related to the Catalan numbers.

COROLLARY 7.5. *Suppose μ is the Wigner semicircle measure in Example 2.13. Then under the weights w from Corollary 7.4, FF_w^* is bounded, with $\|FF_w^*\|_{op} \leq 4\pi$.*

We would also like to apply Theorem 7.3 to Example 2.14, but we cannot since the function B is not in L^∞ . Still, we can provide an estimate. We use a generating function G defined by

$$(7.3) \quad G(\zeta) = \sum_{k \in \mathbb{N}_0} \zeta^k \int |x|^k d\mu(x) = \sum \zeta^k \mathcal{M}_k,$$

where we recall the use of the notation $\mathcal{M}_k$ from Lemma 4.4 in Section 4.1.

LEMMA 7.6. *Suppose $\text{supp}(\mu) \subset [-t, t]$, where μ has finite moments of all orders. Then the radius of convergence R of the generating function G from Equation (7.3) satisfies $R \geq \frac{1}{t}$.*

PROOF. We have

$$\left(\int_{-t}^t |x|^k d\mu \right)^{1/k} \leq t,$$

where we use the fact that $t := \text{ess sup}|x|$ on $L^\infty(\mu)$ and also the fact that μ is a probability measure. Therefore, we are assured that G converges absolutely for $|\xi|t < 1$. Therefore, $R \geq \frac{1}{t}$. $\square$

Pick $w_k := s^k$ for some $t^2 < s < \infty$. We can now establish an estimate for $\|F^* \phi\|_{\ell^2(w)}$ in terms of the generating function G :

$$\begin{aligned} \|F_w^* \phi\|_{\ell^2(w)}^2 &= \sum_k \frac{1}{s^k} \left| \int x^k \phi(x) d\mu(x) \right|^2 \\ &\leq \sum_k \frac{1}{s^k} \left(\int_{-t}^t |x|^k |\phi(x)| d\mu(x) \right)^2. \end{aligned}$$

Then, using Cauchy-Schwarz, we have

$$\begin{aligned} \|F_w^* \phi\|_{\ell^2(w)}^2 &\leq \sum_k \frac{1}{s^k} \mathcal{M}_k \int |x|^k |\phi(x)|^2 d\mu(x) \\ &= \int_{-t}^t G\left(\frac{|x|}{s}\right) |\phi(x)|^2 d\mu(x). \end{aligned}$$

The switch of the sum and integral above is justified by a Fubini argument, since the power series $G(x)$ is continuous.

We apply this observation to Example 2.14, the secant measure. In this case, $G(x) = 2(\sqrt{1-4x^2})^{-1}$ and $t = 2$. Choose $s > 4$ and set $w = \{w_k\}_{k=0}^\infty$ where $w_k = s^k$. Then in $\ell^2(w)$, we have

$$\|F_w^* \phi\|_{\ell^2(w)}^2 \leq \int_{-2}^2 \frac{2}{\sqrt{1 - (\frac{2x}{s})^2}} |\phi(x)|^2 d\mu(x).$$

We next describe a result related to affine iterated function system measures. In particular, we show a sufficient condition under which the matrix A (see Section 4.1) encoding an affine map τ is a Hilbert-Schmidt operator.

PROPOSITION 7.7. Consider the transformation $\tau(z) = cx + b$, and suppose $|c| \leq |b| < \frac{1}{4}$. Then the operator defined by the matrix A which satisfies

$$M^{(\mu \circ \tau)} = A^* M^{(\mu)} A,$$

with is Hilbert-Schmidt; i.e. A is bounded, and $\text{trace}(A^* A) < \infty$.

PROOF. Let k denote the row index in A , and let j denote the column index in A ; as before (3.5), we have

$$A_{k,j} = \begin{cases} \binom{k}{j} c^k b^{j-k} & 0 \leq k \leq j \\ 0 & k > j \end{cases}.$$

The $(i, j)^{\text{th}}$ entry of the infinite matrix $A^* A$ is therefore

$$(A^* A)_{i,j} = \sum_{0 \leq k \leq i \wedge j} \overline{A_{k,i}} A_{k,j} = \sum_{0 \leq k \leq i \wedge j} \binom{i}{k} \binom{j}{k} |c^k|^2 \overline{b^{i-k}} b^{j-k},$$

and if $i = j$, the diagonal entries are

$$(7.4) \quad (A^* A)_{j,j} = \sum_{k=0}^j \binom{j}{k}^2 |c^k|^2 |b^{j-k}|^2.$$

Set $\alpha := |c|^2$ and $\beta := |b|^2$. Then, using the assumption that $\frac{\alpha}{\beta} \leq 1$, we have

$$(7.5) \quad \begin{aligned} \text{trace}(A^* A) &= \sum_{j=0}^{\infty} (A^* A)_{j,j} = \sum_{j=0}^{\infty} \sum_{k=0}^j \binom{j}{k}^2 \alpha^k \beta^{j-k} \\ &\leq \sum_{j=0}^{\infty} \beta^j \sum_{k=0}^j \binom{j}{k}^2 = \sum_{j=0}^{\infty} \beta^j \binom{2j}{j} \end{aligned}$$

The last series has radius of convergence $\frac{1}{4}$. □

7.2. Projection-valued measures

This section provides an overview of the properties of projection-valued measures, leading to the spectral decomposition for unbounded self-adjoint operators on a Hilbert space $\mathcal{H}$. These ideas will be used in Section 7.3 and later in Chapter 8 in order to discuss the Kato-Friedrichs extension of operators. For far more detailed treatment of this theory, we refer the reader to [Bag92, RS80, Rud91].

Recall from Definition 1.4 that an operator H with dense domain $\text{dom}(H)$ in a Hilbert space $\mathcal{H}$ is said to be *self adjoint* if $H^* = H$ and $\text{dom}(H^*) = \text{dom}(H)$.

DEFINITION 7.8. A Borel *projection-valued measure* E on $\mathbb{R}$ is a map from the σ -algebra $\mathcal{B}$ of Borel subsets of $\mathbb{R}$ to the orthogonal projections on a Hilbert space $\mathcal{H}$ such that

- (i) $E(\emptyset) = 0$,
- (ii) if $S = \cup_{i=1}^{\infty} S_i$ and $S_i \cap S_j = \emptyset$ for $S_i \in \mathcal{B}$ and $i \neq j$, then

$$E(S) = \sum_{i=0}^{\infty} E(S_i) = \lim_{i \rightarrow \infty} \sum_{j=1}^i E(S_j),$$

where the limit is in the strong operator topology.

A projection-valued measure E is said to be *orthogonal* if

$$(7.6) \quad E(S_1 \cap S_2) = E(S_1)E(S_2)$$

for all Borel sets S_1, S_2 . We will assume all projection-valued measures are orthogonal unless otherwise stated.

Recall from the definition of an orthogonal projection that for every Borel set S , we have $E(S) = E(S)^* = E(S)^2$.

There is a well-known theory which develops the notion of integration of measurable real-valued functions against a projection-valued measure E , resulting in an operator on the Hilbert space $\mathcal{H}$. We will denote such an integral by $\int_{\mathbb{R}} f(\lambda)E(d\lambda)$. Quite naturally, we start by defining integrals of characteristic functions,

$$E(S) = \int_{\mathbb{R}} \chi_S(\lambda)E(d\lambda),$$

then extend appropriately to measurable functions. The operator is a bounded operator if and only if there exists an $M > 0$ such that $E(|f|^{-1}(M, \infty)) = 0$ (the trivial projection). We say that E is a *resolution of the identity* if

$$(7.7) \quad I_{\mathcal{H}} = \int_{\mathbb{R}} E(d\lambda) = E(\mathbb{R}).$$

EXAMPLE 7.9. [**Bag92**] Given μ a σ -finite Borel measure on $\mathbb{R}$, let $\mathcal{H} = L^2(\mu)$. Given a Borel set $S \subseteq \mathbb{R}$, define $E(S)$ to be a multiplication operator which multiplies by the characteristic function χ_S :

$$E(S)(f) = \chi_S f.$$

Then E is a projection-valued measure on $\mathbb{R}$, called the *canonical projection-valued measure*. Given f a measurable real-valued function on $\mathbb{R}$, $\int_{\mathbb{R}} f(\lambda)E(d\lambda)$ has domain $\{g \in \mathcal{H} : fg \in \mathcal{H}\}$ and on that domain,

$$\int_{\mathbb{R}} f(\lambda)E(d\lambda)(g) = fg.$$

◇

One useful property of projection-valued measures is that integration is multiplicative, where the product of operators is composition.

$$\int f(\lambda)g(\lambda)E(d\lambda) = \int f(\lambda)E(d\lambda) \int g(\lambda)E(d\lambda)$$

Corresponding to a projection-valued measure E , there is a parameterized family of real-valued measures $\{\nu_v\}_{v \in \mathcal{H}}$ given by the $\mathcal{H}$ -inner product:

$$(7.8) \quad \nu_v(S) = \langle v | E(S)v \rangle.$$

Since orthogonal projections are positive operators, the quantity $\langle v | E(S)v \rangle$ is non-negative for any choice of v and any set S . Given a projection-valued measure which is a resolution of the identity, we have

$$\int_{\mathbb{R}} d\nu_v(\lambda) = \langle v | E(\mathbb{R})v \rangle = \|v\|^2.$$

The following theorem gives a spectral decomposition for self-adjoint operators.

THEOREM 7.10 ([DS88, Rud91]). *An operator H in a Hilbert space $\mathcal{H}$ is self adjoint if and only if there exists an orthogonal projection-valued measure E supported on the spectrum $\sigma(H)$ such that*

$$(7.9) \quad Hv = \left[\int_{\mathbb{R}} \lambda E(d\lambda) \right] v = \left[\int_{\sigma(H)} \lambda E(d\lambda) \right] v$$

for all $v \in \text{dom}(H)$. Moreover, $v \in \text{dom}(H)$ if and only if

$$(7.10) \quad \|Hv\|^2 = \int_{\sigma(H)} \lambda^2 d\nu_v(\lambda) < \infty,$$

where the measure ν_v is defined in Equation (7.8).

While we leave the details of the proof to the cited literature, it may be helpful here to include the computation which demonstrates the equality given in Equation (7.10). Given $v \in \text{dom}(H)$,

$$\begin{aligned} \left\| \int \lambda E(d\lambda) v \right\|^2 &= \left\langle \int \lambda E(d\lambda) v \middle| \int \lambda E(d\lambda) v \right\rangle \\ &= \left\langle \left[\int \lambda E(d\lambda) \right]^* \left[\int \lambda E(d\lambda) \right] v \middle| v \right\rangle \\ &= \left\langle \left[\int \lambda E(d\lambda) \right]^2 v \middle| v \right\rangle \quad \text{since the op. is self adjoint} \\ &= \left\langle \left[\int \lambda^2 E(d\lambda) \right] v \middle| v \right\rangle \quad \text{multiplicative prop. of integrals} \\ &= \int \lambda^2 d\nu_v(\lambda). \end{aligned}$$

□

The formulas from Theorem 7.10 together help us define the domains of (possibly unbounded) self-adjoint operators defined by integrals of measurable real-valued functions f against a projection-valued measure E . We say that v is in the domain of an operator $H = \int f(\lambda) E(d\lambda)$ if

$$(7.11) \quad \left\| Hv - \left[\int_{-n}^n f(\lambda) E(d\lambda) \right] v \right\| \rightarrow 0$$

as $n \rightarrow \infty$. If H is a self-adjoint operator, we can use the equality in (7.10) which gives

$$\left\| Hv - \left[\int_{-n}^n \lambda E(d\lambda) \right] v \right\|^2 = \int_{|\lambda| > n} \lambda^2 d\nu_v(\lambda).$$

Therefore, a given vector $v \in \mathcal{H}$ is in $\text{dom}(H)$ if and only if

$$(7.12) \quad \lim_{n \rightarrow \infty} \int_n^\infty \lambda^2 d\nu_v(\lambda) = 0.$$

7.3. Spectrum of the Kato operator

In this section, we show how spectral analysis of the Kato-Friedrichs operator translates into spectral data for the initially given moment matrix $M^{(\mu)}$ and the measure μ . We then illustrate the theory in several examples. Our main result (Theorem 7.11) is that atoms in the spectrum of the Kato-Friedrichs operator pair

up with atoms in the spectrum of the measure μ itself. Using this we arrive at our own proof of the well-known fact (Theorem 7.13) that the Hilbert matrix has continuous spectrum.

In fact, in their spectral picture, the Kato-Friedrichs operators are directly related to the associated moment matrices $M^{(\mu)}$; this correspondence is especially transparent for Examples 7.19 and 7.20. In these examples, the agreement of the rank of the Kato-Friedrichs operators with the rank of the measure, which we define in Section 7.4, can be seen by inspection.

THEOREM 7.11. *Let μ be a measure with compact support on the real line. Then the atoms in the spectrum of the Kato-Friedrichs operator of the moment matrix $M = M^{(\mu)}$ (if any) pair up with atoms in the spectrum of the measure μ itself.*

Before we begin the proof, we will a few key points about spectral resolutions and operator theory.

Let H be a densely defined, positive self-adjoint operator on the Hilbert space $\mathcal{H}$. Then there exists (unique, up to unitary equivalence) a projection-valued measure E such that

$$I_{\mathcal{H}} = \int E(d\lambda)$$

and

$$H = \int_{\sigma(H)} \lambda E(d\lambda).$$

In addition, for any $x \in \mathcal{H}$, given the spectral measure ν_x defined in as in Equation (7.8), we have

$$\|Hx\|^2 = \int_{\sigma(H)} \lambda^2 d\nu_x.$$

DEFINITION 7.12. An *atom* in H or in E is a point $\lambda_1 \in \mathbb{R}$ such that $E(\{\lambda_1\}) \neq 0$, i.e. is not the trivial projection.

If $\mathcal{H}$ is a complex Hilbert space, then we can define an order on self-adjoint operators on $\mathcal{H}$:

$$H \leq K \text{ if and only if } \langle x|Hx \rangle \leq \langle x|Kx \rangle \text{ for all } x \in \mathcal{H}.$$

With this order, we observe that if E_1 and E_2 are projections on $\mathcal{H}$, then

$$E_1 \leq E_2 \Leftrightarrow E_1 = E_1 E_2 = E_2 E_1.$$

In this case, we say that E_1 is a subprojection of E_2 .

PROOF. Let $\text{supp}(\mu) \subset \mathbb{R}$, where the measure μ has compact support. Denote the Kato-Friedrichs operator for the moment matrix $M = M^{(\mu)}$ by H . Let E be the projection-valued measure for H . Recall that in the unweighted case, the mapping $F^* : L^2(\mu) \rightarrow \ell^2$ which takes $f_c(x) = \sum_{i \in \mathbb{N}_0} c_i x^i$ to $\{c_i\}_{i \in \mathbb{N}_0}$ is an isometry.

Next, assume that λ_1 is an atom in the spectrum of H . Therefore, $E(\lambda_1)$ is a nontrivial projection, hence there exists $\xi_1 \in \text{Range}(E(\lambda_1))$, $\|\xi_1\|_{\ell^2} = 1$. We denote the rank-one projection onto ξ_1 (using Dirac notation; see Chapter 1) by

$$(7.13) \quad E_{\lambda_1} = |\xi_1\rangle\langle\xi_1|.$$

H is a positive operator, hence $\lambda_1 > 0$. Since H is the Kato-Friedrichs operator, we have

$$\int |f_c|^2 d\mu = \|H^{1/2}c\|_{\ell^2}^2$$

for all $c \in \mathcal{D}$. Therefore, there exists $A_1 \in \mathbb{R}^+$ such that

$$(7.14) \quad A_1 E_{\lambda_1} \leq H.$$

Equivalently, by the idempotent property of projections, for all $c \in \mathcal{D}$ we have

$$(7.15) \quad A_1 \|E_{\lambda_1} c\|_{\ell^2}^2 \leq \|H^{1/2}c\|_{\ell^2}^2 = \int |f_c|^2 d\mu.$$

Recall that for all $c \in \ell^2$,

$$(7.16) \quad \langle c | E_{\lambda_1} c \rangle_{\ell^2} = \|E_{\lambda_1} c\|_{\ell^2}^2 = |\langle \xi_1 | c \rangle_{\ell^2}|^2.$$

We now make use of the Hankel property from Definition 1.6 in the moment matrix M . Let

$$S : \{c_0, c_1, c_2, \dots\} \rightarrow \{0, c_0, c_1, \dots\}$$

be the right shift operator in ℓ^2 . Then the Hankel condition is equivalent on $\mathcal{D}$ to

$$(7.17) \quad HS = S^*H.$$

By the spectral theorem, then

$$(7.18) \quad E_{\lambda_1} S = S^* E_{\lambda_1}.$$

Looking ahead to the argument in Lemma 8.8, we see that this implies that the vector ξ_1 is of the form

$$\xi_1 = [1 \quad b \quad b^2 \quad b^3 \quad \dots]^{tr}.$$

Equation (7.15) gives that for all $c \in \mathcal{D}$,

$$(7.19) \quad A_1 \|E_{\lambda_1} c\|_{\ell^2}^2 = A_1 \sum_{i=0}^{\infty} \left| \sum_{j=0}^{\infty} \overline{\xi(i)} \xi(j) c_j \right|^2 = A_1 \sum_{i=0}^{\infty} \left| \sum_{j=0}^{\infty} b^{i+j} c_j \right|^2 \leq \int |f_c|^2 d\mu.$$

Using Example 7.16 regarding the Dirac point-mass measure δ_b at the point b , we have

$$A_1 \int |f_c|^2 d\delta_b = A_1 |f_c(b)|^2 = A_1 \left| \sum_{j=0}^{\infty} c_j b^j \right|^2.$$

Since the expression above is equal to the $i = 0$ term from Equation (7.19), we also have

$$(7.20) \quad A_1 \int |f_c|^2 d\delta_b \leq \int |f_c|^2 d\mu \text{ for all } c \in \mathcal{D}.$$

Since the support of μ is compact, the set $\{f_c\}_{c \in \mathcal{D}}$ is dense in $L^2(\mu)$. Therefore, for all Borel sets $E \in \mathcal{B}(\mathbb{R})$ there exists a sequence $\{c_k\}_{k \in \mathbb{N}_0} \subset \mathcal{D}$ such that

$$f_{c_k}(x) \xrightarrow{L^2(\mu)} \chi_E(x).$$

Therefore, using the sequence $\{f_{c_k}\}$ to approximate χ_E , we get

$$(7.21) \quad A_1 \int_E d\delta_b \leq \int_E d\mu,$$

or equivalently,

$$(7.22) \quad A_1 \delta_b(E) \leq \mu(E).$$

Since $\delta_b(\{b\}) = 1$, it must be true that $\mu(\{b\}) > 0$. Therefore, when the Kato-Friedrichs operator H has an atom, the measure μ has a corresponding atom. $\square$

The Kato-Friedrichs operator for the moment matrix for Lebesgue measure on the interval $[0, 1]$ is the bounded operator defined relative to the standard ONB in ℓ^2 by the Hilbert matrix. In particular, we need not introduce weights into the ℓ^2 space in order to produce a closable quadratic form. This is a case when the Kato-Friedrichs operator has infinite rank (see Section 7.4). We now see that the well-known fact that the spectrum of this operator is purely continuous follows as an immediate corollary of Theorem 7.11. (This result was first shown in [Mag50].)

THEOREM 7.13. *The spectrum of the Hilbert matrix is continuous, i.e., there are no atoms in the spectral resolution of the Hilbert matrix.*

PROOF. We saw that the action of the Hilbert matrix M on ℓ^2 defines a bounded (see [Hal67]) self-adjoint operator, and M is the moment matrix $M^{(\mu)}$ where μ is taken to be Lebesgue measure on the unit interval $(0, 1)$. Moreover, the quadratic forms below coincide on the space $\mathcal{D}$ of finite vectors in ℓ^2 :

$$(7.23) \quad \|f_c\|_{L^2(\mu)} = Q_M(c) = \langle c | M c \rangle_{\ell^2} \text{ for all } c \in \mathcal{D}.$$

Therefore, M is the Kato-Friedrichs operator for the quadratic form Q_M .

If $M = M^{(\mu)}$ contained an atom, then there would be a rank-one projection in the spectral resolution for $M^{(\mu)}$, and by Theorem 7.11 this would imply that Lebesgue measure restricted to $(0, 1)$ would contain an atom. That is a clear contradiction, and therefore M has continuous spectrum. $\square$

In Section 7.5, we explicitly compute the Kato-Friedrichs operators in several examples and study their spectra. Recall that in general, the Kato-Friedrichs operator H is a self-adjoint operator on $\ell^2(w)$ for some set of weights $w = \{w_k\}_{k \in \mathbb{N}_0}$. Note that the operator $H(= H_w)$ depends on the choice of w , and the choice of w depends on the measure μ . The lemma below demonstrates how the weights in a Hilbert space $\ell^2(w)$ arise in the equations governing the point spectrum of the Kato-Friedrichs operator.

Recall our notation for the weighted Hilbert spaces, where $w = \{w_i\}_{i \in \mathbb{N}_0}$ are the weights:

$$\ell^2(w) = \left\{ c = \{c_i\}_{i \in \mathbb{N}_0} \left| \sum_{i \in \mathbb{N}_0} w_i |c_i|^2 < \infty \right. \right\}$$

and

$$\langle b | c \rangle_{\ell^2(w)} := \sum_{i \in \mathbb{N}_0} w_i \bar{b}_i c_i.$$

In the following computations, we assume that

- (1) the measure μ is a positive Borel measure on $\mathbb{R}$ with compact support having moments of all orders,
- (2) the weights $\{w_i\}_{i \in \mathbb{N}_0}$ satisfy $w_i > 0$ for all $i \in \mathbb{N}_0$,
- (3) the monomials x^i are in $L^2(\mu) \cap L^1(\mu)$ for all $i \in \mathbb{N}_0$.

LEMMA 7.14. *Let μ be a Borel measure on $\mathbb{R}$ with compact support and finite moments of all orders. Let $M = M^{(\mu)}$ be the moment matrix for μ , and let H be the Kato-Friedrichs operator for M under an appropriate choice of weights $w = \{w_i\}_{i \in \mathbb{N}_0}$. There exists λ in the point spectrum of H , i.e.*

$$Hc = \lambda c$$

for some nonzero c in the domain of H , if and only if for each $i \in \mathbb{N}_0$,

$$(7.24) \quad \sum_{j \in \mathbb{N}_0} M_{i,j} c_j = \lambda w_i c_i.$$

PROOF OF LEMMA 7.14: The (nonnegative) real number λ is in the point spectrum of H if and only if there is a nonzero c in the domain of H such that $Hc = \lambda c$. By definition of the Kato-Friedrichs operator, H is related to the quadratic form Q_M arising from M via

$$Q_M(c) = \sum_j \sum_k \bar{c}_j M_{j,k} c_k = \|H^{1/2} c\|_{\ell^2(w)}^2, \quad \forall c \in \mathcal{D}.$$

When we pass to the completion, given $c \in \text{dom}(H^{1/2})$ and $f_c \in L^2(\mu)$, we have the identity

$$\int |f_c|^2 d\mu = \|H^{1/2} c\|_{\ell^2(w)}^2.$$

Recall that $f_c \in L^2(\mu)$ if and only if $c \in \text{dom}(H^{1/2})$; this follows from the completion in the Kato-Friedrichs construction.

There is also a sesquilinear form S_M determined by M (connected to Q_M by the polarization identity) which satisfies

$$(7.25) \quad S_M(b, c) = \sum_j \sum_k \bar{b}_j M_{j,k} c_k = \langle b | Hc \rangle_{\ell^2(w)}$$

for all $b \in \mathcal{D}, c \in \text{dom}(H)$.

For each $i \in \mathbb{N}_0$, let $b = e_i$, the i -th standard basis vector. Equation (7.25) becomes

$$\sum_j M_{i,j} c_j = \langle e_i | Hc \rangle_{\ell^2(w)} = w_i (Hc)_i.$$

We now have that $Hc = \lambda c$ for some nonzero vector c in the domain of H if and only if

$$\sum_j M_{i,j} c_j = w_i (Hc)_i = w_i \lambda c_i.$$

□

In Equations (7.24), we will seek solutions for λ , w , and c . We require

$$c \in \ell^2(w) \quad \text{and} \quad \|c\|_{\ell^2(w)} = 1,$$

i.e.,

$$(7.26) \quad \sum_{i \in \mathbb{N}_0} w_i |c_i|^2 = 1.$$

In the examples shown in Section 7.5, we will find that it is relatively easy to solve (7.26) for a single Dirac mass $\mu = \delta_b$ for any $b \in \mathbb{R}$. However, in the case of convex combinations of point masses, finding solutions to (7.24) and (7.26) is more tricky.

7.4. Rank of measures

Consider a probability measure μ on $\mathbb{C}$ or on $\mathbb{R}^d$, and assume that μ has finite moments of all orders. Pick an order of the index set for the monomials, and let $\{p_k\}_{k \in \mathbb{N}_0^d}$ be the associated orthogonal polynomials in $L^2(\mu)$ defined from μ ; further, let G be the corresponding Gram matrix. Let $\mathcal{P}$ be the closed subspace in $L^2(\mu)$ spanned by the monomials.

DEFINITION 7.15. We say that the *rank of G* , and hence the *rank of the measure μ* , is the dimension of the Hilbert space $\mathcal{P}$.

We note two things in Definition 7.15. First, $\{p_k\}_{k \in \mathbb{N}_0^d}$ is an ONB in $\mathcal{P}$. Second, $\mathcal{P} = L^2(\mu)$ if and only if the monomials are dense in $L^2(\mu)$; this is known to be true if μ has compact support, but it need not be true in general. We will see this more in Chapter 8.

EXAMPLE 7.16. A Dirac measure has rank 1.

Let b be a fixed complex number, and let $\mu := \delta_b$ be the corresponding Dirac measure. Then the moments m_k of μ are

$$(7.27) \quad m_k = \int_{\mathbb{C}} z^k d\mu(z) = b^k, \quad k \in \mathbb{N}_0;$$

the moment matrix is

$$(7.28) \quad M_{j,k}^{(\mu)} = \bar{b}^j b^k.$$

If $\mathcal{D}$ is the space of finite sequences, $c = \{c_j\}_{j \in \mathbb{N}_0}$, then

$$(7.29) \quad \langle c | M^{(\mu)} c \rangle_{\ell^2} = \left| \sum_{j \in \mathbb{N}_0} c_j \bar{b}^j \right|^2.$$

We showed in Example 6.6 that $\dim(L^2(\mu)) = 1$, so it follows that $p_0(z) \equiv 1$ and $p_k \equiv 0$ for $k \geq 1$. We use Equations (4.2) and (4.6) together with Lemma 4.3 to compute the following formulas for the infinite matrices G and G^{-1} :

$$(7.30) \quad G = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix} \quad \text{and} \quad G^{-1} = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ b & 0 & 0 & \cdots \\ b^2 & 0 & 0 & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix}.$$

We see here that G has rank 1. Note that we are using the notion of inverse described in Section 1.6. We see that GG^{-1} is an idempotent rank-one matrix. A direct computation further yields

$$(7.31) \quad G^{-1}(G^{-1})^* = (b^j \bar{b}^k) = (M^{(\mu)})^{tr},$$

as predicted by Lemma 4.3 and Equation (4.14).

EXAMPLE 7.17. A measure with rank 2.

Let $\mu := \frac{1}{2}(\delta_0 + \delta_1)$. The first moment of μ is 1, and all the other moments are all $\frac{1}{2}$. The associated orthogonal polynomials are

$$(7.32) \quad p_0(z) \equiv 1, \quad p_1(z) = 2z - 1, \quad \text{and} \quad p_2(z) = p_3(z) = \dots \equiv 0.$$

We then compute the Gram matrix and an inverse (in the sense of Definition 1.16 and Example 1.18):

$$(7.33) \quad G = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ -1 & 2 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix} \quad \text{and} \quad G^{-1} = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & 0 & \cdots \\ \frac{1}{2} & \frac{1}{2} & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Again we can compute directly the two idempotents E_1 and E_2 .

$$(7.34) \quad E_1 = GG^{-1} = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ 0 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

and

$$(7.35) \quad E_2 = G^{-1}G = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}.$$

Note that E_1 is a projection; i.e., $E_1 = E_1^* = E_1^2$, while E_2 is only an idempotent; i.e., $E_2^2 = E_2$. Note that these rules are statements about infinite matrices. In fact, if E_2 is viewed as an operator in ℓ^2 , it is unbounded ($E_2(\mathbf{e}_2) \notin \ell^2$.) So,

$$\text{rank}(G) = \text{rank}(\mu) = \text{rank}(M^{(\mu)}) = \dim(L^2(\mu)) = 2.$$

◇

7.5. Examples

In Example 7.18, we use Lebesgue measure and the corresponding Hilbert matrix to illustrate the case where the moment matrix is a bounded operator. We also can demonstrate the IFS techniques from Chapter 3 here. Following that, we use the measures $\mu = \delta_1$ and $\mu = \frac{1}{2}(\delta_0 + \delta_1)$ to illustrate the spectral results from this chapter. In both cases, we find appropriate choices for the weights which allow us to find Kato-Friedrichs operators on the Hilbert space $\ell^2(w)$ space. Since the operator depends on the choice of weights, we will denote the Kato-Friedrichs operator by H_w . In Example 7.19, for a particular choice of w , we get the associated Kato-Friedrichs operators to be a rank-one projection. We similarly analyze the spectrum of the Kato-Friedrichs operator (which has rank 2) in Example 7.20.

EXAMPLE 7.18. Lebesgue measure and the Hilbert matrix.

Let μ be Lebesgue measure restricted to $[0, 1]$, hence the moment matrix $M^{(\mu)}$ is the Hilbert matrix previously discussed in Example 1.7 and Section 6.1. Recall that it is known ([Wid66]) that the Hilbert matrix represents a bounded operator on ℓ^2 with operator norm $\|M^{(\mu)}\|_{op} = \pi$.

It is well known that μ is exactly the equilibrium measure arising from the real affine IFS $\{\tau_i\}_{i=0,1}$ where

$$\tau_0(x) = \frac{x}{2} \text{ and } \tau_1(x) = \frac{x+1}{2},$$

together with equal weights $\frac{1}{2}$. The IFS invariance property for moment matrices is

$$(7.36) \quad M^{(\mu)} = \frac{1}{2} \left(A_0^* M^{(\mu)} A_0 + A_1^* M^{(\mu)} A_1 \right),$$

where Lemma 3.1 gives the matrices A_0 and A_1 (from [EST06]):

$$A_0 = \begin{bmatrix} 1 & 0 & 0 & \cdots \\ 0 & \frac{1}{2} & 0 & \cdots \\ 0 & 0 & \frac{1}{4} & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix} \quad A_1 = \begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{4} & \cdots \\ 0 & \frac{1}{2} & \frac{1}{2} & \cdots \\ 0 & 0 & \frac{1}{4} & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix}.$$

Since the infinite matrices A_0 and A_1 are upper triangular (and hence A_0^* and A_1^* are lower triangular), the matrix multiplication computations in (7.36) involve only *finite* sums. In addition, when we check the appropriate convergence issues, each of the matrices $A_i^* M^{(\mu)} A_i = M^{(\mu \circ \tau_i^{-1})}$, for $i = 0, 1$, represents a bounded operator on ℓ^2 . We compute directly $A_0^* M^{(\mu)} A_0$:

$$\begin{aligned} (A_0^* M^{(\mu)} A_0)_{i,j} &= \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} (A_0^*)_{i,k} M_{k,\ell}^{(\mu)} (A_0)_{\ell,j} \\ &= (A_0^*)_{i,i} \frac{1}{1+i+j} (A_0)_{j,j} \\ &= \left(\frac{1}{2^{i+j}} \right) \frac{1}{1+i+j} = M^{(\mu \circ \tau_0^{-1})}. \end{aligned}$$

The entries of the matrix $A_0 M^{(\mu)} A_0$ together with the IFS invariance (7.36) property

$$\begin{aligned} \frac{1}{2} \left(A_0^* M^{(\mu)} A_0 + A_1^* M^{(\mu)} A_1 \right)_{i,j} &= \frac{1}{2} \left[\left(\frac{1}{2^{i+j}} \right) \frac{1}{1+i+j} + \left(1 - \frac{1}{2^{i+j}} \right) \frac{1}{1+i+j} \right] \\ &= \frac{1}{1+i+j} = M_{i,j}^{(\mu)} \end{aligned}$$

allow us to compute the entries of the matrix $A_1^* M^{(\mu)} A_1$:

$$\begin{aligned} (A_1^* M^{(\mu)} A_1)_{i,j} &= \sum_{k=0}^{\infty} \sum_{\ell=0}^{\infty} (A_1^*)_{i,k} M_{k,\ell}^{(\mu)} (A_1)_{\ell,j} \\ (7.37) \quad &= \frac{1}{2^{i+j}} \sum_{k=0}^i \sum_{\ell=0}^j \binom{i}{k} \binom{j}{\ell} \frac{1}{1+k+\ell} \\ &= \left(1 - \frac{1}{2^{i+j}} \right) \frac{1}{1+i+j}. \end{aligned}$$

We note that (7.37) yields an interesting identity on binomial coefficients. Given $i, j \in \mathbb{N}_0$,

$$(7.38) \quad \sum_{k=0}^i \sum_{\ell=0}^j \binom{i}{k} \binom{j}{\ell} \frac{1}{1+k+\ell} = \frac{2^{i+j} - 1}{1+i+j}.$$

The infinite matrices A_0 and A_1 represent bounded operators on the Hilbert space ℓ^2 with

$$\|A_i\|_{op} = \|A_i^*\|_{op} = 1 \quad i = 0, 1.$$

These are therefore contractive operators on ℓ^2 . $\diamond$

EXAMPLE 7.19. The spectrum of the Kato-Friedrichs operator for $\mu = \delta_1$.

Let $\mu = \delta_1$, the Dirac point mass at 1. We computed the Kato-Freidrichs operator H_w in Example 5.15, where the weights $w = \{w_j\}_{j \in \mathbb{N}_0} \subset \mathbb{R}^+$ are chosen such that $\sum_j \frac{1}{w_j} < \infty$. We found that H_w is a (bounded) rank-one operator on $\ell^2(w)$ defined by

$$(H_w c)_j = \frac{1}{w_j} \sum_k c_k.$$

Recall that we also know from Example 6.6 that the dimension of $L^2(\mu)$ must be exactly 1.

If we choose weights $\{w_i\} \subset \mathbb{R}^+$ such that

$$\sum_{i \in \mathbb{N}_0} \frac{1}{w_i} = 1,$$

then $H_w : \ell^2(w) \rightarrow \ell^2(w)$ is in fact a rank-one projection—that is,

$$(7.39) \quad H_w^2 = H_w = (H_w)^*.$$

In this case, we know from the spectral theory of projections that the spectrum of H_w is exactly $\{0, 1\}$, and the operator norm of H_w is 1.

If we generalize to the case where $\mu = \delta_b$ for some real value b , then the moment matrix is

$$M_{j,k} = b^{j+k},$$

and the Kato-Friedrichs operator H_w for weights w is a rank-one operator with range the span of

$$\xi_b = \begin{bmatrix} \frac{1}{w_0} & \frac{b}{w_1} & \frac{b^2}{w_2} & \cdots \end{bmatrix}.$$

$\diamond$

EXAMPLE 7.20. The Kato-Friedrichs operator associated with $\mu = \frac{1}{2}(\delta_0 + \delta_1)$

When $\mu = \frac{1}{2}(\delta_0 + \delta_1)$, the moment matrix M is given by

$$(7.40) \quad M = \frac{1}{2} \begin{bmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 0 & \cdots \\ 0 & 0 & 0 & 0 & \cdots \\ \vdots & \vdots & & & \ddots \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 & \cdots \\ 1 & 1 & 1 & 1 & \cdots \\ 1 & 1 & 1 & 1 & \cdots \\ \vdots & \vdots & & & \ddots \end{bmatrix}.$$

We wish to solve the eigenvalue equations (7.24), which in this case become

$$(7.41) \quad \begin{aligned} c_0 + \frac{1}{2}(c_1 + c_2 + \cdots) &= \lambda w_0 c_0, & k = 0 \\ \frac{1}{2}(c_0 + c_1 + c_2 + \cdots) &= \lambda w_k c_k, & k \geq 1. \end{aligned}$$

which can be transformed into

$$(7.42) \quad \frac{1}{2}c_0 = \lambda(w_0 c_0 - w_k c_k), \quad k \geq 1.$$

by subtracting the later equations in (7.41) from the first.

To solve for c in Equation (7.42), we consider the cases where $\lambda = 0$ and where $\lambda \neq 0$.

Case 1: If $\lambda = 0$, c_0 must also be 0. In addition, referring back to the equations in the system (7.41), we easily see that Equation (7.24) is true whenever

$$(7.43) \quad \sum_{k=1}^{\infty} c_k = 0.$$

(For example, $c = (0, 1, -1, 0, 0, \dots)$.) There is no restriction on the weights $w = \{w_k\}_{k \in \mathbb{N}_0}$ coming from the equations here. So for any choice of weights, the eigenspace for $\lambda = 0$ is infinite-dimensional.

Case 2: If $\lambda \neq 0$, the k^{th} equation for $k > 0$ in the system of equations (7.42) can be solved for c_k in terms of λ , c_0 , and the weights $w = \{w_k\}_{k \in \mathbb{N}}$:

$$(7.44) \quad c_k = \frac{c_0}{w_k} \left(w_0 - \frac{1}{2\lambda} \right), \quad k \geq 1.$$

We see that $c_0 = 0$ implies that $c_k = 0$ for all k , so we require $c_0 \neq 0$. Substituting (7.44) into the first equation in (7.41), and cancelling c_0 from both sides, we obtain a condition on λ and the weights w :

$$(7.45) \quad 1 + \frac{1}{2} \left(\sum_{k=1}^{\infty} \frac{1}{w_k} \right) \left(w_0 - \frac{1}{2\lambda} \right) = \lambda w_0.$$

Define $T = \sum_{k=1}^{\infty} \frac{1}{w_k}$, where this sum being finite imposes a restriction on our choice of weights w . Then the equation becomes

$$1 + \frac{1}{2} T \left(w_0 - \frac{1}{2\lambda} \right) = \lambda w_0.$$

Observe that the two conditions

$$1 = \lambda w_0 \quad \text{and} \quad 2\lambda w_0 = 1$$

will lead to inconsistent systems, so we must rule these conditions out. Otherwise, there are many solutions to Equations (7.24).

We now demonstrate that there will be two distinct nonzero eigenvalues for the Kato-Friedrichs operator H_w , where the weights w are chosen so that T is finite. Without loss of generality, take $w_0 = c_0 = 1$. Our eigenvalues $\lambda \neq 0$ must satisfy Equation (7.45), which is now

$$(7.46) \quad 1 + \frac{1}{2} T \left(1 - \frac{1}{2\lambda} \right) = \lambda.$$

This yields the quadratic equation in λ :

$$\lambda^2 - \left(1 + \frac{T}{2} \right) \lambda + \frac{T}{4} = 0.$$

This equation has distinct positive roots λ_+, λ_- for $T > 0$, and they are given by

$$(7.47) \quad \lambda_{\pm} = \frac{1 + \frac{T}{2} \pm \sqrt{1 + \left(\frac{T}{2} \right)^2}}{2};$$

and we also see

$$(7.48) \quad \lambda_+ \lambda_- = \frac{T}{4} \quad \text{and} \quad \lambda_+ + \lambda_- = 1 + \frac{T}{2}.$$

Since the vector c satisfying $Hc = \lambda c$ (where $\lambda \neq 0$) is uniquely determined from λ, c_0 , and w , given a choice of weights and taking $c_0 = 1$, each eigenspace for nonzero λ is one-dimensional. Denote by c_+ and c_- these eigenvectors in the eigenspaces for λ_+ and λ_- respectively.

Going back to (7.44) we have the simplification for the eigenvectors:

$$(7.49) \quad c_k = \frac{1}{w_k} \left(1 - \frac{1}{2\lambda} \right) = \frac{1}{w_k} \left(\frac{\lambda - \frac{1}{2}}{\lambda} \right), \quad k \geq 1.$$

With respect to our eigenvalues $\lambda_{\pm}$, this gives

$$c_{\pm} = \left(1, \frac{\lambda_{\pm} - \frac{1}{2}}{w_1 \lambda_{\pm}}, \frac{\lambda_{\pm} - \frac{1}{2}}{w_2 \lambda_{\pm}}, \dots \right);$$

then

$$\|c_{\pm}\|_{\ell^2(w)}^2 \stackrel{(7.49)}{=} 1 + \left(\frac{\lambda_{\pm} - \frac{1}{2}}{\lambda_{\pm}} \right)^2 T.$$

The reader can verify the orthogonality of the eigenvectors.

In order to do more explicit computations, let us choose the weights $w = \{w_k\}_{k \in \mathbb{N}_0}$ such that $T = 2$. This yields

$$\lambda_{\pm} = 1 \pm \frac{1}{\sqrt{2}},$$

which are both positive.

Substituting back into (7.49) we get

$$(7.50) \quad c_{\pm} = \left(1, \pm \frac{1}{\sqrt{2}w_1}, \pm \frac{1}{\sqrt{2}w_2}, \dots \right),$$

with

$$\|c_{\pm}\|_{\ell^2(w)}^2 = 2.$$

Normalize to obtain unit vectors

$$(7.51) \quad \xi_{\pm} := \frac{1}{\sqrt{2}} c_{\pm}$$

which yield the rank-one projections

$$E_{\pm} := |\xi_{\pm}\rangle\langle\xi_{\pm}|$$

onto the eigenspaces. In other words, the spectral resolution of the self-adjoint operator H can be written

$$(7.52) \quad H = \left(1 + \frac{1}{\sqrt{2}} \right) E_+ + \left(1 - \frac{1}{\sqrt{2}} \right) E_-.$$

We next make two observations regarding the connections between $H^{1/2}$, the square root of the Kato-Friedrichs operator, and the Hilbert space $L^2(\mu)$. First, we recall from Equation (5.13) and Lemma 5.8 the isometry

$$(7.53) \quad \int |f_c|^2 d\mu = \|H^{1/2}c\|_{\ell^2(w)}^2;$$

that is, the identification $f_c \leftrightarrow H^{1/2}c$ is isometric. By choosing specific weights, we can verify this isometry using our Kato-Friedrichs operator for $\mu = \frac{1}{2}(\delta_0 + \delta_1)$.

Let $w = \{w_k\}$ be given by the sequence with $w_0 = 1$ and

$$w_k = 2^{k-1}, \quad k \geq 1,$$

which gives $T = 2$. Denote $F(\xi_{\pm})$ by the shorthand $f_{\pm}$. Using (7.50) and (7.51), we get

$$(7.54) \quad f_{\pm}(x) = \sum_{k=0}^{\infty} (\xi_{\pm})_k x^k = \frac{1}{\sqrt{2}} \pm \frac{x}{2-x},$$

and, because $\mu = \frac{1}{2}(\delta_0 + \delta_1)$, we have

$$\int |f_{\pm}(x)|^2 d\mu(x) = 1 \pm \frac{1}{\sqrt{2}} = \frac{\sqrt{2} \pm 1}{\sqrt{2}}.$$

Using Equation (7.52), we have

$$H^{1/2} = \left(1 + \frac{1}{\sqrt{2}}\right)^{1/2} E_+ + \left(1 - \frac{1}{\sqrt{2}}\right)^{1/2} E_-.$$

Moreover,

$$(7.55) \quad H^{1/2} \xi_{\pm} = \left(1 \pm \frac{1}{\sqrt{2}}\right)^{1/2} \xi_{\pm}$$

and hence

$$\|H^{1/2} \xi_{\pm}\|_{\ell^2(w)}^2 = 1 \pm \frac{1}{\sqrt{2}}$$

which verifies the isometry in (7.53).

Define the map $W : L^2(\mu) \rightarrow \ell^2(w)$ given by

$$(7.56) \quad W(f_c) = H^{1/2} c \text{ for } c \in \mathcal{D} \subset \ell^2(w).$$

Recall from Example 6.6 that the dimension of $L^2(\mu)$ is exactly 2. Our second observation is that the map W is an isometry into a two-dimensional subspace of $\ell^2(w)$.

We will check this fact directly. Recall $E_{\pm} = |\xi_{\pm}\rangle\langle\xi_{\pm}|$, with $\xi_{\pm}$ as in (7.51). We only need to check that W takes an ONB in $L^2(\mu)$ to an orthonormal family in $\ell^2(w)$. First, observe that the orthogonal polynomials in $L^2(\mu)$ are $p_0(x) \equiv 1$, $p_1(x) = 2x - 1$, and $p_k(x) \equiv 0$ for $k \geq 2$. Moreover,

$$\begin{aligned} Wp_0 &= Wf_{(1,0,0,\dots)} \\ &\stackrel{(7.56)}{=} \left(1 + \frac{1}{\sqrt{2}}\right)^{1/2} E_+(1, 0, 0, \dots) + \left(1 - \frac{1}{\sqrt{2}}\right)^{1/2} E_-(1, 0, 0, \dots) \\ &\stackrel{(7.51)}{=} \left(1 + \frac{1}{\sqrt{2}}\right)^{1/2} \frac{1}{\sqrt{2}} \xi_+ + \left(1 - \frac{1}{\sqrt{2}}\right)^{1/2} \frac{1}{\sqrt{2}} \xi_-, \end{aligned}$$

and

$$\begin{aligned} Wp_1 &= Wf_{(-1,2,0,0,\dots)} \\ &= \left(1 + \frac{1}{\sqrt{2}}\right)^{1/2} \left(1 - \frac{1}{\sqrt{2}}\right) \xi_+ + \left(1 - \frac{1}{\sqrt{2}}\right)^{1/2} \left(-1 - \frac{1}{\sqrt{2}}\right) \xi_- \\ &= \left(1 - \frac{1}{\sqrt{2}}\right)^{1/2} \frac{1}{\sqrt{2}} \xi_+ - \left(1 + \frac{1}{\sqrt{2}}\right)^{1/2} \frac{1}{\sqrt{2}} \xi_-. \end{aligned}$$

Since Wp_0 and Wp_1 are orthonormal in $\ell^2(w)$, we have that W is an isometry. $\diamond$

EXAMPLE 7.21. The Kato-Friedrichs operator associated with $\mu = \frac{1}{2}(\delta_0 + \delta_b)$, $0 < b < 1$

This example is a more general case where μ is a finite convex combination of Dirac masses, hence the associated infinite Hankel matrix will be of finite rank. To simplify matters we pick the atoms in μ in such a way that no weights will be needed. When we work with an unweighted ℓ^2 space, $w_k = 1$ for all $k \in \mathbb{N}_0$, and $\ell^2 = \ell^2(w)$. The moment matrix $M = M^{(\mu)}$ for the convex combination μ is equal to the Kato-Friedrichs operator, as in the case of the Hilbert matrix. This last example also has the advantage of illustrating an ONB in $L^2(\mu)$ consisting of rational functions; not polynomials. The choice of the rational functions is dictated by our Kato-Friedrichs operator $H = M$.

As in the previous Example 7.20, H has a spectrum consisting of 0 and two points on the positive real line, and $L^2(\mu)$ is two-dimensional. The point 0 has infinite multiplicity, and the two positive eigenvalues are simple, i.e., have multiplicity one.

For $\mu = \frac{1}{2}(\delta_0 + \delta_b)$, $0 < b < 1$, the moment matrix M is

$$(7.57) \quad M = H = \begin{bmatrix} 1 & \frac{1}{2}b & \frac{1}{2}b^2 & \cdots \\ \frac{1}{2}b & \frac{1}{2}b^2 & \frac{1}{2}b^3 & \cdots \\ \frac{1}{2}b^2 & \frac{1}{2}b^3 & \frac{1}{2}b^4 & \cdots \\ \vdots & \vdots & & \ddots \end{bmatrix}.$$

The orthogonal polynomials in $L^2(\mu)$ are

$$(7.58) \quad p_0(x) \equiv 1, \quad p_1(x) = 1 - \frac{2x}{b}, \quad p_k(x) \equiv 0, k \geq 2.$$

However, we also have an orthogonal basis in $L^2(\mu)$ consisting of the two rational functions $f_{\pm}(x)$, where

$$(7.59) \quad f_{\pm}(x) := \alpha_{\pm} + \frac{bx}{1-bx}.$$

These correspond to the eigenvectors for H . We see that if

$$\xi_{\pm} = [\alpha_{\pm} \quad b \quad b^2 \quad b^3 \quad \cdots]^{tr},$$

then $H\xi_{\pm} = \lambda\xi_{\pm}$, where the parameters $\alpha_{\pm}$ are given by

$$\alpha_{\pm} = 1 - \frac{p}{2} \pm \sqrt{\left(\frac{p}{2}\right)^2 + 1}$$

and the two positive eigenvalues of H are

$$\lambda_{\pm} = \frac{1}{2} \left(1 + \frac{p}{2} \pm \sqrt{\left(\frac{p}{2}\right)^2 + 1} \right),$$

where p is the constant

$$p = \frac{b^2}{1-b^2}.$$

The functions above are the images of $\xi_{\pm}$ under the operator F :

$$f_{\pm}(x) = (F\xi_{\pm})(x) = \alpha_{\pm} + \frac{bx}{1-bx}.$$

The eigenvectors $\xi_{\pm}$ are orthogonal, and it follows that $f_{\pm}$ are orthogonal in $L^2(\mu)$:

$$\begin{aligned} \langle f_+ | f_- \rangle_{L^2(\mu)} &= \langle F\xi_+ | F\xi_- \rangle_{L^2(\mu)} \\ &= \langle \xi_+ | F^* F \xi_- \rangle_{\ell^2} \\ &= \langle \xi_+ | H \xi_- \rangle_{\ell^2} \\ &= \lambda_- \langle \xi_+ | \xi_- \rangle_{\ell^2} = 0 \end{aligned}$$

Direct computation of $\langle f_+ | f_- \rangle_{L^2(\mu)}$ also verifies that these functions are orthogonal.

This example also demonstrates the correspondence from Theorem 7.11 between atoms of the measure μ and atoms in the projection-valued measure E associated to H . The measure μ has two atoms at 0 and b , while the projection valued measure E for H has atoms at the two nonzero eigenvalues $\lambda_{\pm}$. $\diamond$

CHAPTER 8

The moment problem revisited

Given a positive definite Hankel matrix M with real entries, we showed in Section 2.1 that there exists a measure μ (not unique in general) such that M is the moment matrix of μ , i.e., $M = M^{(\mu)}$. In this chapter, we will describe a setting in which the solution measure is not unique. Differing measures solving the same moment problem will arise from nontrivial self-adjoint extensions of a symmetric shift operator S . In fact, we will show that if μ is not unique, the self-adjoint extensions of S yield a one-parameter family of measures satisfying the moment problem for M . This will yield a necessary condition for non-uniqueness of measure for a given Hankel matrix M . For more details on the theory of self-adjoint extensions of unbounded symmetric operators and moments, see [Con90, RS75, Rud91].

Given a Hankel matrix M with real entries, let Q_M be the quadratic form from Equation (5.1) and let $\mathcal{H}_Q$ be the Hilbert space completion of Q_M . Then the inner product on $\mathcal{H}_Q$ is defined on the finite sequences $\mathcal{D}$ by

$$\langle c|d \rangle_{\mathcal{H}_Q} = \sum_i \sum_j \bar{c}_i M_{i,j} d_j.$$

We define the shift operator S by

$$(8.1) \quad Sc = S(c_0, c_1, \dots) = (0, c_0, c_1, \dots).$$

The domain of S contains $\mathcal{D}$ which is dense in $\mathcal{H}_Q$, so S is densely defined.

In Section 8.1 we observe some of the interplay between the matrix M , the shift operator S , and the isometry F . Then, in Sections 8.2 and 8.3 we describe how the self-adjoint extensions of S yield solutions to the moment problem $M = M^{(\mu)}$. We also use the shift operator in Section 8.4 in order to find a Jacobi matrix corresponding to a given Hankel matrix M .

8.1. The shift operator and three incarnations of symmetry

Suppose M is a positive definite Hankel matrix and μ is a measure such that $M = M^{(\mu)}$. Recall that we have defined an isometry $F : \mathcal{H}_Q \rightarrow L^2(\mu)$ by

$$Fc(x) = f_c(x) = \sum_{n \in \mathbb{N}_0} c_n x^n \text{ for all } c \in \mathcal{D}.$$

The shift operator S plays a major role in this chapter, and in this section we study how F and S behave with respect to each other. The operators F and S reveal three different incarnations of symmetry in the Hankel matrix M . In turn, these incarnations will be used in subsequent sections to prove results about non-uniqueness of measures which solve the moment problem, particularly in Theorem 8.12.

Incarnation 1: A symmetric operator. We see here the connection between the Hankel property of M and the shift operator in the Hilbert space $\mathcal{H}_Q$.

LEMMA 8.1. *The shift operator S is symmetric in $\mathcal{D} \subset \mathcal{H}_Q$ if and only if the matrix M is a Hankel matrix.*

PROOF. ($\Rightarrow$): Setting $b = e_i$, $c = e_j$, and $\langle Sb|c \rangle_{\mathcal{H}_Q} = \langle b|Sc \rangle_{\mathcal{H}_Q}$, we see that $M_{i-1,j} = M_{i,j-1}$.

($\Leftarrow$): Let $b, c \in \mathcal{D}$. Then

$$\begin{aligned} \langle Sb|c \rangle_{\mathcal{H}_Q} &= \sum_i \sum_j \overline{b_{i-1}} M_{i+j} c_j = \sum_i \sum_j \overline{b_i} M_{i+1+j} c_j \\ (8.2) \quad &= \sum_i \sum_j \overline{b_i} M_{i+j} c_{j-1} = \langle b|Sc \rangle_{\mathcal{H}_Q}. \end{aligned}$$

□

Incarnation 2: Multiplication by x . We notice that if p and q are polynomials in $L^2(\mu)$, then

$$\int \overline{xp(x)} q(x) d\mu(x) = \int \overline{p(x)} xq(x) d\mu(x).$$

We can therefore state the interaction between the isometry F and the shift S .

LEMMA 8.2. *Define $\tilde{S} := F S F^*$. Then $\tilde{S}$ is a symmetric operator on $\mathcal{P} \subset L^2(\mu)$, and in addition, $\tilde{S}$ is a restriction (to its domain) of the multiplication operator M_x :*

$$[M_x f](x) = x f(x).$$

PROOF. Let M_x be the operator which takes $f(x)$ to $x f(x)$. Using the definitions of F and S , we see that

$$(8.3) \quad M_x(Fc) = FSc \text{ for all } c \in \mathcal{D},$$

or stated equivalently,

$$x f_c(x) = f_{Sc}(x) \text{ for all } c \in \mathcal{D}.$$

Since F is an isometry, we know that F^*F is the identity on $\mathcal{H}_Q$. As a result, applying F^* on the left to both sides of Equation (8.3) yields

$$S = F^* M_x F \text{ on } \mathcal{D}.$$

Also, $P = F F^*$ is the projection of $L^2(\mu)$ onto the closure of the polynomials $\mathcal{P} \subset L^2(\mu)$. Applying F^* on the right to both sides of Equation (8.3) yields

$$M_x P = F S F^* = \tilde{S} \text{ on } \mathcal{P},$$

which is the desired conclusion. □

Incarnation 3: Jacobi matrices and orthogonal polynomials. Looking ahead in Section 8.4, we see that given the space $L^2(\mu)$, there is a Jacobi matrix J which encodes the multiplication operator M_x in terms of a recursion relation for orthogonal polynomials $\{p_k\}_{k \in \mathbb{N}_0}$ in $L^2(\mu)$. Specifically,

$$J \begin{bmatrix} p_0 \\ p_1 \\ \vdots \end{bmatrix} = \begin{bmatrix} M_x p_0 \\ M_x p_1 \\ \vdots \end{bmatrix}.$$

Lemma 8.20 then gives an intertwining relationship between the shift S and this Jacobi matrix J .

8.2. Self-adjoint extensions of a shift operator

Recall from Definition 1.4 that a densely defined operator S on a Hilbert space $\mathcal{H}$ is *symmetric* on $\text{dom}(S) \subseteq \mathcal{H}$ if $\langle Sh|k \rangle = \langle h|Sk \rangle$ for all $h, k \in \text{dom}(S)$. Also recall that we can define the *adjoint* S^* of S , as in Definition 1.3, and we call S a *self-adjoint* operator if $S = S^*$, in particular, if $\text{dom}(S) = \text{dom}(S^*)$.

DEFINITION 8.3. Suppose S is a densely defined operator on $\mathcal{D}$ in a Hilbert space $\mathcal{H}$. A *self-adjoint* extension T of S satisfies the following properties:

- (1) T is self-adjoint with $\text{dom}(T) = \text{dom}(T^*)$
- (2) $Tc = Sc$ for all $c \in \mathcal{D}$.

It may be the case that no self-adjoint extensions of S exist.

If S is essentially self-adjoint, we call the closure of S a trivial self-adjoint extension of S . Finally, we say that S is *maximally symmetric* if S has no proper self-adjoint extensions.

Let S be the closure of the shift operator from Equation (8.1). By Lemma 8.1, S is symmetric in $\mathcal{D}$ if and only if the matrix M is a Hankel matrix. The deficiency indices of S will allow us to describe its self-adjoint extensions, if any exist.

DEFINITION 8.4. Let S be a closed symmetric operator on a Hilbert space $\mathcal{H}$ and let $\alpha \in \mathbb{C}$ with $\text{Im}(\alpha) \neq 0$. We define the *deficiency subspace* of S at α by

$$(8.4) \quad \mathcal{L}(\alpha) = \text{null}(S^* - \alpha) = \{\xi \in \text{dom}(S^*) : S^*\xi = \alpha\xi\}.$$

Due to a beautiful argument of von Neumann, the dimension of the deficiency subspace will be the same for any α with $\text{Im}(\alpha) > 0$, and the dimension will be the same for any α with $\text{Im}(\alpha) < 0$. It is therefore sufficient to consider $\alpha = \pm i$. We will denote the respective deficiency subspaces for the shift operator $\mathcal{L}_+$ and $\mathcal{L}_-$.

DEFINITION 8.5. The dimensions of $\mathcal{L}_+$ and $\mathcal{L}_-$ for a closed symmetric operator S are called the *deficiency indices* of S and are denoted by the ordered pair $(\dim(\mathcal{L}_+), \dim(\mathcal{L}_-))$. Note that the indices can take on any value in $\mathbb{N}_0$ or ∞ .

DEFINITION 8.6. Let S and T be linear operators with dense domains in a Hilbert space, and let $\text{Gr}(S), \text{Gr}(T)$ be their corresponding graphs. We say that $S \subseteq T$ if $\text{Gr}(S) \subseteq \text{Gr}(T)$.

Note that an operator S is self-adjoint if and only if $S \subseteq S^*$. If T is a self-adjoint extension of S , then

$$S \subseteq T \subseteq T^* \subseteq S^*.$$

Given a bounded operator J , we write $JS \subseteq SJ$ if J maps the domain of S into itself and $JSv = SJv$ for all $v \in \text{dom}(S)$.

Another theorem of von Neumann gives a condition under which the deficiency indices of a symmetric operator are equal.

THEOREM 8.7. (von Neumann, as stated in [Con90, Prop. 7.2, p. 343]) *Given an operator S on a Hilbert space $\mathcal{H}$, if there exists a function $J : \mathcal{H} \rightarrow \mathcal{H}$ satisfying the following properties:*

- (1) J^2 is the identity on $\mathcal{H}$,

(2) J is conjugate linear—that is, for all $\alpha \in \mathbb{C}$, $J(\alpha h) = \overline{\alpha}J(h)$,

(3) $\|Jh\| = \|h\|$ for all $h \in \mathcal{H}$,

(4) $J\text{dom}(S) \subseteq \text{dom}(S)$ and $JS \subseteq SJ$,

then S has equal deficiency indices, i.e. $\dim(\mathcal{L}_+) = \dim(\mathcal{L}_-)$.

The main idea in the proof of this theorem is that the operator J restricts as an isometry between the deficiency subspaces $\mathcal{L}_+$ and $\mathcal{L}_-$, thus showing they have equal dimension.

We apply this theorem to our shift operator S on the space $\mathcal{H}_Q$. Let J be the conjugation operator:

$$(8.5) \quad J : \mathcal{H}_Q \rightarrow \mathcal{H}_Q \text{ with } J(c_0, c_1, c_2, \dots) := (\overline{c_0}, \overline{c_1}, \overline{c_2}, \dots),$$

If $\xi \in \mathcal{L}_+$, then $J\xi \in \mathcal{L}_-$. It is readily verified that J satisfies the properties in Theorem 8.7. In particular we see that on $\mathcal{D} = \text{dom}(S)$, J commutes with S .

We now calculate the deficiency indices for S , the closure of the shift operator.

LEMMA 8.8. *The closed shift operator S (8.1) defined on $\mathcal{D} \subset \mathcal{H}_M$ is either self-adjoint or has deficiency indices $(1, 1)$ —i.e.*

$$\dim\{\xi \in \text{dom}(S^*) : S^*\xi = i\xi\} = 1.$$

In fact, if $\xi \in \mathcal{L}_+$ with $\xi \neq 0$, then $H\xi$ is a multiple of the vector $(1, i, i^2, i^3, \dots)$, where H is the self-adjoint Kato operator for the quadratic form Q_M on $\mathcal{H}_Q$.

PROOF. If S is self-adjoint, its deficiency indices are $(0, 0)$. Suppose there exists $\xi \neq 0$ such that $\xi \in \mathcal{L}_+$. We will compute $(H\xi)_i$ via the inner product $\langle \cdot | \cdot \rangle_{\mathcal{H}_Q}$ to show that

$$(8.6) \quad H\xi \in \mathbb{C}(1, i, i^2, i^3, \dots).$$

Note e_i belongs to $\mathcal{D} = \text{dom}(S)$ for every $i \in \mathbb{N}_0$. We now compare $(H\xi)_i = \langle e_i | \xi \rangle_{\mathcal{H}_Q}$ and $(H\xi)_{i+1} = \langle e_{i+1} | \xi \rangle_{\mathcal{H}_Q}$:

$$(8.7) \quad \langle e_{i+1} | \xi \rangle_{\mathcal{H}_Q} = \langle Se_i | \xi \rangle_{\mathcal{H}_Q} = \langle e_i | S^*\xi \rangle_{\mathcal{H}_Q} = i\langle e_i | \xi \rangle_{\mathcal{H}_Q},$$

which implies by induction that $H\xi$ is a multiple of the vector (8.6).

Since H has trivial kernel in $\mathcal{H}_Q$, we can conclude that $\dim \mathcal{L}_+ = 1$ and by Theorem 8.7, $\dim \mathcal{L}_- = 1$ as well. $\square$

We next use a theorem, again quoted almost verbatim from [Con90], which states that the self-adjoint extensions of S are determined by the partial isometries from $\mathcal{L}_+$ to $\mathcal{L}_-$.

THEOREM 8.9. [Con90, Theorem 2.17, p. 314] *Let S be a closed symmetric operator. If W is a partial isometry with initial space in $\mathcal{L}_+$ and final space in $\mathcal{L}_-$, then there is a closed symmetric extension S_W of S on the domain*

$$\{f + g + Wg : f \in \text{dom}(S), g \in \text{initial}(W)\}$$

given by

$$S_W(f + g + Wg) = Sf + ig - iWg.$$

Conversely, if T is any closed symmetric extension of S , then there is a unique partial isometry W such that $T = S_W$ as defined above.

In the case of our shift operator S , we see that the only nontrivial partial isometries from $\mathcal{L}_+$ to $\mathcal{L}_-$ are isometries between the one-dimensional spaces. These isometries are given by multiplication by $z \in \mathbb{C}$ where $|z| = 1$.

THEOREM 8.10. *Given a Hankel matrix M and the associated Hilbert space $\mathcal{H}_Q$, let S be the closure of the symmetric shift operator (8.1). If S is not self-adjoint, then it has self-adjoint extensions T_z which have domain $\mathcal{D} + \mathcal{L}_+ + \mathcal{L}_-$ and are exactly given by*

$$T_z(c + \xi + J\xi) = Sc + i\xi - izJ\xi \quad \text{for } z \in \mathbb{C}.$$

PROOF. By Lemma 8.8, if S is not self-adjoint, it has deficiency indices $(1, 1)$. If $\xi \in \mathcal{L}_+$, then $J\xi \in \mathcal{L}_-$. Given $z \in \mathbb{C}$ with $|z| = 1$, we can define an isometry $T_z : \mathcal{L}_+ \rightarrow \mathcal{L}_-$ by $T\xi = zJ\xi$. In fact, every such isometry is of this form since the spaces are one-dimensional. The result follows from Theorem 8.9. $\square$

8.3. Self-adjoint extensions and the moment problem

Next, we describe how the self-adjoint extensions T_z to S described in Theorem 8.10 yield solutions μ_z to the moment problem $M = M^{(\mu)}$. (For background, see [Con90].)

As we discussed in Section 7.2, every self-adjoint operator T can be written in terms of a projection-valued measure E such that

$$(8.8) \quad T = \int_{\mathbb{R}} \lambda E(d\lambda).$$

PROPOSITION 8.11. [Con90, Prop. 7.2, p. 343] *Given an infinite Hankel matrix M , suppose T is a self-adjoint nontrivial extension of the shift operator S on the Hilbert space $\mathcal{H}_Q$ with corresponding projection-valued measure E (8.8). Given E , we can define a real measure as we did in Equation (7.8), by*

$$\mu(\cdot) := \langle e_0 | E(\cdot) e_0 \rangle_{\mathcal{H}_Q},$$

where e_0 is the first standard basis vector $(1, 0, 0, \dots)$. Then the real-valued measure μ is a solution to the moment problem for M —that is,

$$\int x^i d\mu(x) = M_i \text{ for all } i \in \mathbb{N}_0.$$

PROOF. From the multiplicative property of integrals against projection-valued measures,

$$\int_{\mathbb{R}} \lambda^i E(d\lambda) = T^i$$

for each $i \in \mathbb{N}_0$. This gives the corresponding result for μ :

$$\int_{\mathbb{R}} x^i d\mu(x) = \langle e_0 | T^i e_0 \rangle.$$

Since $e_0 \in \mathcal{D}$ and T maps $\mathcal{D}$ to $\mathcal{D}$, we have the following calculation:

$$(8.9) \quad \begin{aligned} \int x^i d\mu(x) &= \langle e_0 | T^i e_0 \rangle_{\mathcal{H}_Q} = \langle e_0 | S^i e_0 \rangle_{\mathcal{H}_Q} = \langle e_0 | e_i \rangle_{\mathcal{H}_Q} \\ &= M_{0,i} = M_i. \end{aligned}$$

$\square$

We can now associate to each self-adjoint extension to the shift operator S a measure which satisfies the moment problem for the matrix M , in the case where the shift operator is not essentially self-adjoint.

THEOREM 8.12. *Given an infinite Hankel matrix M with $M_{0,0} = 1$, let S be the closure of the shift operator on $\mathcal{H}_Q$. Then the following statements are equivalent.*

- (1) *S is not self-adjoint.*
- (2) *The set of distinct solutions to the moment problem $M = M^{(\mu)}$ is a one-parameter family of probability measures $\{\mu_z : z \in \mathbb{C}, |z| = 1\}$.*
- (3) *There exist two nonequivalent probability measures μ_1 and μ_2 which are both solutions to the moment problem, i.e. $M = M^{(\mu_1)} = M^{(\mu_2)}$.*
- (4) *Given any measure μ solving the moment problem $M = M^{(\mu)}$, the polynomials are not dense in the space $L^2(\mu)$.*

PROOF. (1) $\Rightarrow$ (2): If S is not self-adjoint, then by Theorem 8.10 there is a one-parameter family $\{T_z : z \in \mathbb{C}, |z| = 1\}$ of distinct (and not unitarily equivalent) self-adjoint extensions to S . For each T_z , we can define a measure μ_z as in Proposition 8.11 which satisfies the moment problem $M = M^{(\mu)}$. It remains to be shown that these measures are distinct.

The isometry $F : \mathcal{H}_Q \rightarrow L^2(\mu_z)$ which maps $T_z^k e_0$ to x^k extends to map $\psi(T_z)(e_0)$ to the function ψ . Thus, because the span of the functions $\{\frac{1}{x-\alpha} : \alpha \in \mathbb{C}, \text{Im}(\alpha) \neq 0\}$ is dense in each space $L^2(\mu_z)$, the measure μ_z is uniquely determined by these functions. The extensions T_z are cyclic operators on $\mathcal{H}_Q$, which means (see [Con90]) that the set of vectors $\{(T_z - \alpha I)^{-1} e_0 : \alpha \in \mathbb{C}, \text{Im}(\alpha) \neq 0\}$ is dense in $\mathcal{H}_Q$. It follows that μ_z is determined uniquely by the isomorphism F mapping the vector $(T_z - \alpha)^{-1} e_0 \in \mathcal{H}_Q$ to the function $\frac{1}{x-\alpha} \in L^2(\mu)$. Since the operators T_z are distinct, this proves the measures $\{\mu_z : z \in \mathbb{D}, |z| = 1\}$ are distinct.

(2) $\Rightarrow$ (3): Follows because each μ_z is distinct.

(3) $\Rightarrow$ (4): We prove the contrapositive. Suppose the space of polynomials $\mathcal{P}$ is dense in $L^2(\mu)$ where $M = M^{(\mu)}$. Recall the map $F : \mathcal{D} \rightarrow \mathcal{P}$ given by $Fc = \sum_i c_i x^i$ is an isometry on $\mathcal{D}$ which extends to an isometry on $\mathcal{H}_Q$. Assume $\xi \in \mathcal{L}_+$, so $S^* \xi = i\xi$ and let $c \in \mathcal{D}$. Then

$$\begin{aligned} \int_{\mathbb{R}} \overline{xFc(x)} F\xi(x) d\mu(x) &= \langle FSc | F\xi \rangle_{L^2(\mu)} \\ &= \langle Sc | \xi \rangle_{\mathcal{H}_Q} = \langle c | S^* \xi \rangle_{\mathcal{H}_Q} = i \langle c | \xi \rangle_{\mathcal{H}_Q} \\ &= i \langle Fc | F\xi \rangle_{L^2(\mu)} = i \int_{\mathbb{R}} \overline{Fc(x)} F\xi(x) d\mu(x). \end{aligned}$$

Because F gives a one-to-one correspondence between $\mathcal{D}$ and $\mathcal{P}$, we can say for any polynomial $p \in \mathcal{P}$ and $\xi \in \mathcal{L}_+$,

$$(8.10) \quad \int_{\mathbb{R}} (x-i) \overline{p(x)} F\xi(x) d\mu(x) = 0.$$

The function $\frac{1}{x-i} \in L^\infty(\mu)$, hence $\frac{F\xi}{x-i} \in L^2(\mu)$. Let $\{p_n\} \subset \mathcal{P}$ converge in $L^2(\mu)$ to $\frac{F\xi}{x-i} \in L^2(\mu)$. Substituting into Equation (8.10) then gives

$$\int_{\mathbb{R}} |F\xi(x)|^2 d\mu(x) = 0,$$

hence $\xi = 0$. Therefore, S has deficiency indices $(0,0)$.

(4) $\Rightarrow$ (1): Assume the polynomials are not dense in $L^2(\mu)$. As before, $M = M^{(\mu)}$. Let $\psi \in L^2(\mu)$ be a nonzero bounded vector orthogonal to the subspace $\mathcal{P}$ spanned by the polynomials. Recall that $L^2(\mu) \ominus \mathcal{P} \neq 0$ if and only if $\mathcal{P}$ is not dense in

$L^2(\mu)$. Given $\alpha \in \mathbb{C}$ with $\text{Im}(\alpha) \neq 0$, define

$$\xi_\alpha = \frac{\psi(x)}{x - \alpha}.$$

Then, because each function $\frac{1}{x - \alpha}$ is bounded, we have $\xi_\alpha \in L^2(\mu)$. Note that the function $[M_x \xi_\alpha](x) = x \xi_\alpha(x)$ is also an $L^2(\mu)$ function because $x \xi_\alpha(x) = \psi(x) + \alpha \xi_\alpha(x)$.

The isometry F between $\mathcal{H}_Q$ and $L^2(\mu)$ resulting from Equation (5.2) ensures that FF^* is the projection onto the range of F , which is the closed space spanned by the polynomials in $L^2(\mu)$. Therefore, we know $F^*\psi = 0$. Given the function ξ_α , let $\phi_\alpha \in \mathcal{H}_Q$ be defined by $\phi_\alpha = F^*\xi_\alpha$. We first must show that for at least one choice of α , $\phi_\alpha \neq 0$.

Suppose that for all $\alpha \in \mathbb{C} \setminus \mathbb{R}$, ξ_α is orthogonal to the polynomials. Define Ψ to be the linear span of the functions $\{\frac{1}{x - \alpha} : \alpha \in \mathbb{C}, \text{Im}(\alpha) \neq 0\}$. It is well-known that Ψ is dense in $L^2(\mu)$ when μ is a finite real measure, so let $\{\psi_k\}_{k \in \mathbb{N}_0} \subset \Psi$ be a sequence of functions which converges to $\bar{\psi}$ in $L^2(\mu)$. This gives

$$\begin{aligned} & \int_{\mathbb{R}} x^\ell \psi(x) [\overline{\psi(x)} - \psi_k(x)] d\mu(x) \\ &= \int_{\mathbb{R}} x^\ell |\psi(x)|^2 d\mu(x) - \int_{\mathbb{R}} x^\ell \psi(x) \psi_k(x) d\mu(x) \\ & \rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned}$$

The terms $\psi(x) \psi_k(x)$ are linear combinations of ξ_α functions, so the second integral in the sum above is zero for all $k, \ell \in \mathbb{N}_0$. Therefore, for all $\ell \in \mathbb{N}_0$,

$$\int_{\mathbb{R}} x^\ell |\psi(x)|^2 d\mu(x) = 0.$$

In particular, the $\ell = 0$ case implies that $\psi = 0$. This contradicts our choice of ψ , hence we know there must be some function ξ_α which is not orthogonal to the polynomial space $\mathcal{P}$.

Because

$$\int_{\mathbb{R}} (x - \alpha) \overline{p(x)} \xi_\alpha(x) d\mu(x) = \int_{\mathbb{R}} \overline{p(x)} \psi(x) d\mu(x) = 0$$

for all $p \in \mathcal{P}$, we have

$$\langle M_x p | \xi_\alpha \rangle_{L^2(\mu)} = \alpha \langle p | \xi_\alpha \rangle_{L^2(\mu)},$$

where M_x is the multiplication operator by x . Every polynomial p is the image of a finite sequence $d \in \mathcal{D}$ under F . Recall that $FS = M_x F$ for all $d \in \mathcal{D}$, and let $\phi_\alpha = F^*\xi_\alpha \neq 0$. Applying F^* in the above equation gives

$$\langle F^* M_x p | F^* \xi_\alpha \rangle_{\mathcal{H}_Q} = \langle S d | \phi_\alpha \rangle_{\mathcal{H}_Q} = \alpha \langle d | \phi_\alpha \rangle_{\mathcal{H}_Q} = \alpha \langle F^* p | F^* \xi_\alpha \rangle_{\mathcal{H}_Q}.$$

Therefore, ϕ_α satisfies the equation $S^*\phi = \alpha\phi$, hence $\phi_\alpha \in \mathcal{L}(\alpha)$ for S which proves S is not self-adjoint. $\square$

The following example shows a Hankel matrix M which does not have a unique moment problem solution.

EXAMPLE 8.13. A non-unique measure.

Set

$$f(x) := \int_{\mathbb{R}} \cos(xt) e^{-(t^2+1/t^2)} dt.$$

Then

$$\int x^i f(x) dx = \left(\frac{d}{dt} \right)^i e^{-(t^2+1/t^2)} \Big|_{t=0} = 0 \text{ for all } i \in \mathbb{N}_0.$$

Let f_+ be the function $\max(f, 0)$ and let f_- be the function $-\min(f, 0)$. Set $f = f_+ - f_-$ and $d\mu_{\pm}(x) = f_{\pm}(x) dx$. Then

$$\int x^i d\mu_+(x) = \int x^i d\mu_-(x).$$

◇

Next, we see that the converse of Lemma 8.8 also holds.

THEOREM 8.14. *Let M be an infinite Hankel matrix with $M_{0,0} = 1$. The following statements are equivalent.*

- (1) *The moment problem $M = M^{(\mu)}$ does not have a unique solution.*
- (2) *Given any $\alpha \in \mathbb{C}$ with $0 < \text{Im}(\alpha) < 1$, there exists a vector $\xi \in \mathcal{H}_Q$ such that $H\xi = \lambda(1, \alpha, \alpha^2, \alpha^3, \dots)$ for some $\lambda \in \mathbb{C}$.*

PROOF. Let $\mathcal{H}_Q$ be the Hilbert space completion of the quadratic form Q_M as defined previously, and let S be the closure of the shift operator on $\mathcal{H}_Q$. The solution to the moment problem is unique if and only if the shift operator has no self-adjoint extensions, by Theorem 8.12. Using Lemma 8.8, this is true if and only if S is self-adjoint, i.e. its deficiency indices are $(0, 0)$.

(1 $\Rightarrow$ 2): This is a restatement of Lemma 8.8, using α for the deficiency spaces instead of i .

(2 $\Rightarrow$ 1): Fix $\alpha \in \mathbb{C}$ such that $0 < \text{Im}(\alpha) < 1$ and denote

$$s = (1, \alpha, \alpha^2, \alpha^3, \dots) \in \mathcal{H}_Q.$$

Assume there exists a nonzero $\xi \in \mathcal{H}_Q$ such that $H\xi = \lambda s$ for some nonzero scalar $\lambda \in \mathbb{C}$. Then for e_k a standard basis vector,

$$\langle Se_k | \xi \rangle_{\mathcal{H}_Q} = \langle Se_k | H\xi \rangle_{\ell^2} = \lambda \alpha^{k+1},$$

where we have selected α so that $H\xi$ is in ℓ^2 as well as in $\mathcal{H}_Q$. We then compute

$$\langle e_k | \xi \rangle_{\mathcal{H}_Q} = \langle e_k | H\xi \rangle_{\ell^2} = \lambda \alpha^k.$$

By linearity, we have for every $d \in \mathcal{D}$,

$$\langle Sd | \xi \rangle_{\mathcal{H}_Q} = \alpha \langle d | \xi \rangle_{\mathcal{H}_Q},$$

hence $\xi \in \mathcal{L}(\alpha)$ and we know the moment problem for μ does not have a unique solution. $\square$

8.4. Jacobi representations of matrices

The big picture in our work is an analysis of measures, passing from moments to spectra. It turns out that a number of our problems may be studied with the use of unbounded operators in Hilbert space, which fits in with our multi-faceted operator-theoretic approach to moment problems.

In this section we focus on the special relationship between banded matrices which represent unbounded operators and their associated moment problems. Beginning with a Hankel matrix M , we find a (nonunique) banded Jacobi matrix T

which encodes the information in M (Theorem 8.23). On the other hand, given a banded matrix T , we can find an associated moment matrix M (Theorem 8.25). We conclude with a discussion of higher-dimensional analogues and banded matrices which arise in quantum mechanics.

DEFINITION 8.15. We say that a matrix T indexed by $\mathbb{N}_0 \times \mathbb{N}_0$ is *banded* if its nonzero entries are restricted to the main diagonal and some number of diagonal bands adjacent to the main diagonal. Specifically, T is banded if there exist b_1, b_2 such that for all $j, k \in \mathbb{N}_0$,

$$T_{j,k} \neq 0 \Rightarrow j - b_1 \leq k \leq j + b_2.$$

When T_1 and T_2 are banded matrices, then

$$(T_1 T_2)_{s,t} := \sum_{n \in S} (T_1)_{s,n} (T_2)_{n,t}$$

for each pair $(s, t) \in \mathbb{N}_0 \times \mathbb{N}_0$. It is not hard to show that the banded matrices form an algebra with composition as the multiplication operation.

Recall that a matrix A with real entries is called *symmetric* if $A = A^{\text{tr}}$ and if A has complex entries, it is called *hermitian* if $A = \overline{A}^{\text{tr}}$. Banded hermitian (or symmetric) matrices $T : \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{C}$ define symmetric operators on $\ell^2(\mathbb{N}_0)$ with $\mathcal{D}$ as dense domain.

EXAMPLE 8.16. The matrix P which represents the momentum operator in quantum mechanics is a banded symmetric matrix:

$$(8.11) \quad P = \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 & 0 & & & \\ 1 & 0 & \sqrt{2} & 0 & & & \\ 0 & \sqrt{2} & 0 & \sqrt{3} & & & \\ & & \ddots & \ddots & \ddots & & \\ & & & \sqrt{n-2} & 0 & \sqrt{n-1} & \\ & & & 0 & \sqrt{n-1} & 0 & \sqrt{n} \\ & & & & \ddots & \ddots & \ddots \end{bmatrix}.$$

Given $v \in \ell^2$, we have

$$(Pv)_n = \frac{1}{2}(\sqrt{n-1}v_{n-1} + \sqrt{n}v_{n+1}).$$

◇

The following definition generalizes to higher dimensions, but for clarity we will state everything in a one-dimensional form. We begin with a Hankel matrix M , and again work in the Hilbert space $\mathcal{H}_Q$. We denote the standard orthonormal basis in $\ell^2(\mathbb{N}_0)$ by $\{\delta_n\}_{n \in \mathbb{N}_0}$. Even though $\{\delta_n\}_{n \in \mathbb{N}_0}$ is not an ONB in $\mathcal{H}_Q$, we also use δ_n to denote the element of $\mathcal{H}_Q$ with 0's in every place except the $(n+1)^{\text{st}}$, which contains a 1.

DEFINITION 8.17. We say that a Hankel matrix M is of *T-class* if there exists some banded hermitian matrix T , representing a symmetric operator, such that

$$(8.12) \quad M_{j,k} = \langle \delta_0 | T^{j+k} \delta_0 \rangle_{\ell^2}.$$

Recall from Proposition 8.11 that if M is Hankel and positive semidefinite, then M satisfies the equation

$$M_{j,k} = \langle \delta_0 | \tilde{S}^{j+k} \delta_0 \rangle_{\mathcal{H}_Q},$$

where $\tilde{S}$ is a self-adjoint extension of the shift operator on the Hilbert space $\mathcal{H}_Q$. We will demonstrate the correspondence between this shift operator S (when a moment matrix M is given) and a symmetric Jacobi matrix T .

LEMMA 8.18. *Every Hankel matrix of T -class is positive semidefinite.*

PROOF. Let $v \in \mathcal{D}$. Then because T is banded, all summations are finite:

$$\begin{aligned} \sum_j \sum_k \overline{v_j} M_{j,k} v_k &= \sum_j \sum_k \overline{v_j} \left\langle \delta_0 | T^{j+k} \delta_0 \right\rangle_{\ell^2} v_k \\ (8.13) \quad &= \left\langle \sum_j v_j T^j \delta_0 \middle| \sum_k v_k T^k \delta_0 \right\rangle_{\ell^2} \\ &= \left\| \sum_j v_j T^j \delta_0 \right\|_{\ell^2}^2 \geq 0. \end{aligned}$$

□

LEMMA 8.19 (One-dimensional version). *Suppose M is Hankel of T -class. Define $W : \mathcal{H}_Q \rightarrow \ell^2(\mathbb{N}_0)$ by*

$$W(\delta_k) := T^k \delta_0 \quad \forall k \in \mathbb{N}_0.$$

Then W is an isometry.

PROOF. We first show that $\|W(\delta_k)\|_{\ell^2}^2 = \|\delta_k\|_{\mathcal{H}_Q}^2$. Recall from the norm on $\mathcal{H}_Q$ that

$$\|\delta_k\|_{\mathcal{H}_Q}^2 = Q_M(\delta_k),$$

and

$$Q_M(\delta_k) = M_{k,k} = \langle \delta_0 | T^{k+k} \delta_0 \rangle_{\ell^2}.$$

On the other hand,

$$\|W(\delta_k)\|_{\ell^2}^2 = \langle T^k \delta_0 | T^k \delta_0 \rangle_{\ell^2} = \langle \delta_0 | T^{k+k} \delta_0 \rangle_{\ell^2}.$$

Since every element v of $\mathcal{D}$ is a finite linear combination of δ_k s, the rest of the proof follows from the same computation used in the proof of Lemma 8.18. □

As in Equation (8.1), let S will refer to the closure of the shift operator, which is defined on $\mathcal{D}$ by

$$S(c_0, c_1, c_2, \dots) := (0, c_0, c_1, c_2, \dots).$$

Because we will consider deficiency indices of two different operators S and T , we let $\mathcal{L}_{\pm}(S)$ denote the dimension of the deficiency subspace of S , and we let $\mathcal{L}_{\pm}(T)$ denote the dimension of the deficiency subspace of T .

LEMMA 8.20. *Let M be of T -class and let the Hilbert space $\mathcal{H}_Q$ be as defined above. On the dense domain $\mathcal{D} \subset \mathcal{H}_Q$, $WS = TW$.*

TABLE 1. Relationships among S , T , and W .

$$\begin{array}{ccc}
\mathcal{H}_{\min} & \xrightarrow{W} & \ell^2(\mathbb{N}_0) \\
S \curvearrowright & & \curvearrowright T \\
\mathcal{H}_{\min} & \xrightarrow{W} & \ell^2(\mathbb{N}_0)
\end{array}$$

PROOF. Let $c = (c_0, c_1, \dots) \in \mathcal{D}$. Then

$$\begin{aligned}
(8.14) \quad WS(c) &= W(0, c_0, c_1, \dots) \\
&= c_0 T \delta_0 + c_1 T^2 \delta_0 + c_2 T^3 \delta_0 + \dots \\
&= T(c_0 \delta_0 + c_1 T \delta_0 + c_2 T^2 \delta_0 + \dots) = TW(c).
\end{aligned}$$

□

An immediate result of Lemma 8.20 is the following:

LEMMA 8.21. *Suppose the set $\{Wc\}_{c \in \mathcal{D}}$ is dense in $\text{dom}(T)$. Then the isometry W maps the S -defect subspaces $\mathcal{L}_{\pm}(S)$ into the T -defect subspaces $\mathcal{L}_{\pm}(T)$.*

PROOF. Suppose $f_{\pm}$ denotes an element of $\mathcal{L}_{\pm}(S)$. We know that $S^* f_{\pm} = \pm i f_{\pm}$ if and only if $\langle Sc | f_{\pm} \rangle_{\mathcal{H}_Q} = \pm i \langle c | f_{\pm} \rangle_{\mathcal{H}_Q}$ for all $c \in \mathcal{D}$. With the assumption of density, $T^* W f_{\pm} = \pm i W f_{\pm}$ if and only if

$$\langle TWc | W f_{\pm} \rangle_{\ell^2} = \pm i \langle Wc | W f_{\pm} \rangle_{\ell^2}$$

for all $c \in \mathcal{F}$.

Let $c \in \mathcal{D}$. Then by Lemmas 8.20 and 8.19,

$$\begin{aligned}
(8.15) \quad \langle TWc | W f_{\pm} \rangle_{\ell^2} &= \langle WSc | W f_{\pm} \rangle_{\ell^2} = \langle Sc | f_{\pm} \rangle_{\mathcal{H}_Q} \\
&= \langle c | S^* f_{\pm} \rangle_{\mathcal{H}_Q} = \pm i \langle c | f_{\pm} \rangle_{\mathcal{H}_Q} = \pm i \langle Wc | W f_{\pm} \rangle_{\ell^2}.
\end{aligned}$$

□

Recall that we established in Lemma 8.8 that the deficiency indices of S are either $(0, 0)$ or $(1, 1)$.

COROLLARY 8.22. *Suppose the set $\{Wc \mid c \in \mathcal{F}\}$ is dense in $\text{dom}(T)$. Then*

$$\mathcal{L}_{\pm}(T) \geq \mathcal{L}_{\pm}(S).$$

Table 1 shows the relationships among S , T , and W .

Because

$$(8.16) \quad TW = WS,$$

when $S(c_0, c_1, \dots) := (0, c_0, c_1, \dots)$ in $\mathcal{H}_{\min}$, we can take adjoints in (8.16) to see that

$$(8.17) \quad W^* T^* = S^* W^*.$$

Finally, W^* maps $\mathcal{L}_{\pm}(T)$ into $\mathcal{L}_{\pm}(S)$. To see this, suppose $T^* f_{\pm} = \pm i f_{\pm}$. Apply W^* and use (8.17):

$$W^* T^* f_{\pm} = \pm i W^* f_{\pm} = S^* W^* f_{\pm},$$

which implies that $W^* f_{\pm} \in \mathcal{L}_{\pm}(S)$.

We now come to the main result of this section: every positive definite Hankel matrix M is of T -class, and the matrix T such that

$$M_{j,k} = \langle \delta_0 | T^{j+k} \delta_0 \rangle_{\ell^2}$$

can be chosen to be banded with respect to the canonical ONB in $\ell^2(\mathbb{N}_0)$.

THEOREM 8.23. *Let M be a positive definite Hankel matrix, where M is normalized so that $M_{0,0} = 1$.*

- (a) *Then there exists a banded symmetric matrix T operating on $\ell^2(\mathbb{N}_0)$ such that (8.12) is satisfied.*
- (b) *We may choose T of the banded form*

$$(8.18) \quad T = \begin{bmatrix} b_0 & a_0 & 0 & 0 & 0 & \cdots \\ \overline{a_0} & b_1 & a_1 & 0 & 0 & \cdots \\ 0 & \overline{a_1} & b_2 & a_2 & 0 & \\ 0 & 0 & \overline{a_2} & b_3 & a_3 & \\ \vdots & \vdots & & & & \ddots \end{bmatrix}.$$

PROOF. By Theorem 2.2 there exists μ such that

$$M_{j,k} = \int_{\mathbb{R}} x^{j+k} d\mu(x).$$

Now select the orthogonal polynomials $p_0(x), p_1(x), p_2(x), \dots$ in $L^2(\mu)$ with $p_0(x) = 1$. Then the mapping

$$(8.19) \quad \sum_{k=0}^{\infty} c_k p_k(x) \mapsto \{c_k\}_{k \in \mathbb{N}_0}$$

is an isometry of a subspace in $L^2(\mu)$ onto $\ell^2(\mathbb{N}_0)$. Specifically,

$$(8.20) \quad \left\| \sum_k c_k p_k \right\|_{L^2(\mu)}^2 = \sum_k |c_k|^2.$$

It is well-known (see, for example [Akh65, AAR99]) that there exist sequences $\{a_n\}_{n \in \mathbb{N}_0}$ and $\{b_n\}_{n \in \mathbb{N}_0}$ such that the following three-term recursion formulas are satisfied:

$$\begin{aligned} xp_0 &= b_0 p_0 + \overline{a_0} p_1 \\ xp_1 &= a_0 p_0 + b_1 p_1 + \overline{a_1} p_2 \\ &\vdots \\ xp_j &= a_{j-1} p_{j-1} + b_j p_j + \overline{a_j} p_{j+1} \\ &\vdots \end{aligned}$$

Because of the Hankel property assumed for M , in proving (8.12), it is enough to show that

$$(8.21) \quad M_{0,k} = \langle \delta_0 | T^k \delta_0 \rangle_{\ell^2}.$$

Using (8.19) and (8.20), we have for all $k \in \mathbb{N}_0$,

$$\begin{aligned} \langle \delta_0 | T^k \delta_0 \rangle_{\ell^2} &= \int_{\mathbb{R}} p_0 x^k p_0 d\mu(x) \\ &= \int_{\mathbb{R}} x^k d\mu(x) = M_{0,k}, \end{aligned}$$

which is the desired conclusion. □

REMARK 8.24. There are many other choices of banded symmetric or hermitian matrices which solve (8.12). The candidates for T are dictated by applications.

Consider a fixed positive definite Hankel matrix M such that the associated symmetric operator S has deficiency indices $(1, 1)$. Then by Theorem 8.12, there is a one-parameter family of inequivalent measures $\{\mu_z : |z| = 1\}$ such that $M^{(\mu_z)} = M$. Further, for each measure μ_z , we may compute an associated symmetric Jacobi matrix T_z , as in Theorem 8.23. The question we ask next is whether the Jacobi matrix T_z depends on the measure μ_z used to compute the orthogonal polynomials.

Consider the Hilbert space $\mathcal{H}_Q$, in which the shift operator S has dense domain. The Jacobi matrix T_μ we just computed is also a symmetric operator, this time with dense domain in $\ell^2(\mathbb{N}_0)$. Moreover, for each μ solving the M -moment problem, we have an isometry $F = F_\mu$ which maps $\mathcal{H}_Q$ into $L^2(\mu)$ and which intertwines S with multiplication by x (Lemma 8.2). Recall also that T_μ encodes multiplication by x .

By looking at the relevant formulas for the orthogonal polynomials $\{p_n\} \subset L^2(\mu_z)$, we see that T_z does not depend on z . In particular, we use the following well-known formulas for the orthogonal polynomials to justify our answer [Akh65].

Define

$$D_k = \det \begin{bmatrix} m_0 & \dots & m_k \\ m_1 & \dots & m_{k+1} \\ \vdots & & \vdots \\ m_k & \dots & m_{2k} \end{bmatrix}$$

and let $S_M = \{k \in \mathbb{N} : D_k \neq 0\}$. Then if $k \in S_M$, then $p_k(x) = (D_{k-1}D_k)^{-1/2}D_k(x)$, where

$$D_k(x) = \det \begin{bmatrix} m_0 & m_1 & \dots & m_k \\ m_1 & m_2 & \dots & m_{k+1} \\ \vdots & & & \vdots \\ m_{k-1} & m_k & \dots & m_{2k-1} \\ 1 & x & \dots & x^k \end{bmatrix}.$$

As a result, the isometry F_μ is also independent of μ ; the closed subspace spanned by the polynomials in $L^2(\mu_z)$ is independent of z , and it is only the relative orthogonal complement in $L^2(\mu_z)$ which depends on z . The orthogonal complement can be described by

$$\left\{ \psi \in L^2(\mu) \mid \int \psi(x)x^k d\mu(x) = 0 \text{ for all } k \in \mathbb{N}_0 \right\}.$$

Lemma 8.2 tells us that S and T are unitarily equivalent. Moreover,

$$\langle e_0 | S^k e_0 \rangle_{\mathcal{H}_Q} = \langle \delta_0 | T^k \delta_0 \rangle_{\ell^2},$$

so the spectral measures derived from the self-adjoint extensions $\tilde{S}$ of S and $\tilde{T}$ of T produce the same measures μ which solve the M -moment problem. Thus, if we are thus given the Jacobi matrix T , the self-adjoint extensions of T in $\ell^2(\mathbb{N}_0)$ correspond to spectral measures μ which will have a common moment matrix M .

We summarize with a theorem:

THEOREM 8.25. *Let M be a positive definite Hankel matrix, and let μ be a measure such that $M^{(\mu)} = M$. The associated Jacobi matrix T is independent of the choice of measure μ which solves the M -moment problem. Conversely, every symmetric Jacobi matrix T gives rise to a moment problem.*

The following three conditions are equivalent:

- (a) The solution μ to the M -moment problem is unique.
- (b) The deficiency indices for T are $(0, 0)$.
- (c) The polynomials are dense in $L^2(\mu)$.

PROOF. The forward direction is shown in the discussion preceding the statement of the theorem. Given a symmetric Jacobi matrix T , define a Hankel matrix M by taking

$$M_{j,k} = \langle \delta_0 | T^{j+k} \delta_0 \rangle_{\ell^2}.$$

Since powers of banded matrices remain banded, the entries of M are all finite. By Lemma 8.18, M is a positive-definite Hankel matrix, hence yielding a moment problem $M = M^{(\mu)}$ which has at least one solution.

The equivalent statements follow from Theorem 8.12, Corollary 8.22, and the discussion above showing that $\mathcal{L}_{\pm}(T) \leq \mathcal{L}_{\pm}(S)$. $\square$

8.5. The triple recursion relation and extensions to higher dimensions

Let $p_0(x) \equiv 1, p_1(x), p_2(x), \dots$ be the orthogonal polynomials with respect to μ , where $\deg(p_k) = k$. (Here we assume that μ corresponds to a positive definite, not positive semidefinite, linear functional on $L^2(\mu)$.) We can use Gram-Schmidt on $\{1, x, x^2, \dots\}$, so that

$$\text{span}\{1, x, x^2, \dots\} = \text{span}\{p_0(x), p_1(x), p_2(x), \dots\}$$

and

$$\int p_j(x) p_k(x) d\mu(x) = \delta_{j,k} \text{ (the Kronecker delta).}$$

Then for $k \in \mathbb{N}_0$ we have

$$(8.22) \quad xp_{n-1}(x) = b_{n-1}p_{n-2}(x) + a_{n-1}p_{n-1}(x) + \overline{b_n}p_n(x)$$

and

$$(8.23) \quad xp_n(x) = b_n p_{n-1}(x) + a_n p_n(x) + \overline{b_{n+1}} p_{n+1}(x).$$

Note that the $\overline{b_n}$ appearing in (8.22) is the conjugate of the b_n appearing in (8.23). To see this that this is true, suppose

$$\begin{aligned} xp_n(x) &= Bp_{n-1}(x) + \text{terms in } p_n(x) \text{ and } p_{n+1}(x) \\ xp_{n-1}(x) &= Cp_n(x) + \text{terms in } p_{n-2}(x) \text{ and } p_{n-1}(x). \end{aligned}$$

We now take advantage of the orthogonality:

$$\begin{aligned} B &= \langle p_{n-1}(x) | xp_n(x) \rangle_{L^2(\mu)} = \langle xp_{n-1}(x) | p_n(x) \rangle_{L^2(\mu)} \\ &= \langle Cp_n(x) | p_n(x) \rangle_{L^2(\mu)} = \overline{C} \langle p_n(x) | p_n(x) \rangle_{L^2(\mu)} = \overline{C}. \end{aligned}$$

The triple recursion relation can be considered in terms of projections onto finite-dimensional subspaces. Set

$$\mathcal{H}_n := \text{span}\{1, x, \dots, x^n\} = \text{span}\{p_0(x), p_1(x), \dots, p_n(x)\}$$

and let Q_n be the projection onto $\mathcal{H}_n$ with

$$Q_n^* = Q_n = Q_n^2$$

and

$$Q_n(L^2(\mu)) = \mathcal{H}_n.$$

(Note: Q_n is a projection, and is *not* related to our earlier quadratic form.) We are interested in the shift operator, which is the same as multiplication by x on $\mathcal{P} \subset L^2(\mu)$. We have

$$x\mathcal{H}_n \subset \mathcal{H}_{n+1},$$

so

$$Q_{n+1}xQ_n = xQ_n.$$

Set $Q_n^\perp := I - Q_n$ —that is, $Q_n^\perp$ is the projection onto $\mathcal{H}_n^\perp$, where $\mathcal{H}_n^\perp = L^2(\mu) \ominus \mathcal{H}_n$.

If $k < n - 1$, then $\langle p_k, xp_n \rangle_{L^2(\mu)} = 0$. To see this, write

$$\langle p_k, xp_n \rangle_{L^2(\mu)} = \langle xp_k, p_n \rangle_{L^2(\mu)},$$

and p_n is orthogonal to any polynomial with degree less than n .

Now,

$$\begin{aligned} xp_n &= Q_n(xp_n) + Q_n^\perp(xp_n) \\ &= \underbrace{b_n p_{n-1} + a_n p_n}_{Q_n(xp_n)} + \underbrace{\overline{b_{n+1} p_{n+1}}}_{Q_n^\perp(xp_n)}. \end{aligned}$$

If we restrict J^* to $\mathcal{P}$, the subspace of all polynomials, then

$$J^*|_{\mathcal{P}} = J,$$

since $J \subset J^*$. Moreover, since the projections Q_n are self-adjoint,

$$Q_{n+1}JQ_n = JQ_n \Rightarrow Q_nJQ_{n+1} = Q_nJ.$$

Finally,

$$Q_{n-2}J(Q_n - Q_{n-1}) = 0,$$

since

$$Q_{n-2}(JQ_n) = Q_{n-2}JQ_{n-1}Q_n = Q_{n-2}JQ_{n-1}.$$

The projection approach to the recursion relation can be extended to $\mathbb{R}^d$ for $d > 1$. For each of the d coordinate directions, there is a Jacobi matrix

$$J_k = \begin{bmatrix} \frac{a_0^{(k)}}{b_1^{(k)}} & b_1^{(k)} & 0 & 0 & 0 & \cdots \\ b_1^{(k)} & \frac{a_1^{(k)}}{b_2^{(k)}} & b_2^{(k)} & 0 & 0 & \cdots \\ 0 & \frac{a_2^{(k)}}{b_3^{(k)}} & \frac{a_3^{(k)}}{b_4^{(k)}} & b_3^{(k)} & 0 & \cdots \\ 0 & 0 & b_3^{(k)} & \frac{a_4^{(k)}}{b_4^{(k)}} & b_4^{(k)} & \\ 0 & 0 & 0 & b_4^{(k)} & a_4^{(k)} & \\ \vdots & \vdots & \vdots & & & \ddots \end{bmatrix}$$


and a shift operator S_k , which is realized by multiplication in the k^{th} coordinate

$$S_k p(x_1, \dots, x_d) = x_k p(x_1, \dots, x_d).$$

Recall that the degree of $x^\alpha = x_1^{\alpha_1} \cdots x_d^{\alpha_d}$ is $\alpha_1 + \dots + \alpha_d$. With this notation, the finite-dimensional subspaces are

$$\mathcal{H}_n = \{p \in \mathcal{P} | \deg(p) \leq n\}.$$

TABLE 2. Two approaches to moments and banded matrices.



8.6. Concrete Jacobi matrices

We examine the momentum and position operators from quantum mechanics for one degree of freedom and then study some Hamiltonian operators (for example, the polynomials in the momentum and position operators in Table 3). The operators we consider all have a common property: in a natural orthonormal basis their representations take the form of infinite banded matrices; that is, the matrices have zeros outside a band around the diagonal of finite width.

The banded property of the matrices makes matrix multiplication easy; under multiplication, the banded matrices form an algebra of unbounded operators. While such banded matrices follow simple algebraic rules, their spectral theory can be subtle. For example, we show that these operators may not have a well-defined spectral resolution. Using von Neumann's deficiency indices, we showed above the connection of Jacobi matrices to the theory of extensions of symmetric operators with dense domain, and thereby to moment problems.

One often encounters problems in physics, such as the Heisenberg banded matrices T , where the nature of the bands is dictated by applications. A particular infinite matrix T represents an operator in an ℓ^2 sequence space. In fact, in a particular application, T may be realized in a different Hilbert space, for example an L^2 function space, but the function version will be unitarily equivalent to the matrix model. In particular, we refer to the fact that Heisenberg's matrix formulation of quantum mechanics is unitarily equivalent to Schrödinger's wave formulation in function space. For example, the momentum operator P is represented by the matrix (8.11) in ℓ^2 and by the operator $\frac{1}{i} \frac{d}{dx}$ on a dense subspace of $L^2(\mathbb{R})$. See Examples 8.16 and 8.26 and the remark following the two examples.

From our banded matrix T we then get a Hankel matrix M , and we apply our theory to M . In particular, we find the measures μ which solve the moment problem for M . We use operator theory in constructing the family of measures μ which solve the moment problem at hand.

We emphasize further that in applications, one typically encounters a much richer family of banded symmetric or hermitian infinite matrices T ; for example those from Heisenberg's quantum mechanics. In these matrices, the band-size will typically be more than three. In fact the band can be any size, and the deficiency indices can be anything. However this wider class of banded matrices, including for example anharmonic oscillators, may be studied with the aid of the associated Jacobi matrices.

We begin with two Jacobi matrices, the matrix P given in Example 8.16 and Q here.

EXAMPLE 8.26. The operator Q is represented by multiplication by x on $L^2(\mathbb{R})$. The operator Q can be represented on ℓ^2 by a matrix defined by

$$(Qv)_n = \frac{1}{2i}(\sqrt{n-1}v_{n-1} - \sqrt{n}v_{n+1}).$$

REMARK 8.27. The two operators P and Q in Examples 8.16 and 8.26 have dense domains in $L^2(\mathbb{R})$:

$$\text{dom}(P) = \{f \in L^2(\mathbb{R}) : f' \in L^2(\mathbb{R})\}$$

and

$$\text{dom}(Q) = \{f \in L^2(\mathbb{R}) : xf(x) \in L^2(\mathbb{R})\}.$$

The operators P and Q are both self-adjoint, and they are unitarily equivalent via the Fourier transform in $L^2(\mathbb{R})$.

Setting

$$A_{\pm} := \frac{1}{\sqrt{2}}(P \pm iQ),$$

we get $A_+^* = A_-$, and the commutator

$$(8.24) \quad [A_+, A_-] = -I.$$

The Hermite function $h_0(x) = c_0 e^{-x^2/2}$ satisfies

$$A_- h_0 = 0.$$

An application of Equation (8.24) yields

$$(A_+ A_-) A_+^n h_0 = n A_+^n h_0 \text{ for each } n \in \mathbb{N}.$$

The functions $h_n := c_n A_+^n h_0$ diagonalize the harmonic oscillator Hamiltonian

$$H := A_+ A_- = \frac{1}{2}(P^2 + Q^2 - I),$$

and the constants c_n can be chosen so that $\{h_n | n \in \mathbb{N}_0\}$ is an orthonormal basis in $L^2(\mathbb{R})$ consisting of Hermite functions.

Using this ONB, we arrive at the two matrix representations for P and Q in Examples 8.16 and 8.26. Specifically,

$$\begin{aligned} \langle h_{n-1} | P h_n \rangle &= \frac{1}{2} \sqrt{n}, \\ \langle h_n | P h_n \rangle &= 0, \end{aligned}$$

and

$$\langle h_{n+1} | P h_n \rangle = \frac{1}{2} \sqrt{n+1},$$

which is the Jacobi matrix in (8.11).

EXAMPLE 8.28. If $T = QPQ$ in $\ell^2(\mathbb{N}_0)$, then T has deficiency indices both equal to 1.

PROOF. A differential equations problem. □

TABLE 3. Some polynomials in the position and momentum operators.

T	index	spectrum
QPQ	$(1, 1)$	depends on the choice of selfadjoint extension
$P^2 + Q^4$	$(0, 0)$	discrete, anharmonic oscillator
$P^2 - Q^4$	$(2, 2)$	repulsive potential quantum particle shoots to infinity in finite time
$P^2 + Q^2$	$(0, 0)$	$\{2n + 1 \mid n \in \mathbb{N}_0\}$
$P^2 - Q^2$	$(0, 0)$	continuous $\mathbb{R}$ (easier to see in $L^2(\mathbb{R})$ than by using matrix calculations)

Bibliography

- [AAR99] George E. Andrews, Richard Askey, and Ranjan Roy, *Special functions*, Encyclopedia of Mathematics and its Applications, vol. 71, Cambridge University Press, Cambridge, 1999. MR MR1688958 (2000g:33001)
- [Akh65] N. I. Akhiezer, *The classical moment problem and some related questions in analysis*, Translated by N. Kemmer, Hafner Publishing Co., New York, 1965. MR MR0184042 (32 #1518)
- [Aro50] N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337–404. MR MR0051437 (14,479c)
- [Bag92] Lawrence Baggett, *Functional analysis: A primer*, Pure and Applied Mathematics, vol. 153, Marcel Dekker, Inc., New York, NY, 1992.
- [BCR84] Christian Berg, Jens Peter Reus Christensen, and Paul Ressel, *Harmonic analysis on semigroups*, Graduate Texts in Mathematics, vol. 100, Springer-Verlag, New York, 1984, Theory of positive definite and related functions. MR MR747302 (86b:43001)
- [BD06] Yuriy M. Berezansky and Mykola E. Dudkin, *The complex moment problem and direct and inverse spectral problems for the block Jacobi type bounded normal matrices*, Methods Funct. Anal. Topology **12** (2006), no. 1, 1–31.
- [BD07] ———, *On the complex moment problem*, Math. Nachr. **280** (2007), no. 1-2, 60–73.
- [BS52] S. Bergman and M. Schiffer, *Theory of kernel functions in conformal mapping*, Construction and applications of conformal maps. Proceedings of a symposium (Washington, D. C.), National Bureau of Standards, Appl. Math. Ser., No. 18, U. S. Government Printing Office, 1952, pp. 199–206. MR MR0054038 (14,860c)
- [Con90] John B. Conway, *A course in functional analysis*, second ed., Graduate Texts in Mathematics, vol. 96, Springer-Verlag, New York, 1990. MR MR1070713 (91e:46001)
- [Dei99] P. A. Deift, *Orthogonal polynomials and random matrices: a Riemann-Hilbert approach*, Courant Lecture Notes in Mathematics, vol. 3, New York University Courant Institute of Mathematical Sciences, New York, 1999.
- [DJ06] Dorin Ervin Dutkay and Palle E. T. Jorgensen, *Methods from multiscale theory and wavelets applied to nonlinear dynamics*, Wavelets, multiscale systems and hypercomplex analysis, Oper. Theory Adv. Appl., vol. 167, Birkhäuser, Basel, 2006, pp. 87–126. MR MR2240292
- [DJ07] ———, *Harmonic analysis and dynamics for affine iterated function systems*, Houston J. Math. **33** (2007), no. 3, 877–905. MR MR2335741
- [DS88] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part II*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988, Spectral theory. Selfadjoint operators in Hilbert space, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1963 original, A Wiley-Interscience Publication. MR MR1009163 (90g:47001b)
- [Dut04] Dorin Ervin Dutkay, *Positive definite maps, representations and frames*, Rev. Math. Phys. **16** (2004), no. 4, 451–477. MR MR2065233 (2005e:42115)
- [Erd39] Paul Erdős, *On a family of symmetric Bernoulli convolutions*, Amer. J. Math. **61** (1939), 974–976. MR MR0000311 (1,52a)
- [EST06] C. Escribano, M.A. Sastre, and E. Torrano, *Moment matrix of self-similar measures*, Electronic Transactions on Numerical Analysis **24** (2006), 79–87.
- [Fre07] Christopher French, *Transformations preserving the Hankel transform*, J. Integer Seq. **10** (2007), no. 7, Article 07.7.3, 14. MR MR2322498 (2008e:11029)
- [Fug83] Bent Fuglede, *The multidimensional moment problem*, Exposition. Math. **1** (1983), no. 1, 47–65.

- [Hal67] Paul R. Halmos, *A Hilbert space problem book*, D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967. MR MR0208368 (34 #8178)
- [Hav35] E. K. Haviland, *On the Momentum Problem for Distribution Functions in More than One Dimension*, Amer. J. Math. **57** (1935), no. 3, 562–568. MR MR1507095
- [Hav36] ———, *On the Momentum Problem for Distribution Functions in More Than One Dimension. II*, Amer. J. Math. **58** (1936), no. 1, 164–168. MR MR1507139
- [HR79] Edwin Hewitt and Kenneth A. Ross, *Abstract harmonic analysis. Vol. I*, second ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 115, Springer-Verlag, Berlin, 1979, Structure of topological groups, integration theory, group representations. MR MR551496 (81k:43001)
- [Hut81] John E. Hutchinson, *Fractals and self-similarity*, Indiana Univ. Math. J. **30** (1981), no. 5, 713–747. MR MR625600 (82h:49026)
- [JKS07a] Palle E. T. Jorgensen, Keri A. Kornelson, and Karen L. Shuman, *Affine systems: asymptotics at infinity for fractal measures*, Acta Appl. Math. **98** (2007), no. 3, 181–222. MR MR2338387 (2008i:42013)
- [JKS07b] Palle E. T. Jorgensen, Keri A. Kornelson, and Karen L. Shuman, *Harmonic analysis of iterated function systems with overlap*, Journal of Mathematical Physics **48** (2007), 083511, 35pp.
- [JKS08] ———, *Orthogonal exponentials and iterated function systems with overlap*, Representations, Wavelets, and Frames: A Celebration of the Mathematical Work of Lawrence W. Baggett, Applied and Numerical Harmonic Analysis, Birkhäuser, Boston, MA, 2008.
- [JÓ00] Palle E. T. Jorgensen and Gestur Ólafsson, *Unitary representations and Osterwalder-Schrader duality*, The mathematical legacy of Harish-Chandra (Baltimore, MD, 1998), Proc. Sympos. Pure Math., vol. 68, Amer. Math. Soc., Providence, RI, 2000, pp. 333–401. MR MR1767902 (2001f:22036)
- [Jor00] Palle E. T. Jorgensen, *Off-diagonal terms in symmetric operators*, J. Math. Phys. **41** (2000), no. 4, 2337–2349. MR MR1751892 (2001a:47025)
- [Jor06] ———, *Analysis and probability: wavelets, signals, fractals*, Graduate Texts in Mathematics, vol. 234, Springer, New York, 2006. MR MR2254502
- [Jor07] Palle E. T. Jorgensen, *Surprising similarities: Amazon review of “The Umbral Calculus” by Gian-Carlo Rota*, September 2007, http://www.amazon.com/review/R1FKW5NW6Y8ZKN/ref=cm_cr_rdp_perm/.
- [JP98] Palle E. T. Jorgensen and Steen Pedersen, *Dense analytic subspaces in fractal L^2 -spaces*, J. Anal. Math. **75** (1998), 185–228. MR MR1655831 (2000a:46045)
- [Kat80] Tosio Kato, *Perturbation theory for linear operators*, reprinted second ed., Springer-Verlag, Berlin, 1980, Grundlehren der Mathematischen Wissenschaften, Band 132. MR MR0407617 (53 #11389)
- [Kol50] A. N. Kolmogorov, *Foundations of the Theory of Probability*, Chelsea Publishing Company, New York, N. Y., 1950. MR MR0032961 (11,374a)
- [Lan87a] H. J. Landau, *Classical background of the moment problem*, Moments in mathematics (San Antonio, Tex., 1987), Proc. Sympos. Appl. Math., vol. 37, Amer. Math. Soc., Providence, RI, 1987, pp. 1–15.
- [Lan87b] ———, *Maximum entropy and the moment problem*, Bull. Amer. Math. Soc. (N.S.) **16** (1987), no. 1, 47–77.
- [Lay01] John W. Layman, *The Hankel transform and some of its properties*, J. Integer Seq. **4** (2001), no. 1, Article 01.1.5, 11 pp. (electronic). MR MR1848942 (2002k:05021)
- [Mag50] Wilhelm Magnus, *On the spectrum of Hilbert’s matrix*, Amer. J. Math. **72** (1950), 699–704. MR MR0041358 (12,836e)
- [PS72] K. R. Parthasarathy and K. Schmidt, *Positive definite kernels, continuous tensor products, and central limit theorems of probability theory*, Springer-Verlag, Berlin, 1972, Lecture Notes in Mathematics, Vol. 272. MR MR0622034 (58 #29849)
- [Rie23] M. Riesz, *Sur le problème des moments, troisième note*, Ark. Mat. Fys. Astr. **17** (1923), no. 16, 1–52.
- [Rom84] Steven Roman, *The umbral calculus*, Pure and Applied Mathematics, vol. 111, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1984. MR MR741185 (87c:05015)
- [RR78] Steven M. Roman and Gian-Carlo Rota, *The umbral calculus*, Advances in Math. **27** (1978), no. 2, 95–188. MR MR0485417 (58 #5256)

- [RS75] Michael Reed and Barry Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1975. MR MR0493420 (58 #12429b)
- [RS80] ———, *Methods of modern mathematical physics. I. Functional analysis*, Academic Press [Harcourt Brace & Company], San Diego CA, 1980.
- [Rud91] Walter Rudin, *Functional analysis*, second ed., International Series in Pure and Applied Mathematics, McGraw-Hill Inc., New York, 1991. MR MR1157815 (92k:46001)
- [Sho47] J. A. Shohat, *Orthogonal polynomials and polynomials in general*, Math. Mag. **21** (1947), 3–21.
- [SS06] Michael Z. Spivey and Laura L. Steil, *The k -binomial transforms and the Hankel transform*, J. Integer Seq. **9** (2006), no. 1, Article 06.1.1, 19 pp. (electronic). MR MR2175536 (2006m:11022)
- [ST43] J. A. Shohat and J. D. Tamarkin, *The Problem of Moments*, American Mathematical Society Mathematical surveys, vol. II, American Mathematical Society, New York, 1943.
- [Sze75] Gábor Szegő, *Orthogonal polynomials*, fourth ed., American Mathematical Society, Providence, R.I., 1975, American Mathematical Society, Colloquium Publications, Vol. XXIII. MR MR0372517 (51 #8724)
- [Wid66] Harold Widom, *Hankel matrices*, Trans. Amer. Math. Soc. **121** (1966), 1–35. MR MR0187099 (32 #4553)